



Company Name: APPARENT PTY. LTD.
Product Name: APPARENT SWORD 750 WG HERBICIDE
APVMA Approval No: 65285/103782



| | |
|--------------------------------|--|
| Label Name: | APPARENT SWORD 750 WG HERBICIDE |
| Signal Headings: | CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING |
| Constituent Statements: | ACTIVE CONSTITUENT: 750 g/kg CHLORSULFURON |
| Mode of Action: | GROUP B HERBICIDE |
| Statement of Claims: | A selective herbicide for the control of Annual (Wimmera) Ryegrass and certain broadleaf weeds in Wheat, Barley, Oats, Cereal Rye and Triticale. |
| Net Contents: | 1 kg 5 kg 10 kg |
| Restrains: | RESTRAINTS: DO NOT spray emerged crops if rain is expected within four hours. After mixing in the tank, spray within 48 hours if this product is used by itself, or within 24 hours if mixed with another product. DO NOT apply to plants suffering stress. |
| Directions for Use: | This section contains file attachment. File Name: DFU - Sword 750 - 0415.docx File Size: 30230 bytes |
| Other Limitations: | |

| | |
|-----------------------------|---|
| Withholding Periods: | WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED. |
|-----------------------------|---|

| | |
|----------------------|--|
| Trade Advice: | |
|----------------------|--|

| | |
|------------------------------|---|
| General Instructions: | This section contains file attachment. File Name: GI - Sword 750 WG -0415.docx File Size: 26623 bytes |
|------------------------------|---|

| | |
|----------------------------|--|
| Resistance Warning: | <p>RESISTANT WEEDS WARNING</p> <p>GROUP B HERBICIDE</p> <p>Apparent Sword 750 WG Herbicide is a member of the sulfonylurea group of herbicides. Apparent Sword 750 WG Herbicide has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action. For weed resistance management, Apparent Sword 750 WG Herbicide is a Group B herbicide. Naturally-occurring weed biotypes resistant to Apparent Sword 750 WG Herbicide and other ALS inhibitor herbicides (Annual Ryegrass and some broadleaf weeds) are known to exist. They can eventually dominate the weed population if these herbicides are used repeatedly. These herbicides will not be controlled by Apparent Sword 750 WG Herbicide or other ALS inhibitor herbicides. Annual Ryegrass biotypes resistant to diclofop-methyl and other "grass specific" herbicides are often also resistant to Apparent Sword 750 WG Herbicide. Before using Apparent Sword 750 WG Herbicide on a population resistant to "grass specific" herbicides, have a resistance test conducted to ensure that it is still susceptible to Apparent Sword 750 WG Herbicide.</p> <p>Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of Apparent Sword 750 WG Herbicide to control resistant weeds. To prevent, or at least minimise the risk of resistant weeds occurring, use Apparent Sword 750 WG Herbicide in tank mixes (if appropriate) and/or rotations with herbicides having different modes of action effective on the same weed species. Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors.</p> <p>DO NOT make more than one application of an ALS inhibitor herbicide to a crop, either pre-sowing incorporated by sowing or post crop and weed emergence. If the user suspects that an ALS inhibitor-resistant weed is present, Apparent Sword 750 WG Herbicide or other ALS inhibitor herbicides should not be used. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries.</p> |
|----------------------------|--|

| | |
|---------------------|--|
| Precautions: | |
|---------------------|--|

| | |
|---------------------|--|
| Protections: | <p>PROTECTION OF CROPS NATIVE AND OTHER NON-TARGET PLANTS DO NOT apply or drain or flush equipment on or near desirable trees or other plants or on areas where roots may extend or in locations where the chemical may be washed or moved into contact with their roots. DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pasture.</p> <p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p> |
|---------------------|--|

| | |
|------------------------------|--|
| Storage and Disposal: | <p>STORAGE AND DISPOSAL</p> <p>Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.</p> <p>Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> |
|------------------------------|--|

| | |
|---------------------------|--|
| Safety Directions: | <p>SAFETY DIRECTIONS</p> <p>Avoid contact with eyes and skin. DO NOT inhale spray mist. Wash hands after use.</p> |
|---------------------------|--|

| | |
|--------------------------------|---|
| First Aid Instructions: | <p>FIRST AID</p> <p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p> |
|--------------------------------|---|

| | |
|----------------------------|--|
| First Aid Warnings: | |
|----------------------------|--|

DIRECTIONS FOR USE

METHOD OF USE - PRE-SOWING INCORPORATED BY SOWING

Annual Ryegrass

| Crop Situation | Weeds Controlled | State(s) | Rate g/ha | | | Critical Comments |
|--------------------------|---|----------------------------|-----------------------|-----------|-------------|---|
| | | | Soil Type | | | |
| | | | Light to Medium Soils | | Heavy Soils | |
| | | | Soil pH | | | |
| | | | Less than 7 | 7.0 – 8.5 | 8.5 or less | |
| Wheat and Triticale only | Annual (Wimmera) Ryegrass <i>Lolium rigidum</i> | NSW, ACT, Vic, SA, WA only | 20 | 15 or 20* | 20 | * Use the higher rate when paddock history suggests a high weed population can be expected. NOTE: Refer to General Instructions for optimum application timing and conditions. |

| Crop Situation | Weeds Controlled | State(s) | Rate g/ha | Critical Comments |
|--|--|-------------------------------------|----------------------|---|
| Wheat and Triticale Only | African Turnip Weed <i>Sisymbrium thellungii</i> | NSW, ACT and Qld only | 20 | |
| | Amsinckia/Yellow Burrweed <i>Amsinckia</i> spp. | NSW, ACT, Vic, SA, WA only | 15 | |
| | Annual Phalaris <i>Phalaris paradoxa: Phalaris minor</i> | NSW, ACT only | 20 + 1 L Trifluralin | If possible spray and incorporate into the soil in one operation. If this is not possible incorporation should take place within four (4) hours of spraying. Delay may cause inferior weed control. Use only Trifluralin products with an active level of 400 g/L |
| | Barley grass <i>Hordeum leporinum</i> | NSW, ACT and Tas only | | |
| | Silvergrass <i>Vulpia</i> spp. | Tas only | | |
| | Ball Mustard <i>Neslia puniculata</i> | SA only | 15 | |
| | Black Bindweed/Climbing Buckwheat <i>Fallopia convolvulus</i> | Qld only | 20 | Apply to dry soil before the sowing rain. Mechanical incorporation before sowing rains is not necessary. |
| | Brome grass <i>Bromus</i> spp. (Suppression only) | NSW, ACT, Vic, SA, WA, Tas only | | Gives suppression only if populations are 20 plants/m ² or less. |
| | Cape Tulip <i>Homeria</i> spp. | WA only | | |
| | Capeweed <i>Arctotheca calendula</i> | NSW, ACT, Vic, SA, WA, Tas only | | On acid soils pH 5.5 or less, this product will give a shorter period of control in wet years. |
| | Charlock Sinapis arvensis | Vic, SA, Tas only | 15 | |
| | Common Iceplant <i>Mesembryanthemum crystallinum</i> | SA only | | |
| | Corn Gromwell/Sheepweed/ White Ironweed <i>Buglossoides arvensis</i> | Qld, NSW, Vic, SA, WA, ACT only | 20 | |
| | Deadnettle <i>Lamium amplexicaule</i> | All States | 15 or 20 | Use the higher rate when paddock history suggests a high weed population can be expected. |
| | Docks <i>Rumex</i> spp. | NSW, Vic, SA, WA, ACT, Tas only | 20 | |
| | Fat-hen <i>Chenopodium album</i> | NSW, Tas, ACT only | | |
| | Fumitory <i>Fumaria</i> spp. | NSW, Vic, SA, WA, ACT, Tas only | 15 or 20 | Use the higher rate when paddock history suggests a high weed population can be expected. |
| | Guildford grass/Onion grass <i>Romulea rosea</i> | WA only | 15 | |
| | Indian Hedge Mustard <i>Sisymbrium orientale</i> | All States | | |
| | King Island Melilot <i>Melilotus indicus</i> | Vic, SA only | | |
| | Lincoln weed <i>Diploaxis tenuifolia</i> | SA only | | |
| | Loosestrife <i>Lysimachia</i> spp. | Vic only | | |
| | Mintweed <i>Salvia reflexa</i> | Qld, ACT and NSW only | 20 | |
| | Mouse-ear Chickweed <i>Cerastium</i> spp. | NSW, Vic, SA, WA, ACT, Tas only | 15 | |
| | New Zealand Spinach <i>Tetragonia tetragonoides</i> | Qld only | 20 | |
| | Paradoxa grass <i>Phalaris paradoxa</i> | Nth NSW (Soil pH >7.5) and Qld only | | Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary. |
| | Paterson's Curse/Salvation Jane <i>Echium plantagineum</i> | NSW, Vic, SA, WA, ACT, Tas only | 15 | |
| | Pimpernels <i>Anagallis arvensis</i> | NSW, Vic, SA, ACT, Tas only | | |
| | Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i> | Vic, SA only | 20 | |
| | Rough Poppy <i>Papaver hybridum</i> | NSW, SA, WA, ACT, Tas only | 15 or 20 | Use the higher rate when paddock history suggests a high weed population can be expected. |
| Saffron Thistle <i>Carthamus lanatus</i> (Suppression only) | Qld, NSW, Vic, SA, Tas, ACT only | 20 | | |
| Saltbush <i>Atriplex muelleri</i> | Qld, ACT and NSW only | | | |

| Crop Situation | Weeds Controlled | State(s) | Rate g/ha | Critical Comments |
|---------------------------------|--|------------------------------------|-----------|--|
| Wheat and Triticale Only (cont) | Shepherd's Purse <i>Capsella bursa pastoris</i> | NSW, Vic, SA, WA, ACT, Tas only | 15 or 20 | Use the higher rate when paddock history suggests a high weed population can be expected. |
| | Slender Celery <i>Apium leptophyllum</i> | Qld, ACT and NSW only | 20 | |
| | Slender Thistle <i>Carduus tenuiflorus</i> | Tas only | | |
| | Soursob <i>Oxalis pes-caprae</i> | NSW, Vic, SA, ACT only | 15 | Apply only to soils of pH 7.5 or above. Apply after majority of Soursobs have emerged and leave soil undisturbed for 1-4 weeks prior to cultivation or sowing. The most effective and reliable control is achieved with early post-emergence applications (EPE) after crop and weed emergence. |
| | Spear Thistle <i>Cirsium vulgare</i> | Tas only | 20 | |
| | Stemless Thistle <i>Onopordum acaulon</i> | SA only | 15 or 20 | Use the higher rate when paddock history suggests a high weed population can be expected. |
| | Storksbill/Wild Geranium <i>Erodium</i> spp. | Vic, SA, WA, Tas only | 15 | |
| | Three-Cornered Jack(s)/Doublegee/ Spiny Emex <i>Emex australis</i> | NSW, Vic, SA, ACT, WA only | 20 | |
| | Tree Hogweed <i>Polygonum patulum</i> | Vic, SA only | | |
| | Turnip weed <i>Rapistrum rugosum</i> | Qld and SA only | 15 | |
| | Wireweed/Hogweed <i>Polygonum aviculare</i> | All States | 15 or 20 | |
| | Wild Turnip <i>Brassica tournefortii</i> | NSW, Vic, SA, WA, ACT and Tas only | 15 | Use the higher rate when paddock history suggests a high weed population can be expected. |

METHOD OF USE – POST CROP AND WEED EMERGENCE

Annual Ryegrass

| Crop Situation | Weeds Controlled | State(s) | Rate g/ha | | | Critical Comments |
|--|--|----------------------------|-----------------------|-----------|-------------|--|
| | | | Soil Type | | | |
| | | | Light to Medium Soils | | Heavy Soils | |
| | | | Soil pH | | | |
| | | | Less than 7 | 7.0 – 8.5 | 8.5 or less | |
| Wheat, Barley, Oats, Cereal Rye and Triticale only | Annual (Wimmera) Ryegrass <i>Lolium rigidum</i> | NSW, ACT, Vic, SA, WA only | 20 or 25* | 15 or 20* | 20 or 25* | * Use the higher rate under heavy weed pressure. Apply no later than the 3 leaf stage of Annual Ryegrass. * Application of this product to Annual Ryegrass 2 leaf or greater with water volumes less than 50 L/ha may result in reduced efficacy. |

| Crop Situation | Weeds Controlled | State(s) | Rate g/ha | Critical Comments |
|--|--|------------------------------------|-----------|---|
| Wheat, Barley, Oats, Cereal Rye and Triticale only | African Turnip Weed <i>Sisymbrium thellungii</i> | NSW, ACT and Qld only | 20 | Apply at cotyledon to 4 leaf stage. |
| | Amsinckia/Yellow Burrweed <i>Amsinckia</i> spp. | NSW, ACT, Vic, SA, WA only | 15 | |
| | Ball Mustard <i>Neslia puniculata</i> | SA only | | |
| | Bilora/Carrot Weed <i>Cotula australis</i> | | 25 | |
| | Black Bindweed/Climbing Buckwheat <i>Fallopia convolvulus</i> | Qld, NSW, ACT only | 20 | Apply at cotyledon to 2 leaf stage of weeds. |
| | Cape Tulip <i>Homeria</i> spp. | WA only | | |
| | Charlock <i>Sinapis arvensis</i> | NSW, ACT, Vic, SA, Tas only | 15 | |
| | Corn Gromwell/Sheepweed/White Ironweed <i>Buglossoides arvensis</i> | NSW, ACT, Vic, SA, WA only | 20 | Apply at cotyledon to 2 leaf stage. If applied at a later stage only suppression will occur. |
| | Deadnettle <i>Lamium amplexicaule</i> | Qld, NSW, ACT, Vic, SA, Tas only | 15 or 20 | Use the higher rate under heavy weed pressure. |
| | Docks <i>Rumex</i> spp. | Vic, SA, WA and Tas only | 15 | |
| | Fat-hen <i>Chenopodium album</i> | NSW, Tas, ACT only | 20 | |
| | Fumitory Densiflower <i>Fumaria densiflora</i> | NSW, ACT, Vic, SA, WA, Tas only | | Apply at cotyledon to 2 leaf stage. |
| | Guildford grass/Onion grass <i>Romulea rosea</i> | WA only | 15 | |
| | Hoary Cress <i>Cardaria draba</i> | Vic, SA, Tas only | 20 | Apply when plants are fully emerged. |
| | Lincoln weed <i>Diploaxis tenuifolia</i> | SA only | | |
| | Matricaria <i>Matricaria matrecoarioides</i> | WA and Tas only | | |
| | Mintweed <i>Salvia reflexa</i> | Qld, ACT and NSW only | | Apply at cotyledon to 4 leaf stage. |
| | Mouse-ear Chickweed <i>Cerastium</i> spp. | NSW, Vic, SA, WA, ACT, Tas only | 15 | |
| | Mustards <i>Sisymbrium</i> spp. | All States | | |
| | New Zealand Spinach <i>Tetragonia tetragonoides</i> | Qld only | 20 | |
| | Paterson's Curse/Salvation Jane <i>Echium plantagineum</i> | NSW, Vic, SA, WA, ACT, Tas only | 15 | |
| | Pimpernels <i>Anagallis arvensis</i> | NSW, Vic, SA, ACT, Tas only | | |
| | Prickly Lettuce/Whip Thistle <i>Lactuca serriola</i> | Vic, SA only | 20 | |
| | Rough Poppy <i>Papaver hybridum</i> | NSW, SA, WA, ACT, Tas only | 20 | |
| | Saltbush <i>Atriplex muelleri</i> | Qld, ACT and NSW only | | Apply at cotyledon to 4 leaf stage. |
| | Shepherd's Purse <i>Capsella bursa pastoris</i> | NSW, Vic, SA, WA, ACT, Tas only | | |
| | Slender Celery <i>Apium leptophyllum</i> | Qld, ACT and NSW only | | Apply at cotyledon to 4 leaf stage. |
| | Soursob <i>Oxalis pes-caprae</i> | NSW, Vic, SA, ACT only | | Apply when the majority of soursobs have emerged. |
| | Spear Thistle <i>Cirsium vulgare</i> | Tas only | | |
| | Stemless Thistle <i>Onopordum acaulon</i> | Vic only | 25 | |
| | Storksbill/Wild Geranium <i>Erodium</i> spp. | Vic, SA, WA, Tas only | 15 | |
| | Stagger Weed <i>Stachys arvensis</i> | Qld, NSW, ACT, WA, Tas only | 20 | |
| | Tree Hogweed <i>Polygonum patulum</i> | Vic only | | |
| | Turnip weed <i>Rapistrum rugosum</i> | Qld, NSW, ACT, SA only | 15 | |
| | Wireweed/Hogweed <i>Polygonum aviculare</i> | All States | 20 | |
| | Wild Turnip <i>Brassica tournefortii</i> | NSW, Vic, SA, WA, ACT and Tas only | 15 | |
| | Wild Radish <i>Raphanus rapharistrum</i> | All States | 15 or 20 | Use the higher rate under heavy weed pressure. A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations. |

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS
AUTHORISED UNDER APPROPRIATE LEGISLATION.**

GENERAL INSTRUCTIONS

This product is a selective herbicide designed to control certain weeds in wheat, triticale, barley, oats and cereal rye. This product is suitable as a pre-sowing treatment for wheat and triticale, and as a post-sowing treatment for wheat, triticale, barley, oats and cereal rye. When used on emerged weeds, the product is absorbed by foliage and green stems and moves into the root system. Prior to using this product, careful consideration should be given to soil pH. As soil pH increases, rate of breakdown decreases. This product should not be used on soil pH 8.6 or higher as extended soil residual activity could adversely affect crop rotation options beyond normal intervals. This product should not be used on soils with a pH of 8.6 or higher as soil residual activity could adversely affect following crops and crop rotation intervals may be extended beyond normal intervals. Crops other than wheat, barley, oats, triticale and cereal rye can be extremely sensitive to low concentrations of this product in the soil. See Crop Rotation Recommendations. Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks of application.

Pre-sowing incorporated by sowing:

WA only - Avoid applying to dry sandy soils as rapid leaching may occur with early season rains.

SA only - Before using rates greater than 15 g/ha on light to medium soils pH 7 to 8.5, seek further advice.

Conventionally Sown Crops - on soils less than 7, apply a spray just before sowing or in conjunction with the sowing operation. On soils of pH of 7 or greater it is not critical to time the spray just before sowing. Spray onto a non-ridged surface free of large clods. Use low profile 10 cm combine points for sowing. Sow at speeds of 10 kph or greater. Use light covering harrows at sowing. If applied to dry soil and sowing is to be delayed, incorporate immediately after spraying to, prevent loss by wind erosion.

Direct Drilled Crops - Apply tank mixed with either paraquat/diquat mixture or glyphosate in accordance with manufacturer's label recommendation.

Post Crop and Weed Emergence:

Where treatment is delayed or where weeds are not actively growing due to adverse conditions results may be slow to appear and weeds may be only stunted or suppressed.

Wheat, Triticale and Cereal Rye - Apply after crop emergence when weeds are small and actively growing (**Annual Ryegrass no more than 3 leaves, Broadleaved weeds no more than 5 cm in height or diameter** (for Black Bindweed refer to specific recommendations)).

Barley and Oats - apply between 2 leaf stage of crop (3 leaf stage - SA only) and early tillering when weeds are small and actively growing (**Annual Ryegrass no more than 3 leaves, Broadleaved weeds no more than 5 cm in height or diameter** (for Black Bindweed refer to specific recommendations)).

GRAZING ADVICE

Avoid grazing treated areas within 24 hours of application to optimise weed control. A nil withholding period is applicable for grazing Apparent Sword 750 WG Herbicide treated areas (when used as directed on this label).

CROP SAFETY:

DO NOT use this product for:

- Crops other than cereals
- Cereals irrigated by furrows or flooding.
- Winter cereals undersown with legume pasture crops.
- Weed control where crops are under stress. Damage can occur where crops are stressed due to conditions such as excessive soil alkalinity or acidity, poor nutrient status, disease, nematode or insect infestation, adverse weather conditions, drought or waterlogging. If crops become stressed after spraying, they may turn yellow or become retarded, but usually they will recover with no reduction in yield.

Wheat

DO NOT use this product for:

- Wheat varieties Cranbrook, or Miling.
- The wheat variety Vulcan if on acid soils and under stress conditions caused by waterlogging, frost, aluminium or manganese toxicity; reduced yields may result.
- Pre-sowing treatment of weeds in wheat varieties Avocet and Durati (OK for post-emergent use).
- Pre-sowing treatment of weeds in wheat variety Banks if soil pH is 5.5 or less (OK for post-emergent use).

Barley and Oats:

DO NOT use this product for:

- Application before the crop has reached the 2-leaf stage (3-leaf stage in SA).
- Stirling barley.
- Barley under waterlogged conditions (yield may be reduced).

The application of other sulfonylurea herbicides following this product is not recommended.

CROP ROTATION RECOMMENDATIONS

Land previously treated with this product should not be rotated to other crops other than those listed in the following tables. Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas. The treated areas may be replanted to any of the specified crops after the interval indicated in the following tables:

NB - THE TABLE BELOW APPLIES TO ALL STATES

| MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION) | | | | | | |
|--|--------------------|------------|------|--------|--|---|
| SOIL pH* | 0 | 3 | 6 | 9 | 12 | 18 |
| 6.5 or less | Triticale Wheat | Cereal Rye | Oats | Barley | Subterranean Clover** Faba Beans Field Pea Linseed Lucerne Lupins Medics ** Rapeseed Safflower | Maize Sorghum Soybeans Sunflower |

NB – THE TABLES BELOW APPLY TO Qld, SA, WA & Tas ONLY

| MINIMUM RECROPPING INTERVAL (Months After Application) | | | | | | |
|--|--|--|------------------------------|--|--------------------|--|
| Rainfall Requirement | 0 | 3 | 9 | 15 | 18 | 22 |
| | Minimum 700 mm | | | | | |
| Soil pH* 6.6 to 7.5 | Triticale Wheat | Cereal Rye | Barley Oats | Japanese Millet Maize Panicum Millet Sorghum Sunflower White French Millet | Cotton Soybeans | Faba Beans Field Pea Linseed Medics ** Rapeseed Safflower Subterranean Clover ** |
| MINIMUM RECROPPING INTERVAL (Months After Application) | | | | | | |
| Rainfall Requirement | 0 | 15 | 18 | 24 Months or longer | | |
| | Minimum 700 mm | | | | | |
| Soil pH* 7.6 to 8.5 | Triticale Wheat | Japanese Millet Maize Panicum Millet Sorghum Sunflower White French Millet | Barley Oats Cereal Rye | Rotate to crop other than Cereals (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season. | | |
| Soil pH* 8.6 and above | This product is not recommended for use on soils of pH 8.6 and above | | | | | |

NB – THE TABLES BELOW APPLY TO NSW, ACT & Vic ONLY

| MINIMUM RECROPPING INTERVAL (Months After Application) | | | | | |
|--|--------------------|------------|----------------|--|---|
| Soil pH* | 0 | 3 | 9 | 22 | 26 |
| 6.6 to 7.5 | Triticale Wheat | Cereal Rye | Barley Oats | Faba Beans Field Peas Linseed Lucerne Lupins Medics ** Subterranean Clover ** | Maize Sorghum Soybeans Sunflower |

| MINIMUM RECROPPING INTERVAL (Months After Application) | | | |
|--|--|----|---|
| Soil pH* | 0 | 18 | 24 months or longer |
| 7.6 to 8.5 | | | Rotate to crops other than Cereals (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season. |
| 8.6 and above | This product is not recommended for use on soils of pH 8.6 and above | | |

* Soil pH is to be determined by laboratory analysis using the 1:5 soil:water suspension method.

** Include natural regeneration of Subterranean Clover and Medics.

- Land and previously treated with this product should not be rotated to crops other than those listed in the above table
- Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas

SPRAY PREPARATION

This product is a water dispersible granule.

1. Fill tank partially with water and engage full agitation.
2. Add the required amount. (N.B. The measuring flask provided is graduated in grams of Apparent Sword 750 WG Herbicide only. **DO NOT** use for measuring of other materials.)
3. Top up with water to the required volume.
4. Companion products: If applying this product with another product ensure this product has completely dissolved before adding the companion product.
5. Apparent Sword 750 WG Herbicide must be kept in suspension at all times by continuous agitation. Where prepared spray mixes have been allowed to stand, thoroughly re-agitate before using.

USE OF SURFACTANT/WETTING AGENT

For post-emergence application, always add a non-ionic surfactant (1000 gac/L) at 100 mL/100 L of final spray volume (0.1% volume/volume).

The use of spraying oils is not recommended.

NOTE: DO NOT add a surfactant/wetting agent when this product is tank mixed with another product which already has a surfactant/wetting agent in the formulation.

GROUND SPRAYING EQUIPMENT

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern. Avoid overlapping and shut off spray booms while starting, turning, slowing or stopping as injury to the crop may result. Apply a minimum of 30 L of prepared spray per hectare.

AERIAL APPLICATION

Apply a minimum of 20 L/ha water. Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops. Avoid spraying where drift can go onto areas likely to be sown to sensitive crops - see Crop Rotation Recommendations. Turn off spray boom whilst passing over creeks and dams.

SPRAYER CLEAN-UP

It is essential that the sprayer be properly cleaned after using this product to prevent injury to crops other than wheat, triticale, barley, oats, or cereal rye. All traces of Apparent Sword 750 WG Herbicide should be removed from equipment using the following procedure:

1. Drain tank, then flush tank, boom and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water then add 300 mL household chlorine bleach (containing 4% chlorine) per 100 L of water. Flush through boom and hoses then allow to sit for 15 minutes with agitation engaged, then drain.
3. Repeat Step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through hoses and boom.

CAUTION: DO NOT use chlorine bleach with ammonia. All traces of liquid fertilizer containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour which can cause eye, nose, throat and lung irritation. DO NOT clean equipment in an enclosed area.

COMPATIBILITY

Chlorsulfuron is compatible with glyphosate and paraquat. The product does not control wild oats, however it is compatible with wild oat herbicides Avadex BW*, Mataven* and Puma*. It is also compatible with bromoxynil, MCPA (and bromoxynil/MCPA mixtures), 2,4-D Amine and 2,4-D Ester, Lontrel* L, Tigrex* and Jaguar*. This product is also compatible with Trifluralin and the insecticides omethoate, dimethoate, deltamethrin, fenvalerate and chlorpyrifos.