

**ENVIRONMENTALLY
HARZARDOUS SUBSTANCE,
LIQUID N.O.S (contains
oxyfluorfen)
MARINE POLLUTANT**

UN NO. 3082

Packaging Group III

HAZCHEM 2Z

In a Transport Emergency
**Dial
000**
Police or Fire Brigade



FOR SPECIALIST ADVICE
IN AN EMERGENCY DIAL
1800 033 111
ALL HOURS AUSTRALIA WIDE



Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932
Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone: (08) 9347 0500
Facsimile: (08) 9347 0551

APVMA Approval No: 58835/0704



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONQUEST

ENCORE

240 HERBICIDE

ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN
SOLVENTS: 606 g/L LIQUID HYDROCARBON
108 g/L N-METHYL PYRROLIDONE

GROUP G HERBICIDE

For the selective control of certain broadleaf and
grass weeds as per the Directions for use.

IMPORTANT: READ THIS BOOKLET BEFORE USE



Conquest Crop Protection Pty Ltd.
ABN 84 098 814 932
Level 1, 4 Collingwood St, Osborne Park, WA 6017
Telephone: (08) 9347 0500 ,
Facsimile: (08) 9347 0551

CROPS	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
<p>Tree fruit, Nuts and Vines not less than 3 years old</p> <p>DORMANT APPLICATION Almonds, Apples Apricots, Grapevine Macadamias, Peaches Plums, Pears, Pecans Walnuts</p>	<p>Refer to Weeds Controlled list</p>	<p>Weed free soil</p>	<p>3 or 4 L/ha</p>	<p>DO NOT apply once budswell has occurred. Apply to freshly cultivated weed free soil. Use higher rate for longer residual control (up to 4 months). Where grass weeds are expected to be the major problem, or when control of a wider weed spectrum is needed, mix the lower rate with 4.5 L Oryzalin (500g/L) or 4.5kg Oryzalin (500g/kg) per treated hectare. Read and follow all label directions.</p>
		<p>Weeds at 4-6 true leaf stage</p>		<p>DO NOT apply once budswell has occurred. Use higher rate for longer residual control (up to 4 months). When light infestations of weeds are present, apply as a tank mix with Conquest Glyphosate to give both knockdown and residual control. Read and follow all label directions. Where weed growth is large and dense weeds must be eliminated prior to application by using glyphosate or mechanical. Macadamias: Apply in 250 to 500 L water/ha. Apply after harvest to prevent spray-contacting nuts. Avoid spray contact with the macadamia tree foliage and stem. DO NOT apply to nuts on the ground.</p>
<p>Almonds, Apples Apricots, Grapevine Macadamias, Peaches Plums, Pears, Pecans Walnuts</p>	<p>Refer to Spray Conquest Explode 250 Herbicide and Sanction 200 Non-Residual Herbicide labels</p>	<p>Emerged weeds present</p>	<p>250mL plus Conquest Explode 250 Herbicide or Sanction 200 Non-Residual Herbicide at recommended label rates OR 25 to 30 mL/100L plus Conquest Explode 250 Herbicide or Sanction 200 Non-Residual Herbicide at recommended label rates</p>	<p>The addition of Conquest Encore 240 Herbicide to Conquest Explode 250 Herbicide or Sanction 200 Non-Residual Herbicide will improve control of small flowered mallow, evening primrose and other weeds sensitive to Encore 240.</p> <p>Apply as a tank mix with Conquest Explode 250 Herbicide or Sanction 200 Non-Residual Herbicide, refer to the appropriate label and follow all label directions.</p>

CROPS	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Almonds, Apples Apricots, Grapevine Macadamias, Peaches Plums, Pears, Pecans Walnuts	Refer to Glyphosate label	Apply as a tank mix with glyphosate products	75 mL/ha plus glyphosate at recommended label rates	Addition of Conquest Encore 240 Herbicide to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles <i>Urtica</i> spp., and Marshmallow, <i>Malva parviflora</i> . For rates refer to the appropriate label. Read and follow all label directions. DO NOT apply the tank mix of glyphosate and Conquest Encore 240 Herbicide near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.
Pyrethrum – greater than 4 leaf	Blackberry nightshade (<i>Solanum nigrum</i>) Fumitory (<i>Fumaria</i> spp.) Hogweed (<i>Polygonum aviculare</i>) Sorrel (<i>Rumex acetosella</i>) Volunteer potato (<i>Solanum tuberosum</i>) Cleavers (<i>Galium aparine</i>) Field Bindweed (<i>Convolvulus vulgaris</i>) Groundsel (<i>Senecio vulgaris</i>)	Emerged weeds present	100 – 150 mL/ha	Apply when pyrethrum growth stage is greater than 4 true leaf.
Pyrethrum – established crops, greater than 1 year old	As above plus Sow Thistle (<i>Sonchus</i> spp.) Spear Thistle (<i>Cirsium vulgare</i>)	Emerged weeds present	200 mL to 4 L/ha	Apply when pyrethrum is greater than 100 mm rosettes. Apply rates of greater than 1.0 L/ha ONLY between 1st of February and 31st of March. DO NOT apply later than 25 weeks before harvest.

CROPS	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Duboisia	Refer to Weeds Controlled List	Weed free soil	4 or 8 L	Apply after transplantation of young seedlings or after harvest of mature plants. Use higher rate where longer residual control (up to 6 months) is required. "Over-the-top" application will be tolerated.
		Weeds at 4 to 6 true leaf stage		Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Addition of a non-ionic surfactant such as Conquest Wetter 1000 Surfactant at recommended rates will improve activity. Established weeds must be eliminated prior to application by mechanical means.
Tobacco	Refer to Weeds Controlled List	Weed free soil	4 L	Use to control weeds along spray lines only. DO NOT apply to tobacco crop. Apply to soil after solid-set irrigation system has been laid out in the field. Conquest Encore 240 Herbicide should be applied to moist soil. Where very small weeds (2 to 3 leaf) emerge prior to spraying, the addition of a wetting agent to the spray mixture is necessary for effective control. Should the weeds be more advanced, the addition of 2 L/ha Sanction 200 Non-Residual Herbicide is required. Avoid spray drift.
Coffee	Refer to Weeds Controlled List	Weed free soil	2 to 4 L	Apply after transplanting coffee plants or under established bushes. Use the higher rate where longer residual activity (up to 4 months) is required. Best results are achieved when applied to moist soil, free of weeds with follow-up irrigation.
		Weeds at 2 to 4 true leaf stage		When seedlings are present, apply as a tank mix with Conquest Explode 250 Herbicide to produce both knockdown and residual control. Addition of a non-ionic surfactant such as Conquest Wetter 1000 at 150 mL/100L or Activator Surfactant at 125 mL/100L should be used in the spray mixture. Apply as a directed spray to avoid contact with coffee plants. Mature established weeds must be eliminated by mechanical or chemical means prior to application.

CROPS	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Onion (seeded)	Blackberry Nightshade (Solanum nigrum)	Hook leaf stage	50 to 100 mL/ha	DO NOT exceed 100 mL/ha at the hook leaf stage as excessive crop damage may occur.
	Common Cotula (<i>Cotula australis</i>) Crowsfoot/ Storksbill (<i>Erodium</i> spp.) Deadnettle (<i>Lamium amplexicaule</i>) Docks (<i>Rumex</i> spp.) Fumitory (Pink and white flowered) (<i>Fumaria</i> spp.) Groundsel (<i>Senecio vulgaris</i>) Hogweed/ Wireweed (<i>Polygonum aviculare</i>) Milk Thistle (<i>Sonchus oleraceus</i>) Plantain (<i>Plantago</i> spp.) Sorrel (<i>Rumex acetosella</i>) Potato weed (<i>Galinsoga parviflora</i>) Volunteer potato (<i>Solanum tuberosum</i>)	1½ to 2½ leaf stage	350 – 500 mL/ha	<p>Best results are obtained when the hook leaf stage treatment is followed by a further application of Conquest Encore 240 Herbicide when onions are at the 1½ to 2½-leaf stage. Apply Conquest Encore 240 Herbicide in at least 100 L/ha water at pressures below 150 kPa.</p> <p>DO NOT exceed 150 kPa because excessive crop damage could result. Multiple treatments may be applied provided the total dose does not exceed 2 L/ha in one season. For control of other weeds, use in a program with other registered herbicides.</p> <p>NOTE: Can cause flecking, twisting or stunting of onion plants although such symptoms are normally short-lived.</p> <p>DO NOT apply in conditions of very high humidity.</p> <p>DO NOT apply to weeds or crops, which are stressed by prevailing weather conditions, disease or mechanical damage.</p> <p>DO NOT use wetting agents or</p>

CROPS	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Tropical and Subtropical fruit (inedible peel) Custard Apple Kiwi Fruit, Longan Lychees, Mango Passionfruit, Pawpaw Rambutan	Refer to Weeds Controlled List	Weed free soil	4 L/ha	Best results are obtained when applied to moist soil free of weeds.
		Emerged weeds present		If weeds are present Conquest Encore 240 Herbicide should be applied as a tank mix with the recommended rates of Glyphosate Conquest Explode 250 Herbicide or Sanction 200 Non-Residual Herbicide.
Brassica Crops Broccoli Cabbages Cauliflower	Refer to Weeds Controlled List	Weed free soil	1.5 to 2 L/ha	Apply Conquest Encore 240 Herbicide to prepared ground 4 to 7 days prior to transplanting. If soil is dry irrigation or rainfall is required prior to transplanting for activation of Encore 240. Utilise transplanting techniques, which cause minimal soil disturbance. Excessive soil disturbance will lessen herbicide activity. Use the higher rate in situations where weed pressure is known to be heavy.
Forestry – <i>Eucalyptus</i> spp. <i>Pinus</i> spp.	Refer to Weeds Controlled List	Weed free soil	3 or 4 L/ha	Apply to weed free soil either before or preferably soon after transplanting. Disturbance of the herbicidal barrier on the soil surface by the transplantation process may reduce the length of weed control. Use higher rate for longer residual control. Apply as a directed or over-the-top spray. Conquest Encore 240 Herbicide may be tank mixed with Conquest Simaquest 900 WG Herbicide to extend the spectrum and length of weed control. Application to foliage that is soft and has not hardened off and/or if temperatures exceed 20°C may increase the likelihood of foliar damage to trees, eucalypts especially. Conquest Encore 240 Herbicide has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large-scale application.

CROPS	WEEDS CONTROLLED	TIME OF APPLICATION	RATE	CRITICAL COMMENTS
Forestry – <i>Eucalyptus</i> spp. <i>Pinus</i> spp.	Refer to Weeds Controlled List	Weeds at 4 – 6 true leaf stage	3 or 4 L/ha	Apply as an over-the-top spray to eucalypts and pines. Best results are achieved when applied to moist soil free of weeds. However, some newly germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled at these rates. Addition of a non-ionic surfactant such as Conquest Wetter 1000 at recommended rates will improve activity. Established weeds must be eliminated prior to application. Conquest Encore 240 Herbicide may be tank mixed with Conquest Simaquest 900 WG Herbicide to extend the spectrum and length of weed control. Application to foliage that is soft and has not hardened off and/or if temperatures exceed 20°C may increase the likelihood of foliar damage to trees, eucalypts especially. Conquest Encore 240 Herbicide has been found to be safe to commonly planted species used in forestry. Sensitivity of less common species should be checked prior to large-scale application.
Forestry Trees	Broadleaf weeds and grasses	Weeds at 4 – 6 true leaf stage	4 L/ha or 4 mL/10 m ²	For the establishment of trees for approved farm practices such as windbreaks, erosion control, woodlots and forestry planting. When applying as a post-plant spray, ensure spray is directed to the base of seedlings, or that seedlings are protected. Do not apply under hot or windy conditions.
Prior to sowing winter cereals	Refer to Glyphosate label	Apply as a tank mix with glyphosate products	75 mL/ha plus glyphosate at recommended label rates	Addition of Conquest Encore 240 Herbicide to glyphosate products will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with glyphosate applied alone) and give control of annual nettles <i>Urtica</i> spp., and Marshmallow, <i>Malva parviflora</i> . For rates refer to the appropriate label. Read and follow all label directions.
Prior to sowing Cotton				

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

GRAZING – DO NOT GRAZE TREATED WEEDS

HARVEST- NOT REQUIRED WHEN USED AS DIRECTED

WEEDS CONTROLLED		
BEFORE GERMINATION		SEEDLINGS
Amsinckia <i>Amsinckia</i> spp.	Prickly lettuce <i>Lactuca</i> spp.	Amsinckia <i>Amsinckia</i> spp.
Barley grass <i>Hordeum leporinum</i>	Red natal grass <i>Rhynchelytrum repens</i>	Bellvine <i>Ipomoea</i> spp.
Barnyard grass <i>Echinochloa</i> spp.	Redshank <i>Amaranthus cruentus</i>	Capeweed <i>Arctoteca calendula</i>
Blackberry nightshade <i>Solanum nigrum</i>	Ryegrass <i>Lolium</i> spp.	Common Cotula <i>Cotula australis</i>
Bladder ketmia <i>Hibiscus trionum</i>	Sesbania pea <i>Sesbania cannabina</i>	Crowsfoot grass <i>Eleusine indica</i>
Burrgrass <i>Cenchrus australis</i>	Shepherd's purse <i>Capsella bursa-pastoris</i>	Deadnettle <i>Lamium amplexicaule</i>
Caltrop <i>Tribulus terrestris</i>	Smallflower mallow <i>Malva parviflora</i>	Groundsel <i>Senecio vulgaris</i>
Capeweed <i>Arctoteca calendula</i>	Soursob <i>Oxalis pes-caprae</i>	Liverseed grass <i>Urochloa panicoides</i>
Chickweed <i>Stellaria media</i>	Sowthistle <i>Sonchus oleraceus</i>	Marshmallow <i>Malva parviflora</i>
Crowsfoot grass <i>Eleusine indica</i>	Starburr <i>Acanthospermum hispidum</i>	Pigweed <i>Portulaca oleracea</i>
Deadnettle <i>Lamium amplexicaule</i>	Stinkgrass <i>Eragrostis cillanensis</i>	Potato weed <i>Galinsoga parviflora</i>
Fat hen <i>Chenopodium album</i>	Summer grass <i>Digitaria</i> spp.	Redshank <i>Amaranthus cruentus</i>
Giant pigweed <i>Trianthema portulacastrum</i>	Thornapple <i>Datura stramonium</i>	Shepherd's purse <i>Capsella bursa-pastoris</i>
Liverseed grass <i>Urochloa panicoides</i>	White eye <i>Richardia brasiliensis</i>	Sowthistle <i>Sonchus oleraceus</i>
Lovegrass <i>Eragrostis</i> spp.	Wild mustard <i>Sisymbrium</i> spp.	Stinging nettle <i>Urtica urens</i>
Pigeon grass <i>Setaria</i> spp.	Wild radish <i>Raphanus raphanistrum</i>	Stinkgrass <i>Eragrostis cilianensis</i>
Pigweed <i>Portulaca oleracea</i>	Wireweed <i>Polygonum aviculare</i>	Wild radish <i>Raphanus raphanistrum</i>

GENERAL INSTRUCTIONS

Conquest Encore 240 Herbicide is a selective herbicide for the control of certain annual grasses and broadleaf weeds in dormant apples, grapes, peaches, pears, plums, apricots, almonds, macadamia, dubosia, tobacco, coffee, pyrethrum, winter cereals, tropical/sub-tropical crops, brassicas, onions, Eucalyptus and Pinus spp and other forestry trees used in forestry applied either to a weed free soil or to seedling weeds at the 4 to 6 true leaf stage.

Conquest Encore 240 Herbicide applied to well prepared weed free soil should not be disturbed or incorporated after application. Weed control for up to 6 months can be expected, but spot treatment with knockdown herbicides, of escape weeds are perennial grasses may be necessary.

When Conquest Encore 240 Herbicide is applied to seedling weeds at the 4 to 6 true leaf stage a non-ionic surfactant such as Conquest Wetter 100 surfactant should be added at recommended rates to improve activity.

Conquest Encore 240 Herbicide can also be utilised at low rates to improve the weed spectrum of knockdown herbicides.

Dubosia seedlings and mature plants will tolerate "over-the-top" applications. Eucalypt and pine transplants and established trees can tolerate "over-the-top" applications.

When using Conquest Encore 240 Herbicide in combination with other herbicides, refer to the appropriate label and read and follow all label directions.

RESISTANT WEEDS WARNING

GROUP G HERBICIDE

Conquest Encore 240 Herbicide is a member of the diphenyl ether group of herbicides. The product has the inhibitor of protoporphyrinogen oxidase mode of action. For weed resistance management, the product is a Group G Herbicide. Some naturally occurring weed biotypes resistant to the product and other Group G herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other Group G herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Crop Protection Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds.

TIMING

For optimum residual weed control, Conquest Encore 240 Herbicide should be applied to the soil surface prior to weed emergence after all other agricultural operations have been completed, such as mechanical cultivation and reshaping of irrigation furrows. The area should be left undisturbed during the period of desired weed control. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected but spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary.

MIXING

Shake well before use. When using Conquest Encore 240 Herbicide alone, fill the spray tank, at least one-third full with clean water, add the recommended amount of Conquest Encore 240 Herbicide while the pump and agitator are running, then complete filling the spray tank.

A non-ionic surfactant, if required by label directions, should be added near the end of the filling process to minimise foaming.

When tank mixing with Oryzalin, add to one-third filled tank, then the Conquest Encore 240 Herbicide during the filling operation. Maintain agitation during mixing and until spraying is completed.

When tank mixing with Glyphosate, Conquest Explode 250 Herbicide, add these after Conquest Encore 240 Herbicide during the filling operation.

When using Conquest Encore 240 Herbicide in mixtures always refer to the appropriate label and read and follow all label directions.

APPLICATION

Conquest Encore 240 Herbicide should be applied as a ground spray directed at the base of the crop plants. The spray mixture should be applied evenly with a carefully calibrated, standard low-pressure herbicide sprayer with a flat fan or hollow cone nozzle. When seedling weeds are present ensure thorough coverage of their foliage is required to obtain the maximum knockdown effect. Ensure both the weed foliage and the soil surface are sprayed. When applying to bare soil use 250 to 500 L of water/ha. For application to seedling weeds (4 to 6 leaf stage) use 250 to 1,350 L/ha. Use higher volumes for high weed density.

CROP SAFETY

Conquest Encore 240 Herbicide may be applied as directed around dormant peach, plum, apricot, almond, apple and xpear trees and grape vines of all ages when applied at rates of less than 1.0 L/ha. When applied at 3.0 L/ha and above, the trees and grape vines should be at least 3 years of age. Do not apply Conquest Encore 240 Herbicide once bud swell has occurred when using rates greater than 1.0 L/ha. Duboisia seedlings and mature plants will tolerate "over-the-top" applications of Encore 240. When using the 75-mL/ha rate, Conquest Encore 240 Herbicide may be applied as directed prior to sowing winter cereals.

COMPATIBILITY:

This product is compatible with Glyphosate, Conquest Explode 250 Herbicide, Oryzalin and Conquest Sanction 200 Non-Residual Herbicide.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply Conquest Encore 240 Herbicide under weather conditions or from spraying equipment, which may cause spray to drift onto nearby water bodies, susceptible crops/plants, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Use with care when applying in areas frequented by stock.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will damage eyes and will irritate the skin. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length Butyl Rubber gloves and goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26). If swallowed, DO NOT induce vomiting. Give a glass of water.

MATERIAL SAFETY DATA SHEET

If additional hazard information is required refer to the Material Safety Data Sheet or contact Conquest Crop Protection Pty. Ltd.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.