

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

OzCrop

# Dicamba-M SL

## HERBICIDE

### ACTIVE CONSTITUENTS :

340 g/L MCPA (present as the dimethylamine salt)

80 g/L DICAMBA (present as the dimethylamine salt)

GROUP I HERBICIDE

*For the control of certain broadleaf weeds in winter cereals, pastures and non-crop areas.*

### CONTENTS:

**5L - 20L**

NOT A DANGEROUS GOOD  
ACCORDING TO THE AUSTRALIAN  
DANGEROUS GOODS (ADG) CODE.

BATCH NO.

DATE OF MANUFACTURE:

**OzCrop Pty Ltd** ABN 16 160 656 431  
6C/26 Redland Road, Capalaba QLD 4157 Australia  
Tel: 07 3198 4930 • Fax: 07 3198 4939  
Email: [enquiries@ozcrop.com.au](mailto:enquiries@ozcrop.com.au)  
[www.ozcrop.com.au](http://www.ozcrop.com.au)

**DIRECTIONS FOR USE**

**RESTRAINTS:**

DO NOT spray when rain seems likely to occur within 4 hours.

DO NOT spray when weeds are wet with dew or rain or under stress from drought, low soil fertility, extreme cold or water logging.

DO NOT spray outside recommended crop growth as crop damage may result.

DO NOT apply to crops or grass pastures with clover, lucerne or medics.

DO NOT apply to legume pastures.

DO NOT apply to plants suffering from stress.

Crop & Situation	Weeds Controlled	State	Rate	Critical Comments
<b>Barley, Commercial Rye, Oats, Triticale, Wheat</b> From early tillering stage (Zadoks 21)	Capeweed, Charlock, Common Ice-Plant, Seedling Docks, Doublegee (Threecomered Jack, Spiny Emex), Hexham Scent (Melilotus), Mustards, Soldier Thistle, Sorrel Seedlings, Tares, Tree Hogweed, Turnip Weed ( <i>Rapistrum rugosum</i> ), Vetches, Ward's Weed, Wild Turnip ( <i>Brassica toumefortii</i> ), Wireweed (Hogweed)	All States	1L/ha	Spray when most weeds are in the 2 to 4 leaf stage. In some instances suppression only of Wireweed (Hogweed), Capeweed and Wild Radish will occur.
	Seedling Cotton Thistle, Fat-Hen	Tas		
<b>Barley, Commercial Rye, Oats, Triticale, Wheat</b> From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)	Cotton Thistle to bud formation	Tas	1.7L/ha	Spray when most weeds are still in the young rosette or seedling stage.  Note: Wild Radish may need to be resprayed (with 2,4-D or MCPA) if future germinations occur after spraying. In some instances suppression only of Wireweed (Hogweed), Hoary Cress and Climbing Buckwheat will be obtained.
	Mintweed, New Zealand Spinach	Qld, NSW only		
	Doublegee (Threecomered Jack, Spiny Emex), Wireweed, Vetch, Variegated Thistle, Tares, Sunflower, Hexham Scent (Melilotus), Docks	SA only		
	Capeweed, Charlock, Chickweed, Climbing Buckwheat (Black Bindweed), Clover, Common Ice-Plant, Seedling Docks, Doublegee (Threecomered Jack, Spiny Emex), Fat-Hen, Hexham Scent (Melilotus), Seedling Hoary Cress, Mouse-eared Chickweed, Mustards, Narrow-leaved Plantain, Saffron Thistle, Spurry, Swinecress, Tares, Tree Hogweed, Turnip Weed ( <i>Rapistrum rugosum</i> ), Variegated Thistle, Vetches, Wild Radish, Wild Turnip ( <i>Brassica toumefortii</i> ), Wireweed (Hogweed)	QLD, NSW, Tas, WA, Vic only  SA only	1.4L/ha	
<b>Barley, Commercial Rye, Triticale</b> From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)	Volunteer Peas	SA only	1L/ha plus 60mL/ha dicamba (500 g/L)	Apply when most peas are 5-7cm high.
<b>Oats, Wheat</b> From early tillering stage (Zadoks 21) until the fully tillered stage and before jointing occurs (Zadoks 30)			1L/ha plus 120mL/ha dicamba (500 g/L)	

Crop & Situation	Weeds Controlled	State	Rate			Critical Comments
			Per Hectare	Per 100L	Per 15L Knapsack	
<b>Grass Pastures</b>	Bathurst Burr, Caltrop (Yellow Vine), Chicory, Crofton Weed, Cudweed, Dandelion, Deadnettle, Fleabanes, Fumitory, Groundsel, Heliotrope, Inkweed, Mintweed, Nettles, Noogoora Burr, Pennycress, Potato Weed, Ragwort, Ripplewort, Shepherd's Purse, Smart Weed, Soft Rush, Stinkwort, Thistles including Cockspur, Scotch (Spear), Slender, Star, Variegated, Wireweed (Hogweed)	All States	2.8-4L	190mL-270mL	60mL	Use the higher rate on larger weeds.
	Cotton Thistle	Tas				
	Buttercups, Hoary Cress seedlings, Horehound, Khaki Weed, Purple-Top, Veined Verbena	All States	4-7L	470mL	100mL	When necessary a second treatment may be made after 4 weeks.
<b>Turf</b>	Bindy-eye, Capeweed, Catsear, Chickweed, Clovers, Cotula, Creeping Oxalis, Cudweeds, Dandelion, Dock, Fleabane, Fumitory, Jo-Jo (Onehunga), Lambs Tongue, Medics, Mustards, Narrow-Leafed Plantain, Pearlwort, Pennywort, Peppergrass, Radish, Scarlet Pimpernel, Slender Thistle, Soldier Thistle, Spear Thistle, Toad Rush, Variegated Thistle, Wild Turnip, Wireweed (Hogweed)	All States	Golf Tee, Golf Green: 100mL in 15L water per 150m <sup>2</sup>  Bowling Green: 900mL in 200L water per 1400m <sup>2</sup>  Large Turf Area: (Field, Park, Golf Fairway) 6.5L in 250-400L water per 10 000m <sup>2</sup> (1 hectare)			Apply to actively growing weeds and moist soil. DO NOT use on Lippia or Strawberry Clover. DO NOT mow turf for two days before or after application or fertilise with 2 weeks of spraying. DO NOT reapply to Buffalo Grass within 12 months.
	Fat-Hen, Purple-Top, Red Flowered Mallow, Staggerweed, Swinecress	Qld, NSW, Tas, SA, WA only				
<b>Non-Crop Situations</b>	Artichoke, Bindweed, Fennel, Hemlock, Thistle seedlings	All States	5.2L	350mL	80mL	A repeat spray may be necessary.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIOD:**

**GRAZING - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.**

**HARVEST - NOT REQUIRED WHEN USED AS DIRECTED.**

## GENERAL INSTRUCTIONS

### MIXING DIRECTIONS:

Add the required amount of the product directly to the tank and mix well.

### APPLICATION:

#### For Boom Application:

Apply by a properly calibrated boom spray using not less than 50 litres of water per hectare unless indicated otherwise in Critical Comments column.

#### For Aerial Application:

Apply at least 25 litres total spray volume per hectare. The aircraft should fly as low as practical under the prevailing conditions to minimise drift.

#### For High Volume Application:

Apply a minimum 1500 Litres per hectare.

#### For Knapsack Application:

A 15 Litre Knapsack treats 150 square metres.

### Note:

Spray when weeds are young and actively growing. A temporary wilting may be evident in crops after application. If seasonal conditions are unusually late or dry, seek further advice as damage result from spraying under these conditions. Growers should seek advice before spraying recently released cereal varieties. There is a nil tolerance of pea seed contaminates in cereal grain and less than 100% control of Volunteer Peas could result in dockage.

### COMPATIBILITY:

OZCROP DICAMBA/MCPA Herbicide is compatible with Flowable terbutryn.

### RESISTANT WEEDS WARNING

OZCROP DICAMBA-M SL HERBICIDE contains members of the Benzoic acid and Phenoxy groups of herbicides. OZCROP DICAMBA-M SL HERBICIDE has the disrupters of plant cell growth mode of action. For herbicide resistance management OZCROP DICAMBA-M SL HERBICIDE is a Group I Herbicide. Some naturally occurring weed biotypes resistant to OZCROP DICAMBA-M SL HERBICIDE and other Group I Herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if OZCROP DICAMBA-M SL HERBICIDE or other Group I Herbicides are used repeatedly. These resistant weeds will not be controlled by OZCROP DICAMBA-M SL HERBICIDE or other Group I Herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Ozcrop Pty Ltd accepts no liability for any losses that may result from the failure of OZCROP DICAMBA-M SL HERBICIDE to control resistant weeds.

GROUP	I	HERBICIDE
-------	---	-----------

### PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures. Spray equipment must be calibrated accurately prior to working on crops. Keep container sealed (airtight) when not in use. Avoid spray drift and vapour movement onto susceptible crops such as cotton, tobacco, tomatoes, vines, lupins, fruit trees and ornamentals. Minimise spray drift by using low pressure and nozzles, which DO NOT have a fine droplet size. DO NOT apply this product on or near desirable trees or plants or in locations where the chemical may be washed or moved into contact with their roots. DO NOT use in high wind. All spray equipment must be thoroughly washed out after use and must not be used for spraying horticultural crops. DO NOT use at higher rate than recommended.

### PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT use the container for any other purpose. DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area away from children, animals, food, feedstuffs, herbicides, seed and fertilisers. DO NOT store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

## SAFETY DIRECTIONS

Harmful if absorbed by skin contact or inhaled or swallowed. Will irritate the eyes and skin. Do not inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves and face shield or goggles. When using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water after each days use, wash gloves, face shield or goggles and contaminated clothing.

### SAFE WORK AUSTRALIA

Causes serious eye damage.

Suspected of causing cancer.

Repeated exposure may cause skin dryness and cracking.

Do not eat, drink or smoke when using this product.

IF ON SKIN: Wash with plenty of soap and water.

IF SWALLOWED: Call a POISON CENTER or doctor/physician if you feel unwell.

Rinse mouth.

## FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).

## SAFETY DATA SHEET

Additional information is listed in the safety data sheet (SDS). A safety data sheet for OZCROP DICAMBA-M SL HERBICIDE is available from OZCROP PTY LTD. on request. Call Customer Service on 07 3198 4930.

**CONDITIONS OF SALE:** The use of OZCROP DICAMBA-M SL HERBICIDE being beyond the control of the manufacturer, no warranty expressed or implied is given by OZCROP Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and OZCROP Pty Ltd., accepts no responsibility for any consequence whatsoever resulting from the use of this product.



**OzCrop Pty Ltd** ABN 16 160 656 431

6C/26 Redland Road, Capalaba QLD 4157 Australia

Tel: 07 3198 4930 • Fax: 07 3198 4939

Email: [enquiries@ozcrop.com.au](mailto:enquiries@ozcrop.com.au)

visit our website [www.ozcrop.com.au](http://www.ozcrop.com.au)

APVMA Approval No.: 86775/116544

Label Release: 201906

