

Virbac (Australia) Pty Limited	Page 1 of 3
Product Name:	ORALJECT GOAT & SHEEP WORMER BROAD SPECTRUM ANTHELMINTIC PASTE FOR GOATS AND SHEEP (38791)
Version Date:	29 April 2011
Pack sizes:	30 mL
Reason:	Administrative changes – draft label

Text above this line or in italics between parallel lines is not included in the label copy.

IMMEDIATE CONTAINER LABEL

FOR ANIMAL TREATMENT ONLY

**Oralject® GOAT & SHEEP WORMER
BROAD SPECTRUM ANTHELMINTIC PASTE FOR GOATS AND SHEEP**

ACTIVE CONSTITUENT: 30 mg/mL MORANTEL CITRATE



For the control and treatment of morantel susceptible mature and immature roundworms of sheep and goats including strains resistant to benzimidazole chemicals: Barber's Pole Worm (*Haemonchus contortus*), Black Scour Worm (*Trichostrongylus* spp.), Small Brown Stomach Worm (*Teladorsagia [Ostertagia] circumcincta*), Stomach Hair Worm (*Trichostrongylus axei*), Small Intestinal Worm (*Cooperia curticei*), Thin Necked Intestinal Worm (*Nematodirus* spp.), Large Mouthed Bowel Worm (*Chabertia ovina*), Large Bowel Worm (*Oesophagostomum venulosum*).

30 mL

READ LEAFLET BEFORE USING THIS PRODUCT

DIRECTIONS FOR USE:

Dosage and Administration: Calculate dose as 1 mL per 3 kg bodyweight. Turn ring until required dose setting appears on left margin of ring. Remove cap from nozzle. Insert nozzle into animal's empty mouth. Deposit paste on back of tongue. Lift animal's chin and hold mouth closed for a few seconds to ensure that full dose is swallowed. Replace cap on nozzle. This product is safe for use in young, stud and pregnant animals.

Resistance may develop to any chemical. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practises for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 7 days before slaughter for human consumption.

MILK: DO NOT USE in female sheep or goats that are producing, or may in the future produce milk or milk products for human consumption.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126, New Zealand 0800 764 766.*

DISPOSAL: Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE: Store below 30 °C (Room temperature). **KEEP OUT OF REACH OF CHILDREN.**

[B]

EXP:

APVMA No.: 38791/53574

Code:



VIRBAC (AUSTRALIA) PTY LIMITED
361 Horsley Road MILPERRA NSW 2214
Freecall Customer Service Line: 1800 009 847
www.virbac.com.au