



POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



Apparent Haloxyfop 520 Herbicide

ACTIVE CONSTITUENT: :520 g/L HALOXYFOP present as the haloxyfop-P methyl ester

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions For Use.

IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE

Contents: 250 mL, 1L, 5L, 10L, 20L

Apparent Pty Ltd
Suite G.08, 762 Toorak Rd,
Glen Iris, Vic
PO Box 3092,
Cotham PO, Vic 3101
Phone 03 9822 1321

STORAGE AND DISPOSAL

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as described above.

SAFETY DIRECTIONS

Harmful if swallowed, Will irritate the eyes and skin. Avoid contact with the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves, face shield or goggles and contaminated clothing. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre, Phone Australia 13 1126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet for Apparent Haloxyfop 520 Herbicide.

CONDITIONS OF SALE

The use of Apparent Haloxyfop 520 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequence whatsoever resulting from the use of this product.

APVMA Approval Number: 66422/0714

DOM:
BN:

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Apparent

Haloxypop 520

Herbicide

ACTIVE CONSTITUENT: 520g/L HALOXYFOP present as the haloxypop-P methyl ester

GROUP A HERBICIDE

For the post-emergent control of a wide range of annual and perennial grass weeds in grain legume and oilseed crops, lucerne, medic and clover pasture and seed crops, forestry, bananas, citrus, grapes, pineapples, pome and stone fruit, pyrethrum, tropical fruit and nut crops as specified in the Directions For Use.

IMPORTANT: READ THIS LEAFLET BEFORE USE

APVMA Approval No. 66422/0714

Apparent Pty Ltd
Suite G.08, 762 Toorak Rd,
Glen Iris, Vic
PO Box 3092,
Cotham PO, Vic 3101
Phone 03 9822 1321

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (waterlogged or drought affected), poor nutrition or previous herbicide treatment as reduced levels of control may result.

DO NOT spray if rain is likely to occur within one hour.

Table 1a. Winter crops – Canola, Chickpeas, Faba beans, Field peas, Lentils, Linola, Linseed, Lupins, Lucerne, Vetch, Medic and Clover pastures or seed crops.

| WEEDS CONTROLLED | WEED GROWTH STAGE | RATE (mL/ha) | | CRITICAL COMMENTS |
|--|-------------------|--------------------------|-------------------------|--|
| | | with Uptake Spraying Oil | with a non-ionic wetter | |
| Annual ryegrass | 2 to 4 leaf | 75 | 100 | <p><u>CANOLA, LINOLA AND LINSEED</u> DO NOT apply after the 8 leaf stage of the crop. DO NOT apply after the commencement of stem elongation.</p> |
| | Early Tillering | 100 | 100 | |
| Barley grass Brome grass Paradoxa grass Volunteer cereals | 2 to 4 leaf | 50 | 75 | <p>This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced.</p> |
| | Early Tillering | 75 | 100 | |
| Wild oats WA, SA, Vic, Tas, Southern and Central NSW | 2 to 4 leaf | 37.5 | 50 | <p>DO NOT apply more than 1 application of herbicide containing haloxyfop per crop. DO NOT apply after grazing.</p> |
| | Early Tillering | 50 | 75 | |
| Wild oats | 2 to 4 leaf | 50 | 75 | <p>See GENERAL INSTRUCTIONS, Spraying oils/wetters section.</p> <p><u>FIELD PEAS AND CANOLA:</u> The only oil recommended for use with Apparent Haloxyfop 520 is Uptake* Spraying Oil. Apparent Haloxyfop 520 + Lontrel* 750 SG + Uptake* Spraying Oil are compatible and selective to canola. This tank-mix is also compatible with atrazine and selective to triazine tolerant canola.</p> <p><u>LUPINS AND FIELD PEAS:</u> Mixtures with Brodal* or simazine may cause crop yellowing and separate applications are recommended.</p> <p><u>CHICKPEAS, FABA BEANS, LENTILS AND VETCH, LINOLA, LINSEED:</u> Broadleaf herbicides should not be added to Apparent Haloxyfop 520. Apply Apparent Haloxyfop 520 and broadleaf herbicides at least a week apart.</p> <p><u>LUCERNE, CLOVER OR MEDIC PASTURES:</u> If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/wetters section). If silver grass (<i>Vulpia</i> spp.) is present in pasture, simazine should be tank mixed with the higher rate of Apparent Haloxyfop 520 plus a non-ionic wetter.</p> |
| Northern NSW & Qld | Early Tillering | 75 | 100 | |

Table 1b. Winter crop growth stage application windows

| Crop | Crop Growth Stage |
|---|--|
| Lucerne, Medic and Clover pastures or seed crops | Apply from 2 nd trifoliolate leaf onwards. For <i>Erodium spp.</i> spraying, apply from cotyledon crop stage onwards. |
| Canola, Linola and Linseed | Apply from 2 leaf to 8 leaf stage of crop growth. DO NOT apply after the commencement of stem elongation This means that application must not occur after the 8 leaf stage, or if stem elongation commences before the 8 leaf stage, application must not occur after stem elongation has commenced. |
| Chickpeas, Faba beans, Field peas, Lentils, Lupins, Vetch | Apply from 2 nd leaf, 2 nd node or 2 nd branch to prior to flowering |

Table 2a. Lucerne, Medic and Clover seed crops and pastures. See table 1b for crop stages

| WEEDS CONTROLLED | WEED GROWTH STAGE | RATE (mL/ha) with Uptake Spraying Oil | CRITICAL COMMENTS |
|---|-------------------------------|---------------------------------------|--|
| Prairie grass (<i>Bromus catharticus</i>) | Up to early tillering | 100 | <p>See GENERAL INSTRUCTIONS, Spraying oils/wetters section.</p> <p>³ Use lower rate when growing conditions and crop or pasture competition are good and when weed populations are below 100 plants/m². Use the higher rate when weed populations are above 100 plants/m² or when crop or pasture competition is poor.</p> <p>NOTE : Storksbill may not be controlled if simazine or Broadstrike* are tank-mixed with Apparent Haloxyfop 520.</p> <p>LUCERNE, CLOVER OR MEDIC PASTURES: If grazed or cut for hay immediately prior to treatment delay application until all grasses have fully expanded leaves. Use 75 mL + spraying oil or 100 mL + wetter/ha. (See GENERAL INSTRUCTIONS, Spraying Oils/wetters section). If silver grass (<i>Vulpia spp.</i>) is present in pasture, simazine should be tank mixed with the higher rate of Apparent Haloxyfop 520 plus a non-ionic wetter.</p> |
| Musky or ferny leaf Storksbill: (<i>Erodium moschatum</i>) Common Crowsfoot or Common Storksbill (<i>Erodium cicutarium</i>) | Up to 6 leaf or 5 cm diameter | 50 - 75 ³ | |
| Long or shiny leaf storksbill (<i>E. botrys</i>) | Up to 8 leaf or 5 cm diameter | 75-100 | |

Table 2b. Lucerne, Medic and Clover seed crops only - not to be used for stockfeed. See table 1b for crop stages

| WEEDS CONTROLLED | WEED GROWTH STAGE | RATE (mL/ha) with Uptake Spraying Oil | CRITICAL COMMENTS |
|--|---------------------|---------------------------------------|--|
| Couch grass (suppression), Rhodes grass (control) | Tillering seedlings | 150 + 150 ⁴ | ⁴ For best suppression of couch or control of Rhodes grass, make 2 applications of Apparent Haloxyfop 520 2-4 weeks apart. Time second application to coincide with tillering stage of weeds and just after irrigation or significant rain. |
| Couch grass (control) Rhodes grass (control) | Established stands | 400 - 800 | Only treat actively growing weeds which are not moisture stressed. Use these rates for control of couch and Rhodes grass |

Table 3a. Summer crops – Cotton, Cowpea, Lucerne, Mung bean, Navy beans, , Peanuts, Soybeans, Sunflowers.

| WEEDS CONTROLLED | WEED GROWTH STAGE | RATE (mL/ha) with Uptake Spraying Oil | CRITICAL COMMENTS |
|---|---------------------------------|---------------------------------------|---|
| Australian millet | 2 leaf to tillering up to 15 cm | 150 | See GENERAL INSTRUCTIONS, Spraying oils/wetters section. |
| Barnyard grass | 2 to 5 leaf | 100 | NAVY BEANS, PEANUTS, SOYBEANS: For broadleaf weed control, Apparent Haloxyfop 520 Herbicide at 150mL/ha plus wetter may be tank mixed with Blazer* (except on navy beans) or Basagran*. Tank mixtures may cause transient leaf spotting on the crop but do not normally affect yield. |
| | Tillering up to 15 cm | 150 | |
| Crowsfoot grass Green panic Johnson grass (rhizome) | 2 leaf to tillering up to 15 cm | 150 | DO NOT tank mix broadleaf herbicides with Apparent Haloxyfop 520 if grasses have begun tillering or if the grasses are under moisture stress. |
| Johnson grass (seedling) Liverseed grass (seedling) Mossman river grass | Tillering and up to 15 cm | 150 | |
| Summer grass | 2 leaf to tillering up to 15 cm | 150 | DO NOT add Uptake Spraying Oil when mixing with Blazer* or Basagran*. |
| Volunteer cereals | 2 to 4 leaf | 100 | DO NOT use Blazer or Basagran* tank-mixes on Cowpeas. |
| | Tillering up to 15 cm | 150 | |

Table 3b. Summer crop growth stage application windows

| Crop | Crop Growth Stage |
|--|--|
| Lucerne | Apply from 2 nd trifoliate leaf onwards |
| Cowpea, Mung beans, Navy beans, Soybeans | Apply from 2 nd leaf to flowering |
| Peanuts | Apply from 2 nd leaf to pegging |
| Cotton | Apply from 2 nd leaf to before the onset of flowering |
| Sunflowers | Apply from 2 nd leaf to head initiation |

Table 4. Annual and Perennial grasses and *Erodium* spp. in Orchard, Vine and Plantation crops, forestry, and pyrethrum.

| CROPS | CROP GROWTH STAGE | WEEDS CONTROLLED | WEED GROWTH STAGE | RATE (mL/ha) with Uptake Spraying Oil | CRITICAL COMMENTS |
|---|-------------------|---|-------------------------------|---------------------------------------|---|
| Orchard, vine and plantation Crops including: Apples Avocado Banana Blueberry Citrus Custard apple Feijoa Grapevines Guava Kiwifruit Litchi (Lychee) Longan Mango Nashi Nut trees Passionfruit Paw paw Pear Persimmon Pineapple Rambutan Stone fruit | All growth stages | Perennial grasses: Couch Rhodes grass Slender rats tail grass | Established stands | 400 – 800 | See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Spray should be directed to the base of the tree or vine avoiding contact with fruit and foliage. |
| | | Buffel grass Green panic Johnson grass Kikuyu <i>Paspalum spp</i> <i>Setaria spp</i> | Vegetative to early tillering | 200 | |
| | | | Late tillering | 400 | Annual Grasses: Where treated in association with perennial grasses, these annual grasses will be controlled. |
| | | Annual grasses: Annual ryegrass Barley grass Barnyard grass Brome grass Crowsfoot grass Lesser canary grass Liverseed grass Mossman river grass Paradoxa grass Summer grass Volunteer cereals Wild oats | 2 leaf to tillering | 200 | |
| Forestry: <i>Pinus radiata</i> <i>Eucalyptus spp.</i> | | Annual grasses as above | Vegetative to tillering | 125 - 250 | Forestry: For annual grasses apply lowest rate to newly emerged grasses, increasing the rate as they develop. |
| Forestry: <i>Pinus pineaster</i> | | Barley grass Brome grass Rope twitch Barnyard grass <i>Erodium spp.</i> Volunteer cereals | Vegetative to tillering | 100 - 250 | Pyrethrum Tasmania only: For <i>Erodium spp</i> apply 75-100mL/ha if the main weed is <i>E. botrys</i> . Use 50 - 75 mL/ha if either <i>E. cicutarium</i> or <i>E. moschatum</i> are the main weeds. |
| Pyrethrum | | | | | |

Table 5. Apparent Haloxyfop 520 and Select® Herbicide tank-mixes – Canola, Chickpeas, Faba beans, Field peas, Lupins, Lentils

| WEEDS CONTROLLED | WEED GROWTH STAGE | RATE (mL/ha) | | CRITICAL COMMENTS |
|--|-------------------|------------------------|------------------|--|
| | | Apparent Haloxyfop 520 | Select Herbicide | |
| FOP/DIM susceptible Annual ryegrass + Volunteer barley Volunteer wheat Brome grass Wild oats Barley grass Phalaris | 2 to 4 leaf | 25 | 150 | See GENERAL INSTRUCTIONS, Spraying oils/wetters section. Use Uptake* Spraying Oil at 500mL/100L or Hasten* at 1L/100L. Apply at the same crop growth stages as those in Table 1b Winter Crops. |
| | Early tillering | 38 | 150 | |
| FOP resistant Annual ryegrass + Volunteer barley Volunteer wheat Brome grass Wild oats Barley grass Phalaris | 2 to 4 leaf | 25 | 200 | Lentils: Apply up to 7 node-early branching crop growth stage only. Lupins: Not for Qld. |
| | Early tillering | 38 | 250 | |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

HARVESTING WITHHOLDING PERIODS

Canola, chickpeas, cotton, cowpea, faba beans, field peas, lentils, linola, linseed, lupins, mung beans, navy beans, orchard crops, peanuts, plantation crops, soybeans, sunflowers, vetch or vine crops.

NOT REQUIRED WHEN USED AS DIRECTED

Medic and Clover Seed Crops:

DO NOT HARVEST FOR 7 DAYS AFTER APPLICATION

STOCK FOOD WITHHOLDING PERIODS:

Medic and Clover Pasture:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION

Lucerne:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION

Canola, Chickpeas, Cotton, Cowpea,
Faba Beans, Field Peas, Lentils, Linola,
Linseed, Lupins, Mung Beans, Navy Beans,
Peanuts, Soybeans, Sunflowers, Vetch:

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION

COTTON GIN TRASH MUST NOT BE FED TO ANIMALS.

GENERAL INSTRUCTIONS

Mixing

- Add water to the spray tank to 10 cm above the level of agitation and ensure the agitation device is working vigorously. (There must be a minimum of 100 L of water in the tank before any pesticide is added.)
- If tank mixing, firstly, add any soluble liquid formulations (e.g. LONTREL* Herbicide) and allow agitation for approximately one minute.
- Then add Apparent Haloxyfop 520 at the point where agitation is strongest. (Do not add Apparent Haloxyfop 520 through a strainer or sieve). Allow further agitation for one minute.
- Half fill the spray tank.
- If using wettable powder or water dispersible granules, or other emulsifiable concentration formulations (e.g. LORSBAN™ 750WG or Omethoate) these should be added after the Apparent Haloxyfop 520 to the half full spray tank ensuring vigorous agitation.
- Finally add Uptake* Spraying Oil or approved alternate spraying oil/wetter. (See section on spraying oils/wetters) and continue filling the tank to the required volume maintaining agitation at all times.
- Only mix sufficient solution for immediate use. Apparent Haloxyfop 520 and any other tank mixes should be applied immediately for best results.

Spraying Oils/wetters

Spraying Oils: It is essential to add an adjuvant to Apparent Haloxyfop 520. Best results will be achieved with Uptake™ Spraying Oil at 0.5 L/100 L of spray solution. Alternatively, other oils plus a non-ionic wetter may also be used. When other crop spraying oils are used, mix at 1.0 L/100 L and add a *non-ionic wetter (surfactant)* at 200 mL/100L of spray solution. Use of oil is not always recommended. See Critical Comments for specific situation recommendations.

Non-ionic Wetters: When Uptake or other oils are not used, a 100% concentrate non-ionic wetting agent such as BS-1000™ at 200 mL/100 L must be used along with the higher rate of Apparent Haloxyfop 520 as specified in the Directions For Use.

Where water volumes of less than 50 L/ha are used, DO NOT use less than 250 mL/ha of Uptake or 500 mL/ha for oils other than Uptake or less than 100 mL/ha of wetter.

CANOLA, LUCERNE, MEDIC AND CLOVER PASTURES AND SEED CROPS:

When tank mixing Apparent Haloxyfop 520 with Lontrel herbicides (canola only) or Broadstrike (lucerne, clover and medics), use Uptake Spraying Oil with the lower rates of Apparent Haloxyfop 520 or a wetting agent with the higher rates of Apparent Haloxyfop 520 unless otherwise specified. When mixing Apparent Haloxyfop 520 with other broadleaf herbicides on these crops, DO NOT use an oil, use a wetter instead.

FIELD PEAS AND CANOLA:

The oil recommended is Uptake Spraying Oil. Hasten is also recommended for use with tank-mixtures of Apparent Haloxyfop 520 and Select Herbicide.

For canola, Apparent Haloxyfop 520 + Lontrel 750SG + Uptake Spraying Oil are compatible and selective to canola. This tank-mixture is also compatible with atrazine or simazine and selective to triazine tolerant canola.

NAVY BEANS, PEANUTS, SOYBEANS:

When mixing with Blazer or Basagran DO NOT add spraying oil to these mixtures. DO NOT use these tank-mixes on cowpea,

Compatibility:

Ground use only: Apparent Haloxyfop 520 Herbicide can be tank mixed with:

| | |
|-----------------|--|
| Insecticides: | dimethoate Lorsban* 500 EC Insecticide Lorsban* 750WG Insecticide omethoate |
| Herbicides: | atrazine Basagran* Blazer* Broadstrike Herbicide Lontrel*.Herbicide Lontrel* 750SG MCPA ester (LVE) - <i>DO NOT exceed 700 mL/ha of MCPA LVE</i> Oryzalin Select* Herbicide simazine Starane 200 Herbicide |
| Fungicides: | Dithane DF* Dithane Rainshield |
| Trace elements: | magnesium sulphate zinc sulphate |

Apparent Haloxyfop 520 Herbicide is NOT COMPATIBLE with 2,4-D or MCPA as sodium or amine salts,

Aerial use: No product other than a recommended crop oil or wetter should be mixed with Apparent Haloxyfop 520 Herbicide when applied by air except for addition of Lontrel Forestry Herbicide for use in forestry and Lontrel 750SG for use in canola only.

Application

Apply Apparent Haloxyfop 520 Herbicide in sufficient water to obtain good coverage. It should be applied by an accurately calibrated ground rig or aircraft delivering droplets with a VMD of 200-300 microns.

The following spray volumes are recommended.

Ground application 50-150 L/ha

Aerial application 30 L/ha minimum

Use higher water volumes in orchards and in dense crops where the weeds may be shielded by the crop canopy,

CLEANING SPRAY EQUIPMENT

If broadleaf herbicides, particularly sulfonyleureas have been used in the spray equipment at any time prior to Apparent Haloxyfop 520, particular care should be taken to follow the directions on the relevant broadleaf herbicide label for equipment cleaning, or damage to susceptible crops may occur.

After using Apparent Haloxyfop 520, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse. After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate .. Before spraying cereals, maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. SURF™, Cold Water SURF Concentrate*, DynamoMatic Concentrate*, OMO* or DRIVE*) at 500 mL/100 L of water or the powder equivalent at 500 g/100 l of water, and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Chlorine based cleaners are not recommended.

Rinse water should be discharged onto a designated disposal area, or if this is unavailable, onto unused land away from desirable plants and water sources.

RESISTANT WEEDS WARNING



Apparent Haloxyfop 520 Herbicide is a member of the aryloxyphenoxy propionate group of herbicides. The product has the acetyl CoA carboxylase inhibitor mode of action. For weed resistance management Apparent Haloxyfop 520 Herbicide is a Group A herbicide.

Some naturally occurring weed biotypes resistant to the product and other inhibitors of acetyl CoA carboxylase herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by the product or other inhibitors of acetyl CoA carboxylase.

Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from the failure of the product to control resistant weeds.

Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Apparent Pty Ltd representative.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

- Haloxyfop 520 Herbicide damages cereals and grasses.
- DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.
- Cereal crops or grasses planted within twelve weeks of application may be damaged by the residual effects of Haloxyfop 520 Herbicide, particularly on light and red soils.

PROTECTION OF LIVESTOCK

DO NOT graze or cut treated crops for stock food except as specified under withholding periods.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Apparent Haloxyfop 520 Herbicide is toxic to fish. DO NOT contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Store in the closed, original container, in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. DO NOT store near feedstuffs, fertilisers or seeds.

Triple or preferably pressure rinse containers before disposal. Add rinsate to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

SMALL SPILL MANAGEMENT

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, cat litter or clay granules to the spill. When absorption is complete, sweep up material and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb this wash liquid for disposal as described above.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with the eyes and skin. When preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length PVC gloves and face shield or goggles. After each day's use, wash gloves, face shield or goggles and contaminated clothing. Wash hands after use

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 1126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed on the Material Safety Data Sheet for Apparent Haloxyfop 520 Herbicide.

CONDITIONS OF SALE

The use of Apparent Haloxyfop 520 Herbicide being beyond the control of the manufacturer no warranty expressed or implied is given by Apparent Pty Ltd regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Apparent Pty Ltd accepts with no responsibility for any consequence whatsoever resulting from the use of this product.

Apparent is a registered trademark

* registered trademarks