

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TAKTIC® WP

CATTLE DIP AND SPRAY

ACTIVE CONSTITUENT: 500 g/kg AMITRAZ



For the control of ticks, including organophosphate and synthetic pyrethroid resistant strains, on cattle in NSW and Qld and cattle tick on sheep, goats, deer and certain circus animals in NSW only.

Contents:
TAKTIC® WP 2.5 kg Net (5 x 500 g)
TAKTIC® Stabiliser 50 kg Net (5 x 10 kg)



INTERVET RURAL CO. PTY. LTD.
91-105 HARPIN STREET,
BENDIGO EAST, VICTORIA 3550
PHONE: (03) 5442 5011

® TAKTIC is a Registered trademark
LA3001102/00

Please insert
13 digit 80% size
barcode

9 000000000000

NRA Approval No: 41278/01

GENERAL INSTRUCTIONS

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT

TAKTIC® WP is toxic to fish. Do NOT contaminate dams, rivers or streams with TAKTIC® WP. used containers or dip or spray wash containing TAKTIC® WP. Do NOT use empty containers for any other purpose. If used in accordance with the directions and recommendations, TAKTIC® WP will not give rise to hazards to livestock or wildlife.

Each measure pack contains sufficient material to make up 1,000 L of dip or spray wash.

ROUTINE TICK CONTROL (Qld, NSW and NT only)

For routine control of specific types of ticks on cattle, apply TAKTIC® WP at the following intervals:

CATTLE TICK (*Boophilus microplus*): intervals of 19 to 21 days.

NZ CATTLE TICK (Bush tick) (*Haemaphysalis longicornis*): intervals of 7 to 21 days.

PARALYSIS TICK (*Ixodes holocyclus*) intervals of 7 to 10 days.

Do NOT leave longer intervals between treatments when the ticks are active. For further information see the enclosed usage leaflet or consult your Stock Inspector, District Veterinary Officer or Tick Extension Officer.

QUARANTINED HOLDINGS (NSW only)

For control of cattle tick (*B. microplus*) on cattle, sheep, goats and deer on quarantined holdings in NSW only, apply TAKTIC® WP at 14 to 21 day intervals or as directed by the NSW Board of Tick Control.

ANIMAL MOVEMENT (NSW only)

For control of cattle tick (*B. microplus*) on certain circus animals being moved from tick infested to tick free areas in NSW, apply TAKTIC® WP three times at 3 to 7 day intervals or as directed by the NSW Board of Tick Control.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet available from Intervet Rural Co. Pty. Ltd.

DISPOSAL

Dispose of empty containers by puncturing top, bottom, and sides, then bury in an approved landfill, or bury under at least 500 mm of soil in a non crop, non pasture area away from water courses or homes. Do not burn empty containers or product.

STORAGE

Store in the closed original container in a dry, well ventilated area, out of direct sunlight.
Store below 30°C (Room temperature).



INTERVET RURAL CO. PTY. LTD.
91-105 HARPIN STREET,
BENDIGO EAST, VICTORIA 3550
PHONE (03) 5442 5011

©TAKTIC is a Registered trademark
NRA Approval No: 41278/01

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TAKTIC® WP

CATTLE DIP AND SPRAY

ACTIVE CONSTITUENT: 500 g/kg AMITRAZ

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin.

Do NOT inhale dust or spray mist. When preparing dip or spray, using prepared dip or spray, wear cotton overalls buttoned to the neck and wrist, washable hat, elbow-length PVC gloves and face shield.

If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash contaminated clothing, gloves and face shield.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. PHONE 13 11 26.

OVER DOSAGE The specific antidote for animal toxicity is yohimbine (Reverzine -Parnell; Antagozil-Ilium).

PRECAUTION Do NOT use on horses, other equines, cats, other felines or dogs.

IN A TRANSPORT
EMERGENCY
DIAL 000
POLICE OR
FIRE BRIGADE



See precaution statement
LE3000042/00

DIRECTIONS FOR USE (Qld, NSW, NT and WA only)

	PLUNGE DIPS	STABILISED SPRAY RACES	UNSTABILISED SPRAY RACES
	Stabilisation is always required.	Stabilisation is required when unused spray wash is to be retained for later use.	Applies when all spray wash is to be used within 24 hours of initial charge.
INITIAL CHARGE	500 g TAKTIC® WP plus 10 kg TAKTIC® Stabiliser per 1,000 litres of water.		500 g TAKTIC® WP per 1,000 litres or 100 g TAKTIC® WP per 200 litres of water. Do NOT add TAKTIC® Stabiliser.
MIXING	Empty the contents of one pack TAKTIC® Stabiliser (10 kg) per 1,000 litres of water in dip or sump, taking care to distribute the Stabiliser along the surface of the dip.		
	For each 1,000 litres of water in dip or sump, empty 500 g TAKTIC® WP (one tin) into a bucket containing water and stir into a smooth, creamy suspension. Pour into dip/sump, then rinse bucket adding rinsings to dip/sump.		
STIRRING	Use at least 30 head of cattle through the dip.	Agitate vigorously with pump or manually with paddle.	
	It is essential that the dip/sump be stirred at the commencement of each day's dipping and again if there is a break of more than half an hour during dipping.		
TOPPING UP	When the level of the dip has fallen by 700 litres, add 700 litres of water and 500 g TAKTIC® WP plus 10 kg TAKTIC® Stabiliser. Mix and stir as above. It is not advisable to allow the dip to fall more than 700 litres from its original level.	Topping up-not recommended. See Reinforcement	Topping up-not recommended. See Reinforcement
REINFORCEMENT	Reinforcement not recommended. See Topping up	Each time the level of the spray wash in the sump drops by 300 litres add 100 g TAKTIC® WP and continue spraying. Do NOT add water. Do NOT add TAKTIC® Stabiliser. Continue spraying until no more wash can be pumped out. If spraying is to continue - recharge as before. When the wash is polluted, clean out the sump before recharging.	
	DO NOT RETAIN OPENED TAKTIC® WP TINS OR TAKTIC® STABILISER PACKS FOR LATER USE. Use all contents of each pack immediately after opening since TAKTIC® WP and TAKTIC® Stabiliser deteriorate in opened packs. Similarly, any unused, unstabilised dip wash should be discarded. It should NOT be retained for later use unless stabilised first. (See above)		
MANAGEMENT DIP/SPRAY RACES	It is important that TAKTIC® WP be used at the recommended concentration of 0.025%w/v. This will be achieved by charging and topping up/reinforcement as described above. If the level of wash is allowed to fall excessively without topping up/reinforcing, the concentration of TAKTIC® WP will fall and its efficacy will be reduced.		
	During a period when regular dipping is not in progress, TAKTIC® Stabiliser* must be added to the dip wash at intervals of no more than 2 months after the last topping up. Add one pack TAKTIC® Stabiliser* (10 kg) per 2,000 litres of dip wash and stir as before. *Hydrated (slaked) lime (80% purity) can be used for this purpose.	If after the spraying, unused spray wash is retained in the sump but not to be used during next 7 days, add TAKTIC® Stabiliser* at the rate of one complete pack (10 kg) per 2,000 litres of spray wash in the sump and stir as before. Further additions of TAKTIC® Stabiliser* must be made at intervals of 2 months when regular spraying is not in progress. *Hydrated lime can be used for this purpose.	
SAMPLING	Stir the dip/sump as described above before sampling		Sampling not required
	Take the sample from 1 metre below the surface at the jump-in end immediately after stirring.	Take the sample from a nozzle on the race or from the sump after vigorous stirring.	
	Analysis of dip samples is available through Biochemistry Branch, Animal Research Institute, 665 Fairfield Road, Yeerongpilly, Queensland 4105. Suitable sample bottles are obtainable from your local agent or Intervet Rural Co. Pty. Ltd. representative.		

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION. DO NOT USE ON HORSES, CATS OR DOGS.

WITHHOLDING PERIODS: MEAT: NIL. MILK: NIL. EXPORT SLAUGHTER INTERVAL: NIL.

Provided cattle and calves are treated in accordance with these recommendations, the use of TAKTIC® will not present any hazard to users of milk, milk products, meat or other edible portions.