

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 1L, 5L, 10L, 20L, 24L  
MAIN PANEL – IMMEDIATE CONTAINER LABEL**

---

## **CAUTION**

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY**

COOPERS®

# **Stampede®**

## **POUR ON LOUSICIDE FOR CATTLE AND SHEEP**

**ACTIVE CONSTITUENT: 20 g/L DIFLUBENZURON  
in 300 g/L N-methyl-2-pyrrolidone**

A versatile and easy to use IGR pour-on for use on both CATTLE and SHEEP for the control and treatment of diflubenzuron-susceptible biting and sucking lice.

**[CATTLE AND SHEEP PICTURES]**

**READ DIRECTIONS FOR USE BEFORE OPENING OR USING THIS PRODUCT.**

**1 Litre  
[5 Litres  
10 Litres  
20 Litres  
24 Litres]**

### **COOPERS LOGO**



File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 1 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 5L, 20L, 24L  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL

---

COOPERS®

**Stampede®**

**POUR ON LOUSICIDE FOR CATTLE AND SHEEP**

**Indications – CATTLE**

For the treatment of diflubenzuron-susceptible biting lice (*Bovicola bovis*) and sucking lice (*Linognathus vituli*, *Haemotopinus eurysternus* and *Solenopotes capillatus*) of cattle.

**DIRECTIONS FOR USE: CATTLE**

**Restraint:**

**DO NOT re-treat within 21 days.**

**Dosage and Administration:**

For topical use only.

**Dosage Guide – Cattle:**

Dose rate according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Weigh cattle with scales or use a weighband. If there is a large variation in weight, draft the mob into two or more mobs based on body weight and dose the animals to the heaviest in each mob.

Liveweight (kg)	Dose Volume (mL)	Cattle Treated 5L	Cattle Treated 20L	Cattle Treated 24L
Under 100 kg	25	200	800	960
101-200	40	125	500	600
201-300	50	100	400	480
301-400	60	83	333	400
401-500	70	71	285	342
Over 500 kg	Add 10 mL/100 kg			

**Application:**

Apply along the midline of the back between the wither and the base of the tail.

[DIAGRAM OF APPLICATION]

For better coverage of large cattle (over 500 kg), the calculated dosage may be applied in an 'I' shape, with stripes across the shoulders and across the base of the tail, joined by a stripe along the midline of the back between the wither and the base of the tail.

[DIAGRAM OF APPLICATION]

Treated and non-treated cattle should be kept separate for 6 to 8 weeks after treatment to reduce the risk of reinfestation.

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 2 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 5L, 20L, 24L  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL**

**Indications - SHEEP**

For the control of diflubenzuron-susceptible body lice (*Bovicola ovis*) of shorn sheep up to 7 days off-shears and on unshorn lambs up to 3 months of age.

**DIRECTIONS FOR USE: SHEEP**

**RESTRAINTS:**

**DO NOT use on lactating or pregnant ewes where milk or milk products may be used for human consumption.**

**DO NOT mix treated sheep with untreated sheep until at least 6 weeks after treatment.**

**DO NOT apply to wet sheep.**

**DO NOT use on sheep that have not been cleanly shorn for any reason e.g. Mycotic Dermatitis.**

**Dosage and Administration:**

For topical use only. Using a T-bar applicator apply along the midline of the back from the poll to the base of the tail. Do not deviate from the spine.

[DIAGRAM OF APPLICATION]

**Dosage Guide – Sheep:**

Dose rate according to the heaviest animal by liveweight in the group (rams, ewes, wethers, lambs, etc.) Do not underdose. Weigh with scales. If there is a large variation in weight, draft the mob into two or more mobs based on body weight and dose the animals to the heaviest in each mob. Treat all sheep in the mob. Check the accuracy of the applicator before and during treatment.

Liveweight (kg)	Dose Volume (mL)	Sheep Treated 5L	Sheep Treated 20L	Sheep Treated 24L
10-20	12	416	1666	2000
21-30	15	333	1333	1600
31-55	20	250	1000	1200
56-75	25	200	800	960
Over 75 kg	Add 5 mL/10 kg			

**Use in Lambing Ewe Flocks:**

**COOPERS STAMPEDE** may be used for lice control in the following situations:

- Ewes with lambs at foot (up to 3 months old and greater than 10 kg).** The ewes must be shorn and treated up to 7 days off-shears at the dose rates indicated in the table above. If lambing is complete unshorn lambs (not less than 10 kg) should be treated as per the table above. Re-treat lambs (not less than 10 kg) at shearing to ensure ongoing protection against lice.

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 5L, 20L, 24L  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL**

---

2. **Pregnant Ewes which have not commenced lambing.** The ewes must be shorn and treated up to 7 days off-shears at the dose rates indicated in the table above. Ewes should not be treated less than 4 weeks before lambing commences. Lambs (not less than 10 kg) born to treated ewes should be treated at a convenient time – (eg. lamb marking). At first shearing lambs (not less than 10 kg) should be treated to ensure ongoing protection against lice.

**GENERAL INSTRUCTIONS:**

Treat animals in accordance with good agricultural practice. Animals should not be treated in extremely hot weather (>40°C) or if under heat stress.

Efficacy of **COOPERS STAMPEDE** is not affected by moderate rainfall after application.

**COOPERS STAMPEDE** does not affect mammals, which do not rely on chitin synthesis in normal body functions.

**COOPERS STAMPEDE** contains a yellow dye to identify treated animals. This dye is scourable from wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces. Apply only with the **COOPERS STAMPEDE** applicators and draw-off tubes. Clean out the applicator using vegetable oil, dismantle to wipe clean, re-oil with vegetable oil and assemble. Do not rinse the applicator equipment with water.

**MODE OF ACTION:**

**COOPERS STAMPEDE** is based on a chemical from the benzoyl urea group known as insect growth regulators (IGR). Insect growth regulators prevent the deposition of chitin in larval species of insects, thereby causing death. Immature larvae present at application and those which hatch from eggs post application are prevented from developing to adults thus breaking the breeding cycle.

**COOPERS STAMPEDE** is not a knockdown insecticide. **COOPERS STAMPEDE** produces a rapid decline in lice numbers over several weeks. This is due to **COOPERS STAMPEDE** inhibiting the development of the immature stages of lice, adults take some weeks to die out.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**MEAT (CATTLE): DO NOT USE less than 10 days before slaughter for human consumption.**

**MILK (CATTLE): NIL.**

**MEAT (SHEEP): DO NOT USE less than 7 days before slaughter for human consumption.**

**MILK (SHEEP): DO NOT USE on lactating or pregnant ewes where milk or milk products may be used for human consumption.**

**WOOL HARVESTING INTERVAL: Wool should not be harvested from treated sheep within six months of treatment.**

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 4 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 5L, 20L, 24L  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL

---

**TRADE ADVICE:**

EXPORT SLAUGHTER INTERVAL (ESI): CATTLE – DO NOT USE less than 10 days before slaughter for export. SHEEP – DO NOT USE less than 7 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1 800 226 511 or the APVMA website [www.apvma.gov.au/residues/](http://www.apvma.gov.au/residues/)

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues: adequate treatment records should be kept and made available, if requested by wool buyers.

**SAFETY DIRECTIONS:**

**Will irritate eyes and skin. Avoid contact with eyes and skin. When using the product wear elbow length butyl rubber gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves.**

**ADDITIONAL USER SAFETY INFORMATION**

**SHEEP RE-HANDLING INTERVAL:** After treatment, wait until sheep are dry before rehandling.

**FIRST AID:**

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.**

Additional Information is listed in the Material Safety Data Sheet.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

Do not contaminate dams, ponds, waterways or drains with **COOPERS STAMPEDE**.

**SPECIALIST ADVICE IN EMERGENCY ONLY**

Intervet Australia Pty Limited  
(MSD Animal Health)  
**1 800 226 511**

**ALL HOURS – AUSTRALIA WIDE**

**DRUMMUSTER  
ELIGIBLE  
CONTAINER**

**DISPOSAL**

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 5 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 5L, 20L, 24L  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL**

---

**Warranty**

Intervet Australia Pty Ltd (IAPL), known as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

**COOPERS ANIMAL HEALTH**

Intervet Australia Pty Limited,  
91-105 Harpin Street, Bendigo East VIC 3550

**STORAGE**

Store below 30°C (Room Temperature) in the closed, original container in a dry well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA 61351/0614

® Registered Trademarks

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

**BATCH  
EXPIRY**

**5L [20L, 24L]**

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 6 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 1L, 10L ZIPSEAL TOP  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL**

---

TO OPEN PEEL HERE ↗

**COOPERS®**  
**Stampede®**  
**POUR ON LOUSICIDE FOR CATTLE AND SHEEP**

**READ ENCLOSED DIRECTIONS BEFORE USE**

**CATTLE**

For the treatment of diflubenzuron-susceptible\_biting lice (*Bovicola bovis*) and sucking lice (*Linognathus vituli*, *Haemotopinus eurytarnus* and *Solenopotes capillatus*) of cattle.

**SHEEP**

For the control of diflubenzuron-susceptible\_body lice (*Bovicola ovis*) of shorn sheep up to 7 days off-shears and on unshorn lambs up to 3 months of age.

**SAFETY DIRECTIONS:**

**Will irritate eyes and skin. Avoid contact with eyes and skin. When using the product wear elbow length butyl rubber gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves.**

**ADDITIONAL USER SAFTEY INFORMATION**

**SHEEP RE-HANDLING INTERVAL: After treatment, wait until sheep are dry before rehandling.**

**FIRST AID:**

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.**

Additional Information is listed in the Material Safety Data Sheet.

[1 L]

**Dispose** of empty container by wrapping with paper and putting in garbage.

[10 L]

**DISPOSAL**

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 7 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 1L, 10L ZIPSEAL TOP  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL (CONTINUED)**

---

**COOPERS ANIMAL HEALTH**

Intervet Australia Pty Limited,  
91-105 Harpin Street, Bendigo East VIC 3550

**STORAGE**

Store below 30°C (Room Temperature). Do not freeze. Store in tightly closed original container in a well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval No. 61351/0614

**FOR BATCH AND EXPIRY SEE BELOW**

® Registered Trademarks

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

**1L [10L]**

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 8 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 1L, 10L ZIPSEAL BASE  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL**

---

COOPERS®

**Stampede®**

**POUR ON LOUSICIDE FOR CATTLE AND SHEEP**

**SAFETY DIRECTIONS:**

**Will irritate eyes and skin. Avoid contact with eyes and skin. When using the product wear elbow length butyl rubber gloves. If product on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves.**

**ADDITIONAL USER SAFETY INFORMATION**

**SHEEP RE-HANDLING INTERVAL: After treatment, wait until sheep are dry before rehandling.**

**FIRST AID:**

**If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, wash out immediately with water.**

Additional Information is listed in the Material Safety Data Sheet.

[1 L]

**Dispose** of empty container by wrapping with paper and putting in garbage.

[10 L]

**DISPOSAL**

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

**SPECIALIST ADVICE IN EMERGENCY ONLY**

Intervet Australia Pty Limited

(MSD Animal Health)

**1 800 226 511**

**ALL HOURS – AUSTRALIA WIDE**

**DRUMMUSTER  
RE ELIGIBLE  
CONTAINER**  
[10L only]

**Warranty**

Intervet Australia Pty Ltd (IAPL), known as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 9 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

**STAMPEDE POUR ON LOUSICIDE FOR CATTLE AND SHEEP – 1L, 10L ZIPSEAL BASE  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL (CONTINUED)**

---

**COOPERS ANIMAL HEALTH**

Intervet Australia Pty Limited,  
91-105 Harpin Street, Bendigo East VIC 3550

**STORE** below 30°C (Room Temperature). Do not freeze. Store in tightly closed original container in a well-ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval No. 61351/0614

**FOR BATCH AND EXPIRY SEE BELOW**

® Registered Trademarks

® Yellow (PANTONE® 123) the predominant colour of Coopers packaging is a Registered Trademark. PANTONE is a registered trademark of Pantone, Inc.

**1L [10L]**

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 10 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

STAMPEDE POUR ON LOUSICIDE FOR CATTLE & SHEEP – 1L, 10L ZIPSEAL LEAFLET  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL

---

**Indications – CATTLE**

For the treatment of diflubenzuron-susceptible biting lice (*Bovicola bovis*) and sucking lice (*Linognathus vituli*, *Haemotopinus eurysternus* and *Solenopotes capillatus*) of cattle.

**DIRECTIONS FOR USE: CATTLE**

**RESTRAINTS:**

**DO NOT re-treat within 21 days.**

**Dosage and Administration:**

For topical use only.

**Dosage Guide – Cattle:**

Dose rate according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Weigh cattle with scales or use a weighband. If there is a large variation in weight, draft the mob into two or more mobs based on body weight and dose the animals to the heaviest in each mob.

Liveweight (kg)	Dose Volume (mL)	Cattle Treated 1 L	Cattle Treated 10 L
Under 100 kg	25	40	400
101-200	40	25	250
201-300	50	20	200
301-400	60	16	166
401-500	70	14	142
Over 500 kg	Add 10 mL/100 kg		

**Application:**

Apply along the midline of the back between the wither and the base of the tail.

[DIAGRAM OF APPLICATION]

For better coverage of large cattle (over 500 kg), the calculated dosage may be applied in an 'I' shape, with stripes across the shoulders and across the base of the tail, joined by a stripe along the midline of the back between the wither and the base of the tail.

[DIAGRAM OF APPLICATION]

Treated and non-treated cattle should be kept separate for 6 to 8 weeks after treatment to reduce the risk of reinfestation.

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 11 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

STAMPEDE POUR ON LOUSICIDE FOR CATTLE & SHEEP – 1L, 10L ZIPSEAL LEAFLET  
 ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL (CONTINUED)

**Indications - SHEEP**

For the control of diflubenzuron-susceptible body lice (*Bovicola ovis*) of shorn sheep up to 7 days off-shears and on unshorn lambs up to 3 months of age.

**DIRECTIONS FOR USE: SHEEP**

**RESTRAINTS:**

**DO NOT use on lactating or pregnant ewes where milk or milk products may be used for human consumption.**

**DO NOT mix treated sheep with untreated sheep until at least 6 weeks after treatment.**

**DO NOT apply to wet sheep.**

**DO NOT use on sheep that have not been cleanly shorn for any reason e.g. Mycotic Dermatitis.**

**Dosage and Administration:**

For topical use only. Using a T-bar applicator apply along the midline of the back from the poll to the base of the tail. Do not deviate from the spine.

[DIAGRAM OF APPLICATION]

**Dosage Guide – Sheep:**

Dose rate according to the heaviest animal by liveweight in the group (rams, ewes, wethers, lambs, etc.) Do not underdose. Weigh with scales. If there is a large variation in weight, draft the mob into two or more mobs based on body weight and dose the animals to the heaviest in each mob. Treat all sheep in the mob. Check the accuracy of the applicator before and during treatment.

Liveweight (kg)	Dose Volume (mL)	Sheep Treated 1 L	Sheep Treated 10 L
10-20	12	83	833
21-30	15	66	666
31-55	20	50	500
56-75	25	40	400
Over 75 kg	Add 5 mL/10 kg		

**Use in Lambing Ewe Flocks:**

**COOPERS STAMPEDE** may be used for lice control in the following situations:

- Ewes with lambs at foot (up to 3 months old and greater than 10kg).** The ewes must be shorn and treated up to 7 days off-shears at the dose rates indicated in the table above. If lambing is complete unshorn lambs (not less than 10 kg) should be treated as per the table above. Re-treat lambs (not less than 10 kg) at shearing to ensure ongoing protection against lice.

Text appearing above the header and below the footer lines is not included in the label

---

STAMPEDE POUR ON LOUSICIDE FOR CATTLE & SHEEP – 1L, 10L ZIPSEAL LEAFLET  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL (CONTINUED)

---

2. **Pregnant Ewes which have not commenced lambing.** The ewes must be shorn and treated up to 7 days off-shears at the dose rates indicated in the table above. Ewes should not be treated less than 4 weeks before lambing commences. Lambs (not less than 10 kg) born to treated ewes should be treated at a convenient time – (eg. lamb marking). At first shearing lambs (not less than 10 kg) should be treated to ensure ongoing protection against lice.

**GENERAL INSTRUCTIONS:**

Treat animals in accordance with good agricultural practice. Animals should not be treated in extremely hot weather (>40°C) or if under heat stress.

Efficacy of **COOPERS STAMPEDE** is not affected by moderate rainfall after application.

**COOPERS STAMPEDE** does not affect mammals, which do not rely on chitin synthesis in normal body functions.

**COOPERS STAMPEDE** contains a yellow dye to identify treated animals. This dye is scourable from wool. The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces. Apply only with the **COOPERS STAMPEDE** applicators and draw-off tubes. Clean out the applicator using vegetable oil, dismantle to wipe clean, re-oil with vegetable oil and assemble. Do not rinse the applicator equipment with water.

**MODE OF ACTION:**

**COOPERS STAMPEDE** is based on a chemical from the benzoyl urea group known as insect growth regulators (IGR). Insect growth regulators prevent the deposition of chitin in larval species of insects, thereby causing death. Immature larvae present at application and those which hatch from eggs post application are prevented from developing to adults thus breaking the breeding cycle.

**COOPERS STAMPEDE** is not a knockdown insecticide. **COOPERS STAMPEDE** produces a rapid decline in lice numbers over several weeks. This is due to **COOPERS STAMPEDE** inhibiting the development of the immature stages of lice, adults take some weeks to die out.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**MEAT (CATTLE): DO NOT USE less than 10 days before slaughter for human consumption.**

**MILK (CATTLE): NIL.**

**MEAT (SHEEP): DO NOT USE less than 7 days before slaughter for human consumption.**

**MILK (SHEEP): DO NOT USE on lactating or pregnant ewes where milk or milk products may be used for human consumption.**

**WOOL HARVESTING INTERVAL: Wool should not be harvested from treated sheep within six months of treatment.**

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 13 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				

Text appearing above the header and below the footer lines is not included in the label

---

STAMPEDE POUR ON LOUSICIDE FOR CATTLE & SHEEP – 1L, 10L ZIPSEAL LEAFLET  
ANCILLIARY PANELS – IMMEDIATE CONTAINER LABEL (CONTINUED)

---

**TRADE ADVICE:**

EXPORT SLAUGHTER INTERVAL (ESI): CATTLE – DO NOT USE less than 10 days before slaughter for export. SHEEP – DO NOT USE less than 7 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1 800 226 511 or the APVMA website [www.apvma.gov.au/residues/](http://www.apvma.gov.au/residues/)

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues: adequate treatment records should be kept and made available, if requested by wool buyers.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT**

Do not contaminate dams, ponds, waterways or drains with **COOPERS STAMPEDE**.

File Name: INTERVET - 61351_0614 - Stampede-Diflubenzuron.docx	Prepared By: J. Burgess	Label & Printer Specifications:	Version: 4	Page 14 of 15
Preparation Date: 9/12/2008				
Last Revised: 9 May 2014				