

Product Name: Apparent Triceratops 700 WG Herbicide  
APVMA Approval No: 85750 / 123509



Label Name:	Apparent Triceratops 700 WG Herbicide
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	700 g/kg IMAZAMOX
Mode of Action:	GROUP <b>B</b> HERBICIDE
Statement of Claims:	For the post-emergence control of certain annual grass and broadleafed weeds in field peas, legume-based pastures, lucerne, peanuts and soybeans, as specified in the DIRECTIONS FOR USE table  IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USE
Net Contents:	250g-3kg
Restrains:	DO NOT apply to crops or weeds stressed by factors such as root or foliar diseases, water logging, nutrient deficiencies or extremes of temperature and moisture. DO NOT apply to crops stressed by any previous herbicide treatments.
Directions for Use:	This section contains file attachment.
Other Limitations:	
Withholding Periods:	GRAZING

	<p>Lucerne, pastures: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 7 DAYS AFTER APPLICATION.</p> <p>Peanuts, soybeans: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.</p> <p>Field peas: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.</p> <p>HARVEST</p> <p>Field peas, peanuts, soybeans: NOT REQUIRED WHEN USED AS DIRECTED.</p>
--	--

Trade Advice:	
---------------	--

General Instructions:	
-----------------------	--

Resistance Warning:	<p><b>RESISTANT WEEDS WARNING</b>  <b>GROUP B HERBICIDE</b></p> <p>Apparent Triceratops 700 WG Herbicide is a member of the Imidazolinone group of herbicides. The product has the ALS mode of action. For weed resistance management, the product is a Group B herbicide. Some naturally-occurring weed biotypes resistant to the product and other Group B herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group B herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Apparent Pty Ltd accepts no liability for any losses that may result from failure of this product to control resistant weeds.</p>
---------------------	---

Precautions:	
--------------	--

Protections:	<p><b>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</b>  DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.  DO NOT spray within 50 m of wetlands or waterways.</p> <p>Crop safety: This product may cause shortening of plant internodes and may in some circumstances lead to transient crop yellowing but plants will soon recover. This effect is more pronounced under poor growth conditions including conditions of prolonged moisture stress.</p> <p>Field peas  DO NOT apply to field peas beyond the 4 node stage.  DO NOT apply if frost is forecast.  DO NOT use Apparent Triceratops 700 WG Herbicide after simazine (unregistered in field peas) and DO NOT exceed 250 g ac/ha diuron if use of Apparent Triceratops 700 WG Herbicide is planned.  DO NOT roll field peas after germination if use of Apparent Triceratops 700 WG Herbicide is planned.  Before using this product on field pea varieties other than Bohatyr, Bonzer, Dun, Dundale, Glenroy, Laura, Magnet, Mukta, Santi and Wirrega, contact TGAC Australia Pty Ltd for upto-date information.  DO NOT use on Alma, Excell and Parafield varieties.</p> <p>Pasture legumes and perennial grasses  Safety to medic varieties may be lower than to clover varieties. Safety to seedling pastures less than one year old may be lower than to older established pastures. Transient biomass</p>
--------------	---

	<p>reduction and leaf yellowing may occur to pasture legumes and perennial grasses, with good recovery occurring under favourable seasonal conditions. In some situations a reduction in perennial grass numbers may occur as a result of unfavourable seasonal conditions following application. For up-to-date information on all pasture clover and medic varieties and other temperate pasture species, contact Apparent Pty Ltd before use.</p> <p>Lucerne Before using this product on lucerne varieties other than Hunterfield, Sceptre, Sequel, Siriver and Trifecta, contact Apparent Pty Ltd for up-to-date information.</p> <p><b>DO NOT USE THE HASTEN® OR KWICKIN® MIX ON SEEDLING LUCERNE.</b></p> <p>Replanting: If replanting is necessary in a field previously treated with Apparent Triceratops 700 WG Herbicide, the field may be replanted to field peas (above listed varieties only), soybeans or peanuts. DO NOT apply a second treatment of Apparent Triceratops 700 WG Herbicide.</p> <p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b></p> <p><b>TOXIC TO AQUATIC FLORA. DO NOT</b> contaminate streams, rivers or waterways with the chemical or used containers.</p>
--	---

Storage and Disposal:	Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
-----------------------	---

Safety Directions:	Will irritate the eyes. When opening the container and preparing spray, wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash face shield or goggles.
--------------------	---

First Aid Instructions:	<p><b>FIRST AID</b></p> <p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

## DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	RATE PER ha	WHP	CRITICAL COMMENTS
<p><b>Field peas</b>  <b>Post-emergence</b>            See "Crop Safety" re varieties</p>	<p>Barley grass (<i>Hordeum leporinum</i>), Brome grass (<i>Bromus</i> spp.), Deadnettle (<i>Lamium amplexicaule</i>), Indian hedge mustard (<i>Sisymbrium orientale</i>), Storksbill (<i>Erodium</i> spp.), Turnip weed (<i>Rapistrum rugosum</i>), Volunteer barley (<i>Hordeum vulgare</i>), Volunteer oats (<i>Avena sativa</i>), Volunteer lupins (<i>Lupinus</i> spp.), Volunteer triticale (<i>Triticosecale</i> spp.), Volunteer wheat (<i>Triticum aestivum</i>), Wild oats (<i>Avena</i> spp.), Wild turnip (<i>Brassica tournefortii</i>), *Doublegee (<i>Emex australis</i>), *Shepherds purse (<i>Capsella bursa-pastoris</i>), *Threehorn bedstraw <i>Galium tricornutum</i>, *Wild radish (<i>Raphanus raphanistrum</i>), *Wireweed (<i>Polygonum aviculare</i>)</p>	<p><b>45g plus</b> BS 1000, or equivalent, at 200mL per 100L water</p>	<p>6 weeks (grazing)</p>	<p><b>DO NOT</b> apply to field peas beyond the 4 node stage.  <b>DO NOT</b> use adjuvants other than BS 1000 or equivalent.</p> <p>Apply to actively growing broadleaved weeds in the cotyledon to 3 leaf stage. Apply to grass weeds up to the 2 tiller stage.            Good crop growth will aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded.            * Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds.            Leaf yellowing and height suppression may occur, especially under adverse growing conditions such as moisture stress. Flowering may be delayed.</p> <p><b>IMPORTANT: Refer to CROP SAFETY section of label re use on field peas.</b>            Refer to <b>FOLLOW CROPS</b> section of this label regarding follow crops.</p>

CROP	WEEDS CONTROLLED	RATE PER ha	WHP	CRITICAL COMMENTS
<b>Legume-based pastures Post-emergence clovers, lucerne, medics, serradellas only;</b> see "Crop Safety" re varieties	As for field peas above	<b>45g plus</b> Hasten® or Kwickin® at 500mL per 100L water	7 days (grazing)	<p><b>Seedling</b> Apply to legumes after full emergence of the third trifoliate leaf but well before commencement of flowering.</p> <p><b>Established</b> Apply before commencement of flowering. Apply to actively growing broadleaved weeds in the cotyledon to 3 leaf stage. Apply to grass weeds up to the 2 tiller stage. Good pasture growth will aid weed control. Weeds may not be totally controlled but populations will be significantly reduced and surviving plants will generally be severely retarded. Where pasture stand is poor, weed control may be inadequate due to lack of competition. Ensure sufficient spray coverage of weeds particularly in dense pastures. * Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Very high populations may not be efficiently controlled. Good spray coverage is essential. Ensure that crop agronomy is correct, to provide competition to such weeds.</p> <p><b>Refer to CROP SAFETY section</b> of this label re varieties of clovers and medics and other pasture species.</p> <p><b>Note: Use of the 50g/ha rate plus Hasten® or Kwickin® may severely damage medics.</b> Refer to <b>FOLLOW CROPS</b> section of this label regarding follow crops.</p>
	*Silver grass ( <i>Vulpia</i> spp.), plus above weeds	<b>50g plus</b> Hasten® or Kwickin®, at 500mL per 100L water plus Liase® at 2L per 100L water		
<b>Lucerne Post-emergence</b> See "Crop Safety" re varieties	As for post emergence use in field peas (winter weeds) and soybeans (summer weeds)	<b>45 or 50g plus</b> BS 1000®, or equivalent, at 200mL per 100L water plus Liase® at 2L per 100L water	7 days (grazing)	<p>Apply to actively growing weeds at growth stages as for field peas (winter weeds) and soybeans (summer weeds).</p> <p><b>DO NOT</b> use the Hasten® or Kwickin® mix on seedling lucerne. Use the 50g rate for summer weeds.</p> <p><b>Crop stage:</b> Seedling: apply when crop is at 2 trifoliate leaf stage and before commencement of flowering. Established: apply as above following cutting or grazing.</p> <p><b>Refer to CROP SAFETY section</b> of this label re varieties of lucerne.</p>

CROP	WEEDS CONTROLLED	RATE PER ha	WHP	CRITICAL COMMENTS
<b>Post-emergence</b> Peanuts, soybeans	Amaranth ( <i>Amaranthus</i> spp.), Barley grass ( <i>Hordeum leporinum</i> ), Barnyard grass ( <i>Echinochloa crus-galli</i> ), Bell vine ( <i>Ipomoea plebeia</i> ), Brome grass ( <i>Bromus</i> spp.), Deadnettle ( <i>Lamium amplexicaule</i> ), Fathen ( <i>Chenopodium album</i> ), Fierce thornapple ( <i>Datura ferox</i> ), Indian hedge mustard ( <i>Sisymbrium orientale</i> ), Liverseed grass ( <i>Urochloa panicoides</i> ), Turnip weed ( <i>Rapistrum rugosum</i> ), Volunteer barley ( <i>Hordeum vulgare</i> ), Volunteer oats ( <i>Avena sativa</i> ), Volunteer lupins ( <i>Lupinus</i> spp.), Volunteer triticale ( <i>Triticosecale</i> spp.), Volunteer wheat ( <i>Triticum aestivum</i> ), Wild gooseberry ( <i>Physalis minima</i> ), Wild oats ( <i>Avena</i> spp.), Wild turnip ( <i>Brassica tournefortii</i> ), *Anoda weed ( <i>Anoda cristata</i> ), *Awnless barnyard grass ( <i>Echinochloa colona</i> ), *Blackberry nightshade ( <i>Solanum nigrum</i> ), *Caltrop ( <i>Tribulus terrestris</i> ), *Chickweed ( <i>Stellaria media</i> ), *Crabgrass ( <i>Digitaria ciliaris</i> ), *Doublegee ( <i>Emex australis</i> ), *Noogoora burr ( <i>Xanthium pungens</i> ), *Shepherd's purse ( <i>Capsella bursa-pastoris</i> ), *Threehorn bedstraw ( <i>Galium tricornutum</i> ), *Wild radish ( <i>Raphanus raphanistrum</i> ), *Wireweed ( <i>Polygonum aviculare</i> )	<b>50g plus EITHER BS 1000<sup>®</sup>, (or equivalent), at 200mL per 100L water plus Liase<sup>®</sup> at 2L per 100L water OR Hasten<sup>®</sup> or Kwickin<sup>®</sup>, at 500mL or 1L per 100L water</b>	4 weeks (grazing)	DO NOT apply to crops beyond the 6 leaf stage.  Apply to actively growing weeds in the cotyledon to 2 leaf stage. Apply to grass weeds up to the 2 tiller stage.  If using the Hasten <sup>®</sup> or Kwickin <sup>®</sup> mix, use the 1L per 100L rate where grass weeds are predominate.  * Control will not be 100% but surviving plants will generally be retarded and will not compete with good crop growth. Ensure that crop agronomy is correct, to provide competition to such weeds.  <b>Refer to FOLLOW CROPS section</b> of this label regarding follow crops.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

## GENERAL INSTRUCTIONS

This product is for early post-emergence weed control in field peas, lucerne, legume-based pastures, soybeans and peanuts. Weeds will either die or remain stunted and will not compete with the crop. For weeds marked with an asterisk in the Directions for Use table, the level of control will vary significantly depending on factors such as climatic conditions following application, crop vigour, weed seed depth, etc.

A spray adjuvant must always be added, according to the Directions for Use table, while Liase® Liquid Herbicide Adjuvant should also be added in situations indicated in the table. Good crop competition is essential for effective weed control.

## MIXING

**Apparent Triceratops 700 WG Herbicide** is a water dispersible granule formulation. Part fill the spray tank with water, then with the agitator running, add the required amount of product, then fill the tank with water. When tank mixing this product with other recommended compatible products, first add the other product to the tank and mix thoroughly before adding this product.

## APPLICATION

For ground application only: Apply with boom equipment in not less than 50L/ha water using flat fan nozzles. Avoid overlap and do not over spray headlands.

**DO NOT** apply by aircraft.

**DO NOT** apply by mister.

**Apparent Triceratops 700 WG Herbicide** should be applied a minimum of two hours before rainfall or irrigation.

**DO NOT** apply **Apparent Triceratops 700 WG Herbicide** more than once per growing season.

## EQUIPMENT CLEAN-UP

Thoroughly flush all spray equipment with water following the use of **Apparent Triceratops 700 WG Herbicide** and before use with other products. Rinse water should NOT be discharged where it will reach streams, water bodies or natural vegetation.

## COMPATIBILITY

The product is compatible with MCPA, simazine, dimethoate, Fastac® Duo, Crop Care Dominex® Duo, phosmet, omethoate and endosulfan.

**DO NOT** tank mix with selective post-emergence grass herbicides. **DO NOT** apply these herbicides following the use of **Apparent Triceratops 700 WG Herbicide** until grasses have resumed active growth. Alternatively, apply the grass herbicide at least one day before using **Apparent Triceratops 700 WG Herbicide**.

**DO NOT** tank mix with either MCPA Amine or 2,4-D Amine when using in pastures.

## FOLLOW CROPS

Under conditions, such as very dry seasons, which do not favour breakdown of this product, carry-over soil residues can affect susceptible follow crops. As environmental and agronomic factors make it impossible to eliminate all risks associated with this product, rotational crop injury is always possible. The following minimum re-cropping intervals (months after application) should be observed.

**Following use in lucerne, legume-based pastures and field peas:**

MONTHS AFTER APPLICATION		
0	10	21
field peas, canola varieties tolerant to applications of imazamox, maize varieties tolerant to applications of imazamox, wheat varieties tolerant to applications of imazamox	chickpeas faba beans lucerne lupins pasture legumes vetch *barley *wheat (except varieties tolerant to applications of imazamox) *triticale	All other crops including canola (except varieties of canola tolerant to applications of imazamox) oats, safflower

\* The following additional requirements apply if it is intended to sow WHEAT, BARLEY or TRITICALE during the next winter season:

**DO NOT** apply **Apparent Triceratops 700 WG Herbicide** later than the end of August.

**DO NOT** use **Apparent Triceratops 700 WG Herbicide** in areas where rainfall from the time of spraying to sowing of cereals is expected to be below 200mm.

**Furthermore:**

**DO NOT** use on soils of pH 5.5 (CaCl<sub>2</sub>) or less in areas where rainfall from spraying to sowing of cereals is expected to be below 300mm.

In late, short seasons where the soil is cold for most of the time it is wet, break-down will be slower and plant-back times will be extended.

If expected rainfall is not received following use of **Apparent Triceratops 700 WG Herbicide**, consult your local agronomist before planting wheat, barley or triticale. (In calculating rainfall actually received, exclude single, isolated falls which do not result in periods of continuous soil moisture to allow microbial breakdown to occur in the root zone).

**Following use in summer crops:**

**Irrigated only:**

MONTHS AFTER APPLICATION			
0	5	10	18
mungbeans peanuts soybeans canola varieties tolerant to applications of imazamox maize varieties tolerant to applications of imazamox wheat varieties tolerant to applications of imazamox	*chickpeas *lucerne *lupins *pasture legumes *barley *wheat (except varieties tolerant to applications of imazamox)	**maize (except varieties tolerant to applications of imazamox) **sorghum **cotton **oats **sunflower	All other crops (providing rainfall and irrigation exceeds 2000mm)

\***DO NOT** plant these crops unless interim rainfall (rainfall plus irrigation) from application to sowing is at least 500mm.

\*\***DO NOT** plant these crops unless interim moisture (rainfall plus irrigation) from application to sowing is at least 800mm.

**Dryland only:**

<b>MONTHS AFTER APPLICATION</b>				
<b>0</b>	<b>8</b>	<b>10</b>	<b>15</b>	<b>22</b>
mungbeans peanuts soybeans canola varieties tolerant to applications of imazamox maize varieties tolerant to applications of imazamox wheat varieties tolerant to applications of imazamox	*lucerne *barley *wheat (except varieties tolerant to applications of imazamox)	**chickpeas **maize (except varieties tolerant to applications of imazamox) **sorghum **cotton **oats **sunflower	pasture legumes lupins	All other crops (providing rainfall and irrigation exceeds 2000mm)

**\*DO NOT** plant these crops unless interim rainfall from application to sowing is at least 500mm.

**\*\*DO NOT** plant these crops unless interim rainfall from application to sowing is at least 800mm.

