



Product Name: ECO-OIL ORGANIC MITICIDE/INSECTICIDE BOTANICAL OIL CONCENTRATE
APVMA Approval No: 51403/113438



| | |
|-------------------------|--|
| Label Name: | ECO-OIL ORGANIC MITICIDE/INSECTICIDE BOTANICAL OIL CONCENTRATE |
| Signal Headings: | |
| Constituent Statements: | 850 g/L EMULSIFIABLE BOTANICAL OIL |
| Statement of Claims: | CONTROLS CITRUS LEAFMINER, SCALE AND VARIOUS SAP SUCKING PESTS IN CERTAIN VEGETABLE AND ORNAMENTAL PLANTS IN THE HOME GARDEN IMPROVES WETTING AND STICKING OF SPRAYS |
| Net Contents: | 250mL 500mL 1L 5L 20L |
| Restrains: | DO NOT spray Eco-Oil in temperatures exceeding 35°C. DO NOT spray directly on to sensitive flower heads, apply to foliage only to prevent oil spotting. DO NOT apply to plants suffering from heat or moisture stress. |
| How to Use: | This section contains file attachment. |
| Withholding Periods: | Not required when used as directed |
| How to Prepare: | HOW TO PREPARE Eco-Oil is a botanical oil based miticide/insecticide that controls pests by contact. As such complete coverage of plants is important. For best results spray with Eco-Oil at the first sign of pest infestation. Repeat application may be necessary. Some plant species (like maidenhair ferns) are more sensitive than others so it is recommended that a pre-test be |

conducted on a small area of foliage before using. Do not exceed recommended dose rates.

How to mix

1. Half fill the spray container with clean water.
2. Add the required amount of Eco-Oil, shake, then top up container with water and shake again.

Frequent agitation (shaking) of the spray solution is recommended during application.

Precautions:

CAUTION

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

Protections:

Storage and Disposal:

Store in the closed, original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in plastic bag and putting in garbage.

Safety Directions:

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.

First Aid Warnings:

| INSECTICIDE USE | | | |
|---|---|-------------------------------------|--|
| PLANTS | PESTS | RATE | HOW TO APPLY |
| Tomatoes, cucumbers, capsicums, strawberries and ornamental plants | Two-spotted mite Aphids Sooty mould | Mix 5 mL of product to 1 L of water | Spray when pests first appear. Apply two sprays 3 to 5 days apart. Repeat applications at signs of reinfestation. Spray both upper and underside of leaves to run-off. DO NOT apply more than three sprays to plants within a 4 to 8 week period. |
| Vegetables and ornamental plants | Greenhouse Whitefly | | |
| Ornamental plants | Mealybugs Sooty mould | 10 mL of product to 1 L of water | Apply at early signs of Mealybugs and Scale. Apply two sprays at 7 day intervals. Repeat applications at signs of reinfestation. Spray upper and underside of leaves and branches to run-off. |
| Olive trees | Black scale | | |
| Citrus | Citrus leafminer | 10 mL of product to 1 L of water | Begin applying when the new flush growth is about 4 cm long. Apply every 5 to 14 days with presence of new flush growth. Ensure that the new flush leaves are sprayed both on top and underneath. Do not apply any more than 8 times per season. |
| SPRAY ADDITIVE USE – TO IMPROVE WETTING AND STICKING OF SPRAYS | | | |
| Spray additive for insecticides, fungicides and foliar fertilisers | - | 5 mL per 2 litres of spray mix | Rate is given as a guide only. Always follow directions on pesticide label where addition of a botanical oil concentrate is recommended. |

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.