



**Company Name:** AXICHEM PTY LTD  
**Product Name:** AC Amon Herbicide  
**APVMA Approval No:** 80927/101955



<b>Label Name:</b>	AC Amon Herbicide
<b>Signal Headings:</b>	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
<b>Constituent Statements:</b>	250g/L AMITROL 220g/L AMMONIUM THIOCYANATE
<b>Mode of Action:</b>	GROUP <b>Q</b> HERBICIDE
<b>Statement of Claims:</b>	For the control of weeds in orchards, vineyards, irrigation ditches and drains, roadsides, wheat and barley, and for general industrial situations.
<b>Net Contents:</b>	5L 10L 20L - 1000L
<b>Restrains:</b>	
<b>Directions for Use:</b>	This section contains file attachment. <b>File Name:</b> 28052015 DIRECTIONS FOR USE.pdf <b>File Size:</b> 63277 bytes
<b>Other Limitations:</b>	
<b>Withholding Periods:</b>	ORCHARDS AND VINES: DO NOT APPLY LATER THAN 56 DAYS BEFORE HARVEST OTHER CROPS: NOT REQUIRED WHEN USED AS DIRECTED.

<b>Trade Advice:</b>	
----------------------	--

<b>General Instructions:</b>	<p>This section contains file attachment.  <b>File Name:</b> APPLICATION INFORMATION.pdf  <b>File Size:</b> 48890 bytes</p>
------------------------------	---

<b>Resistance Warning:</b>	<p><b>RESISTANT WEEDS WARNING</b></p> <p><b>GROUP Q HERBICIDE</b></p> <p>AC Amon Herbicide is a member of the triazole group of herbicides. AC Amon Herbicide has the inhibitors of carotenoid biosynthesis mode of action. For weed resistance management AC Amon Herbicide is a Group Q herbicide. Some naturally occurring weed biotypes resistant to AC Amon Herbicide and other Group Q herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by AC Amon Herbicide or other Group Q herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Axichem Pty Limited accepts no liability for any losses that may result from the failure of AC Amon Herbicide to control resistant weeds. AC Amon Herbicide may be subject to specific resistance strategies. For further information contact your farm chemical supplier, consultant, local Department of Agriculture or local Axichem Pty Ltd representative.</p>
----------------------------	---

<b>Precautions:</b>	
---------------------	--

<b>Protections:</b>	<p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT</b>  DO NOT contaminate streams, rivers or waterways with the chemical or used containers.</p> <p><b>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</b>  Drift Warning: This product is a non-selective herbicide and may damage crops, pastures and some ornamentals. However it is safer to use near plants susceptible to phenoxy type herbicides. DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.</p> <p><b>Equipment Maintenance and Usage:</b>  Do not use the spray machine for any other purpose unless thoroughly cleaned by several rinsings of all parts with clean water.</p>
---------------------	---

<b>Storage and Disposal:</b>	<p><b>STORAGE AND DISPOSAL</b></p> <p>Triple or preferably pressure rinse containers before disposal. Add rinsings to tank. Do not dispose of chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or territory government regulations. DO NOT burn empty containers or product.</p> <p>Refillable Containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.</p>
------------------------------	---

<b>Safety Directions:</b>	<b>SAFETY DIRECTIONS</b> DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.
---------------------------	---

<b>First Aid Instructions:</b>	<b>FIRST AID</b> If poisoning occurs contact a doctor or Poisons Information Centre (Phone Australia 13 11 26).
--------------------------------	--

<b>First Aid Warnings:</b>	
----------------------------	--

## DIRECTIONS FOR USE

SITUATION AND CROP	WEEDS CONTROLLED	STATE	Rate/Ha	CRITICAL COMMENTS
Wheat and Barley	Rye Grass, Wild Oats	All States	2.8L	Pre-planting. Sow the crop at a depth of 5cm not less than 5 days after spraying. A mechanical cultivation 4 to 5 days after spraying will assist results.  Apply when grasses are in the 2-3 leaf stage and 5cm high.
			5.6L	Apply when grasses are in the 6 leaf stage and 10cm high. DO NOT use on sandy soils in cropping situations.
Vineyards and Orchards	Couch Grass		1.1L/100L water	Apply when weeds are small and actively growing. Repeat application in 6-8 weeks. Apply as a directed spray to weeds only.
	Broadleaf Weeds and Grasses		570mL/100L water	
Potatoes Pre-Harvest Preparation	Rye Grass, Winter Grass, Capeweed, Paterson's Curse	Vic, Tas only	5.5-11L	Apply approximately 4-5 weeks before harvest after the potato haulms have dried off. Use the higher rates when rye grass is in clumps or other weeds more than 30cm high are present. For boom spraying apply at least 100litres of spray mixture per hectare. If other broadleaf weeds such as Clovers, Variegated Thistle and Cruciferous weeds are also present add a Low Volatile Ester 400 Herbicide. <b>Warning: if used on sandy soils do not plant follow-up crops within 3 months.</b>
<i>Pinus radiata</i> Plantations	Rye Grass, Wild Oats, Yorkshire Fog Grass, Silver Grass, Sand Brome, Capeweed, Clovers, Mouse-ear Chickweed, Sow Thistle and Spear Thistle seedlings	NSW, SA, WA, Tas, Vic only	5.6-8L/ha plus 2.8-4L/ha flowable Atrazine 500g/L	Use highest rates where grasses are a problem. Spray evenly. <b>Pre-planting:</b> Graze area heavily and apply the tank mixture not more than 2 weeks before planting. <b>Post-planting:</b> Apply tank mixture immediately after or within one month of planting. Do not spray pine seedlings using directed spray.
Non-crop areas around buildings, Commercial and Industrial areas, Domestic and Public Service areas, Right-Of-Ways	Blackberry	Vic, Tas, SA only	2L/100L water	Apply twice. First treatment in December or January and regrowth sprayed in autumn. Respraying in the following year is essential. DO NOT apply to blackberry bushes carrying mature or developing fruit.
	Furze or Gorse, Ox-eye Daisy			Apply during summer or before full flowering. Respraying will be necessary.
	Fennel, Hoary Cress, African Daisy, Apple of Sodom, Illyrian Thistle	All States	1.1L/100L	Apply when weeds are actively growing, immediately prior to flowering.
	Russian (creeping) Knapweed, Watsonia, Artichoke Thistle, Prairie Ground Cherry, Onion Weed, Khaki Weed, Chilean Cestrum (Green Cestrum), Oxalis (Soursob), Couch, Kikuyu, Fog Grass, Sorrel			Apply when weeds are actively growing, immediately prior to flowering. Respraying will be necessary to destroy regrowth and seedlings.
Aquatic Areas, Drains and Channels, Margins of Streams, Lakes and Dams	Cumbungi, Phragmites and Nutgrass		2.3L/100L water	Apply during flowering period between January and May. Re-treatment of Nutgrass will be necessary.
	Water Couch		4.5L/100L water	Apply as a single treatment between February and May.
			2.2L/100L water	Apply in November to December with a further application at the same rate 6-8 weeks later.
	Water Hyacinth		280mL/100L water	Apply immediately prior to flowering.

**NOT TO BE USES FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**Before opening carefully read Safety Directions, First Aid Instructions, Directions for Use, Storage and Disposal and Protection Statements.**

### **APPLICATION INFORMATION**

Thoroughly wet all surfaces by aiming to apply 1700-2200 Litres per hectare with power spraying equipment, 900-1100 Litres per hectare (1 Litre per 10m<sup>2</sup>) using a knapsack and 170 Litres per hectare when using a misting machine.

**Note: Misting machines are not recommended for use in Tasmania.**

### **MIXING INSTRUCTIONS**

To improve the efficiency of this product it is recommended that a 700g/L active surfactant be added to the spray mixture at the following rates:

Power Spraying – 250mL/100L spray mixture

Knapsack Spraying – 55mL/15L knapsack

Misting Machine – 60mL/15L spray mixture (Not recommended in Tasmania)

Note: The rates indicated in the Directions for Use table are per 100L of water for power spraying. Refer to the chart below for the required amounts to suit other equipment.

### **CONVERSION CHART FOR RECOMMENDATIONS TO SUIT VARIOUS EQUIPMENT**

<b>Recommended Rate per 100L of water</b>	<b>Per 15L of knapsack</b>	<b>Per 5L Misting</b>
4.5L	1.1L	1.3L
2.2L	570mL	630mL
2.0L	510mL	560mL
1.7L	430mL	480mL
1.1L	280mL	310mL
570mL	110mL	120mL
280mL	55mL	60mL