

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY**

FENNEC™
POUR-ON FOR SHEEP

**Active Constituents: 35 mg/mL IMIDACLOPRID, 4 mg/mL ABAMECTIN
Solvent: 300 mg/mL N-methyl-2-pyrrolidone**

For the treatment and control of susceptible strains of lice (*Bovicola ovis*) on sheep, including synthetic pyrethroid resistant lice when applied up to 7 days after shearing. Sheep will be protected against re-infestation with lice for 5 weeks after treatment when applied within 24 hours of shearing. For the treatment and control of abamectin susceptible gastrointestinal nematodes of sheep.



20L



**Boehringer
Ingelheim**

LF18400

Fennec™ Pour-On for Sheep is effective for the treatment and control of the following abamectin sensitive adult and immature gastrointestinal nematodes:
 Small Brown Stomach Worm (*Teladorsagia circumcincta*)
 Black Scour Worm (*Trichostrongylus* spp.)
 Stomach Hair Worm (*Trichostrongylus axei*)
 Nodule Worm (*Oesophagostomum columbianum*)
 Large Bowel Worm (*Oesophagostomum venulosum*)

Fennec™ Pour-On for Sheep contains imidacloprid, a member of the neonicotinoid family of chemicals and abamectin, a member of the macrocyclic lactone family of chemicals. Resistance may develop to any chemical.

DIRECTIONS FOR USE

Restraints

DO NOT USE on ewes which are producing, or may in the future produce milk that may be used or processed for human consumption.

Contraindications

DO NOT USE on unshorn sheep, or shorn sheep more than 7 days after shearing. DO NOT USE on ewes with unshorn lambs at foot.

This product should not be used on lambs under 6 weeks of age or less than 10 kg bodyweight.

This product should not be used on severely sick, debilitated or dying animals. This product should not be used on sheep which have not been cleanly shorn for any reason e.g. Mycotic Dermatitis.

Precautions

This product is for application as a pour-on only. Do not give orally or by injection. Do not apply this product at the same time as any other pour-on products.

Dosage and Administration

Dose rate: 10 mL for each 10 kg body weight.

Sheep should be weighed before treatment. Apply the entire dose to sheep under 30 kg as a single broadband stripe from poll of the head to the base of the tail.

For sheep over 30 kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the base of the tail. The stripes should be no further than 10 cm apart. Refer to diagrams below.

The following table may be used as a guide:

| Bodyweight (kg) | Dose Volume (mL) | Method | Apply in the pattern below |
|-----------------|------------------|---------------|----------------------------|
| 10 to 12.5 | 12.5 | Single stripe | |
| 12.6 to 15 | 15 | | |
| 15.1 to 20 | 20 | | |
| 20.1 to 25 | 25 | | |
| 25.1 to 30 | 30 | | |
| 30.1 to 40 | 40 | Double stripe | |
| 40.1 to 50 | 50 | | |
| 50.1 to 60 | 60 | | |
| 60.1 to 70 | 70 | | |
| 70.1 to 80 | 80 | | |
| 80.1 to 90 | 90 | | |

Animals heavier than 90 kg to be dosed at 10 mL per 10 kg. A representative sample of animals should be weighed before treatment. Dose the mob according to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs).

Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range.

Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

General Directions

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.

Apply product only with the recommended applicator fitted with a six-hole T-bar nozzle. Check the accuracy of equipment before and during treatment.

Efficacy is not affected by heavy rainfall before and/or after treatment.

Avoid treating ewes less than 6 weeks before lambing commences, while ewes are lambing or have lambs at foot, as live lice can persist and infect the lambs.

Treated sheep should not be mixed with lice free sheep until 6 weeks after treatment.

Management practices which promote good lice control such as complete musters, accurate treatment of every animal according to label, ensuring all animals can be treated at the same time, no split shearings, stock-proof fencing to prevent external introductions, quarantine and lice treatment of all newly purchased stock are essential for effective lice control.

Fennec Pour-On for Sheep contains a blue dye to identify treated sheep. This dye is scorable from wool.

Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted before any parasite treatment is used.

Do not leave container, hose or applicator in the sun for long periods of time.

The product can be detrimental to certain plastics, paints and surface finishes. Avoid contact with such surfaces.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 63 days before slaughter for human consumption.

MILK: DO NOT USE on ewes which are producing, or may in the future produce milk that may be used or processed for human consumption.

WOOL: DO NOT USE less than 6 months before shearing or fibre collection.

Use of this product may result in wool residues that may not comply with European Union environmental quality standards.

Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 84 days before slaughter for export. Before using this product, confirm the current ESI from Boehringer Ingelheim on 1800-808-691 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes. When using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow length chemical resistant gloves and a face shield or goggles. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

Rehandling Interval: DO NOT handle sheep after treatment until dry.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126. If swallowed, do NOT induce vomiting. If in eyes wash out immediately with water.

ADDITIONAL USER SAFETY INFORMATION

Additional information is listed in the Safety Data Sheet.

ENVIRONMENTAL PROTECTION

Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple rinse or pressure rinse container before disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

STORAGE

Store below 30°C (Room Temperature). Protect from light.

Additional GHS statements required by Safe Work Australia: Combustible liquid. Causes skin irritation. Harmful if inhaled. May cause respiratory irritation. May damage fertility or the unborn child. Keep away from heat, sparks, open flames and hot surfaces – No smoking. Avoid breathing fumes, mists, vapours or spray. Do not get on skin or on clothing. Do not eat, drink or smoke when using this product. Use only outdoors or in a well ventilated area. If swallowed: call a Poison Centre or doctor if you feel unwell. If on skin: wash with plenty of soap and water. If inhaled: remove victim to fresh air and keep at rest in a position comfortable for breathing. If exposed or concerned, if skin irritation occurs or if eye irritation persists: get medical advice. Store in a closed container, locked up in a dry, well-ventilated place. Keep cool.

Boehringer Ingelheim Animal Health Australia Pty Ltd
 Level 1, 78 Waterloo Road,
 North Ryde NSW 2113

For more information, contact Boehringer Ingelheim Customer Care on 1800 808 691.

APVMA Approval No.: 85544/113561

™Fennec is a trademark of the Boehringer Ingelheim Group.

Batch:

Expiry:



LB18414