



Company Name: SPECIALIST SALES PTY LTD
Product Name: TICKPRO POUR-ON TICK DEVELOPMENT INHIBITOR
APVMA Approval No: 81640/103523

Label Name:	TICKPRO POUR-ON TICK DEVELOPMENT INHIBITOR
--------------------	--

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
-------------------------	--

Constituent Statements:	25g/L FLUAZURON Also Contains: (Solvents) 100 g/L 1-dodecyl-2-pyrrolidone 100 g/L 1-octyl-2-pyrrolidone 415 g/L 1-methyl-2-pyrrolidone
--------------------------------	--

Claims:	For strategic control of the Cattle Tick, Rhipicephalus (Boophilus) microplus (including all known acaricide resistant strains), on Beef Cattle.
----------------	--

Net Contents:	2.5L (5L, 10L, 20L, 25L)
----------------------	--------------------------

Directions for Use:	DIRECTIONS FOR USE:
----------------------------	---------------------

Restrains:	Restraint: DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.
-------------------	---

Contraindications:	Contraindications This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.
---------------------------	---

Precautions:	
---------------------	--

Side Effects:	
----------------------	--

Dosage and Administration:	<p>Dosage and Administration</p> <p>2.5 L [5 L, 10 L, 20 L, 25 L] Pack [Refer to the TickPro Dosage Guide attached]</p> <p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION</p> <p>This section contains file attachment. File Name: eLabel_TickPro Dosage Guide Tables_V2_9 Oct 15.pdf File Size: 44588 bytes</p>
-----------------------------------	--

General Directions:	<p>General Directions: TICKPRO POUR-ON Treated Areas – Introduction of New Cattle Tick infested cattle introduced onto properties where a TICKPRO POUR-ON treatment regime is in place should be treated with TICKPRO POUR-ON prior to release.</p> <p>Handling of Treated Animals Wear protective clothing recommended under Safety Directions, if handling treated cattle within 24 hours of application.</p> <p>For local tick control recommendations contact your local department of agriculture.</p> <p>Cleaning Instructions After use wash TICKPRO POUR-ON applicator with detergent and hot water. Dispose of the rinsates in a disposal pit away from desirable plants and their roots and watercourses.</p>
----------------------------	---

Withholding Periods:	<p>WITHHOLDING PERIODS: MEAT: DO NOT USE less than 42 days before slaughter for human consumption. Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows. MILK: DO NOT USE on cattle which are producing or may in the future produce milk that may be used or processed for human consumption.</p>
-----------------------------	---

Trade Advice:	<p>TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Specialist Sales on 1800 780 317 or the APVMA website (www.apvma.gov.au/residues).</p>
----------------------	--

Safety Directions:	<p>SAFETY DIRECTIONS Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking and smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.</p>
---------------------------	--

First Aid Instructions:	<p>FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.</p>
--------------------------------	--

First Aid Warnings:	
----------------------------	--

Additional User Safety:	ADDITIONAL USER SAFETY INFORMATION: For further information, refer to the Safety Data Sheet.
--------------------------------	--

Environmental Statements:	ENVIRONMENTAL PROTECTION DO NOT contaminate fish ponds, dams or rivers with the product or used containers. This product is very toxic to aquatic invertebrates.
----------------------------------	---

Disposal:	DISPOSAL: Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, and bury empty containers in a local authority landfill. If no landfill is available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
------------------	---

Storage:	STORAGE: Store below 30°C (Room Temperature). Store in original container tightly closed in a dry, cool place. Store out of direct sunlight.
-----------------	---

TICKPRO DOSAGE GUIDE:

2.5L PACK

Animals and Pest	State	Bodyweight (kg)	Dose (mL)	No. treated per 2.5 L pack	Critical comments	
<u>Beef cattle after weaning</u> Cattle Tick (<i>Rhipicephalus microplus</i>)	QLD, NSW, WA & NT only	101 – 150	9	277	<ol style="list-style-type: none"> Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. Maximum number of treatments per annual tick season: 3. Apply in two bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator. The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose. There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves. TICKPRO POUR-ON should not be applied to show cattle as a light scurfiness may be visible for some time after treatment. TICKPRO POUR-ON is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of TICKPRO POUR-ON commences within 3 days of treatment. Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas. 	Cattle application diagram as per reference product
		151 – 200	12	208		
		201 – 250	15	166		
		251 – 300	18	138		
		301 – 350	21	119		
		351 – 400	24	104		
		401 – 450	27	92		
		451 – 500	30	83		
		501 – 550	33	75		
		551 – 600	36	69		
601 – 650	39	64				
Over 650	3 mL per 50 kg					

5L PACK

Animals and Pest	State	Bodyweight (kg)	Dose (mL)	No. treated per 5 L pack	Critical comments	
<u>Beef cattle after weaning</u> Cattle Tick <i>(Rhipicephalus microplus)</i>	QLD, NSW, WA & NT only	101 – 150	9	555	<ol style="list-style-type: none"> 1. Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. 2. Maximum number of treatments per annual tick season: 3. 3. Apply in two bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator. 4. The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose. 5. There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves. 6. TICKPRO POUR-ON should not be applied to show cattle as a light scurfiness may be visible for some time after treatment. 7. TICKPRO POUR-ON is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of TICKPRO POUR-ON commences within 3 days of treatment. 8. Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas. 	Cattle application diagram as per reference product
		151 – 200	12	416		
		201 – 250	15	333		
		251 – 300	18	277		
		301 – 350	21	238		
		351 – 400	24	208		
		401 – 450	27	185		
		451 – 500	30	166		
		501 – 550	33	151		
		551 – 600	36	138		
		601 – 650	39	128		
Over 650	3 mL per 50 kg					

10L PACK

Animals and Pest	State	Bodyweight (kg)	Dose (mL)	No. treated per 25 L pack	Critical comments	
Beef cattle after weaning Cattle Tick (<i>Rhipicephalus microplus</i>)	QLD, NSW, WA & NT only	101 – 150	9	1111	<ol style="list-style-type: none"> 1. Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. 2. Maximum number of treatments per annual tick season: 3. 3. Apply in two bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator. 4. The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose. 5. There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves. 6. TICKPRO POUR-ON should not be applied to show cattle as a light scurfiness may be visible for some time after treatment. 7. TICKPRO POUR-ON is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of TICKPRO POUR-ON commences within 3 days of treatment. 8. Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas. 	Cattle application diagram as per reference product
		151 – 200	12	833		
		201 – 250	15	666		
		251 – 300	18	555		
		301 – 350	21	476		
		351 – 400	24	416		
		401 – 450	27	370		
		451 – 500	30	333		
		501 – 550	33	303		
		551 – 600	36	277		
		601 – 650	39	256		
Over 650	3 mL per 50 kg					

20L PACK

Animals and Pest	State	Bodyweight (kg)	Dose (mL)	No. treated per 20 L pack	Critical comments	
<p><u>Beef cattle after weaning</u> Cattle Tick (<i>Rhipicephalus microplus</i>)</p>	<p>QLD, NSW, WA & NT only</p>	101 – 150	9	2222	<p>1. Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. 2. Maximum number of treatments per annual tick season: 3. 3. Apply in two bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator. 4. The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose. 5. There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves. 6. TICKPRO POUR-ON should not be applied to show cattle as a light scurfiness may be visible for some time after treatment. 7. TICKPRO POUR-ON is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of TICKPRO POUR-ON commences within 3 days of treatment. 8. Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas.</p>	<p>Cattle application diagram as per reference product</p>
		151 – 200	12	1666		
		201 – 250	15	1333		
		251 – 300	18	1111		
		301 – 350	21	952		
		351 – 400	24	833		
		401 – 450	27	740		
		451 – 500	30	666		
		501 – 550	33	606		
		551 – 600	36	555		
		Over 650	3 mL per 50 kg			

25 L PACK

Animals and Pest	State	Bodyweight (kg)	Dose (mL)	No. treated per 25 L pack	Critical comments	
<p><u>Beef cattle after weaning</u> Cattle Tick (<i>Rhipicephalus microplus</i>)</p>	<p>QLD, NSW, WA & NT only</p>	101 – 150	9	2778	<ol style="list-style-type: none"> 1. Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. 2. Maximum number of treatments per annual tick season: 3. 3. Apply in two bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator. 4. The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose. 5. There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves. 6. TICKPRO POUR-ON should not be applied to show cattle as a light scurfiness may be visible for some time after treatment. 7. TICKPRO POUR-ON is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of TICKPRO POUR-ON commences within 3 days of treatment. 8. Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas. 	<p>Cattle application diagram as per reference product</p>
		151 – 200	12	2083		
		201 – 250	15	1667		
		251 – 300	18	1389		
		301 – 350	21	1190		
		351 – 400	24	1042		
		401 – 450	27	926		
		451 – 500	30	833		
		501 – 550	33	758		
		551 – 600	36	694		
		601 – 650	39	641		
Over 650		3 mL per 50 kg				