


**TEMPLATE FOR RELEVANT LABEL PARTICULARS (RLPs) (Veterinary Products)**

Select appropriate:

 New Product (include all applicable RLPs) OR

 Variation (highlight instructions that are being varied). Approval no. of label being varied: [APVMA No. 67327 /ATS No. 55810]

Signal heading:	<b>POISON</b> <b>KEEP OUT OF REACH OF CHILDREN</b> <b>READ SAFETY DIRECTIONS BEFORE OPENING OR USING</b> <b>FOR ANIMAL TREATMENT ONLY</b>	
Product name:	<b>Coopers® TRIFECTA® Triple Active Drench for Sheep and Cattle Mineralised</b>	
Active constituent/s:	2 g/L Abamectin, 80 g/L Levamisole hydrochloride and 45.3 g/L Oxfendazole 5 g/L Cobalt (as Cobalt EDTA) and 1 g/L Selenium (as Sodium Selenate)	
Statement of claims (short claim):	For the treatment and control of internal parasites of sheep including those resistant to Macrocytic Lactone (ML), Benzimidazole (BZ) or Levamisole (LEV) drench families. For the treatment and control of internal parasites of cattle including those resistant to the Macrocytic Lactone (ML) drench family. To supplement diets which may be deficient in selenium and cobalt.	
(long claim):	COOPERS TRIFECTA is a combination of three active ingredients including abamectin (a member of the Macrocytic Lactone (ML) drench family); oxfendazole (a member of the Benzimidazole (white) drench family) and levamisole hydrochloride (an Imidazothiazole and member of the clear drench family). It also contains two minerals including selenium (as sodium selenate) and cobalt (as cobalt EDTA). COOPERS TRIFECTA provides treatment and control of the following roundworms, sensitive to at least one of the three active ingredients in this product. <b>SHEEP</b> <b>Adult gastro-intestinal roundworms</b> <ul style="list-style-type: none"> <li>• Barber's Pole Worm (<i>Haemonchus contortus</i>)</li> <li>• Small Brown Stomach Worm (<i>Teladorsagia circumcincta</i>)</li> <li>• Stomach Hair Worm (<i>Trichostrongylus axei</i>)</li> <li>• Black Scour Worm (<i>Trichostrongylus colubriformis</i> †; <i>T. vitrinus</i>)            (†including <i>T. colubriformis</i> resistant to a combination of benzimidazoles and levamisole)</li> <li>• Small Intestinal Worm (<i>Cooperia curticei</i>; <i>C. oncophora</i>)</li> <li>• Thin Necked Intestinal Worm (<i>Nematodirus filicollis</i>; <i>N. spathiger</i>; <i>N. abnormalis</i>)</li> <li>• Hookworm (<i>Bunostomum trigonocephalum</i>)</li> <li>• Intestinal Threadworm (<i>Strongyloides papillosus</i>)</li> <li>• Large Bowel Worm (<i>Oesophagostomum venulosum</i>)</li> <li>• Nodule worm (<i>Oesophagostomum columbianum</i>)</li> <li>• Large Mouthed Bowel Worm (<i>Chabertia ovina</i>)</li> <li>• Whipworm (<i>Trichuris ovis</i>)</li> </ul>	

	<p><b>CATTLE</b></p> <p><b>Adult gastro-intestinal roundworms</b></p> <ul style="list-style-type: none"> <li>• Barber's Pole Worm (<i>Haemonchus placei</i>)</li> <li>• Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>)</li> <li>• Stomach Hair Worm (<i>Trichostrongylus axei</i>)</li> <li>• Black Scour Worm (<i>Trichostrongylus longispicularis</i>)</li> <li>• Small Intestinal Worm (<i>Cooperia oncophora</i>; <i>C. pectinata</i>; <i>C. punctata</i>)</li> <li>• Thin Necked Intestinal Worm (<i>Nematodirus helvetianus</i>)</li> <li>• Hookworm (<i>Bunostomum phlebotomum</i>)</li> <li>• Capillarids (<i>Capillaria bovis</i>)</li> <li>• Nodule Worm (<i>Oesophagostomum radiatum</i>)</li> </ul> <p>Resistance may develop to any drench. Ask your local veterinarian or animal health advisor for recommended parasite management practices for your area to reduce resistance development. It is advisable that a drench resistance test should be conducted before any drench is used.</p>
Net contents:	<b>1L, 2.5L, 5L, 6L, 10L, 15L, 20L</b>
Directions for Use Heading:	<p><b>DIRECTIONS FOR USE</b></p> <p><b>Before commencing drenching read the label carefully.</b></p>
Restrains:	<p><b>RESTRAINTS</b></p> <p><b>DO NOT USE</b> in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.</p> <p><b>DO NOT USE</b> in lambs weighing 15 kg or less.</p> <p><b>DO NOT USE</b> in calves weighing 100 kg or less.</p>
Contraindications:	<p>This product is contraindicated for use in lambs under six (6) weeks of age or weighing 15 kg or less.</p> <p>This product is contraindicated for use in calves less than 16 weeks of age or weighing 100 kg or less.</p> <p>This product is contraindicated for use when selenium intake from pasture is high or selenium is provided by other means (such as pasture top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.</p>
Precautions:	<p><b>Precautions</b></p> <p>Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching. Recommended dose should not be exceeded.</p> <p>Selenium can have toxic effects, particularly in cattle and sheep with liver damage including cattle and sheep that graze in areas where plants containing pyrrolizidine alkaloids are prevalent.</p> <p>Safety in breeding cattle and sheep of either sex and in pregnant or lactating cows or ewes has not been established.</p>
Side Effects:	

<p>Dosage &amp; Administration:</p>	<p><b>DOSAGE:</b></p> <p><b>SHAKE WELL BEFORE USE.</b></p> <p>COOPERS TRIFECTA should be administered orally using accurate drenching equipment. Clean equipment thoroughly before and after use.</p> <p>Always dose on a liveweight basis, do not underdose. The heaviest animal in a group should be weighed. Dose the group to the heaviest animal by liveweight (as ewes, wethers, rams, lambs, cows, bulls, calves), do not underdose. For cattle, where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Check the accuracy of drenching equipment before and during drenching procedures.</p> <p><b>Re-Treatment Interval (CATTLE/SHEEP): DO NOT retreat animals for 28 days after last treatment.</b></p> <p><b>Dose Rates:</b> 1 mL per 10 kg bodyweight by a single oral administration.</p> <table border="1" data-bbox="443 725 1453 1055"> <thead> <tr> <th colspan="2">SHEEP:</th> <th colspan="7">PACK TREATS</th> </tr> <tr> <th>LIVEWEIGHT</th> <th>DOSE</th> <th>1L</th> <th>2.5L</th> <th>5L</th> <th>6L</th> <th>10L</th> <th>15L</th> <th>20L</th> </tr> </thead> <tbody> <tr> <td>15 kg or less</td> <td>Do not treat</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>16-20 kg</td> <td>2 mL</td> <td>500</td> <td>1250</td> <td>2500</td> <td>3000</td> <td>5000</td> <td>7500</td> <td>10000</td> </tr> <tr> <td>21-30 kg</td> <td>3 mL</td> <td>333</td> <td>833</td> <td>1666</td> <td>2000</td> <td>3333</td> <td>5000</td> <td>6666</td> </tr> <tr> <td>31-40 kg</td> <td>4 mL</td> <td>250</td> <td>625</td> <td>1250</td> <td>1500</td> <td>2500</td> <td>3750</td> <td>5000</td> </tr> <tr> <td>41-50 kg</td> <td>5 mL</td> <td>200</td> <td>500</td> <td>1000</td> <td>1200</td> <td>2000</td> <td>3000</td> <td>4000</td> </tr> <tr> <td>51-60 kg</td> <td>6 mL</td> <td>166</td> <td>416</td> <td>833</td> <td>1000</td> <td>1666</td> <td>2500</td> <td>3333</td> </tr> <tr> <td>61-70 kg</td> <td>7 mL</td> <td>142</td> <td>357</td> <td>714</td> <td>857</td> <td>1428</td> <td>2142</td> <td>2857</td> </tr> <tr> <td>71-80 kg</td> <td>8 mL</td> <td>125</td> <td>312</td> <td>625</td> <td>750</td> <td>1250</td> <td>1875</td> <td>2500</td> </tr> <tr> <td>Over 80 kg</td> <td>1 mL/10 kg</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table> <table border="1" data-bbox="443 1095 1453 1525"> <thead> <tr> <th colspan="2">CATTLE:</th> <th colspan="7">PACK TREATS</th> </tr> <tr> <th>LIVEWEIGHT</th> <th>DOSE</th> <th>1L</th> <th>2.5L</th> <th>5L</th> <th>6L</th> <th>10L</th> <th>15L</th> <th>20L</th> </tr> </thead> <tbody> <tr> <td>100 kg or less</td> <td>Do not treat</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> <tr> <td>101-150 kg</td> <td>15 mL</td> <td>66</td> <td>166</td> <td>333</td> <td>400</td> <td>666</td> <td>1000</td> <td>1333</td> </tr> <tr> <td>151-200 kg</td> <td>20 mL</td> <td>50</td> <td>125</td> <td>250</td> <td>300</td> <td>500</td> <td>7500</td> <td>1000</td> </tr> <tr> <td>201-250 kg</td> <td>25 mL</td> <td>40</td> <td>100</td> <td>200</td> <td>240</td> <td>400</td> <td>600</td> <td>800</td> </tr> <tr> <td>251-300 kg</td> <td>30 mL</td> <td>33</td> <td>83</td> <td>166</td> <td>200</td> <td>333</td> <td>500</td> <td>666</td> </tr> <tr> <td>301-350 kg</td> <td>35 mL</td> <td>28</td> <td>71</td> <td>142</td> <td>171</td> <td>285</td> <td>428</td> <td>571</td> </tr> <tr> <td>351-400 kg</td> <td>40 mL</td> <td>25</td> <td>62</td> <td>125</td> <td>150</td> <td>250</td> <td>375</td> <td>500</td> </tr> <tr> <td>401-450 kg</td> <td>45 mL</td> <td>22</td> <td>55</td> <td>111</td> <td>133</td> <td>222</td> <td>333</td> <td>444</td> </tr> <tr> <td>451-500 kg</td> <td>50 mL</td> <td>20</td> <td>50</td> <td>100</td> <td>120</td> <td>200</td> <td>300</td> <td>400</td> </tr> <tr> <td>501-550 kg</td> <td>55 mL</td> <td>18</td> <td>45</td> <td>90</td> <td>109</td> <td>181</td> <td>272</td> <td>363</td> </tr> <tr> <td>551-600 kg</td> <td>60 mL</td> <td>16</td> <td>41</td> <td>83</td> <td>100</td> <td>166</td> <td>250</td> <td>333</td> </tr> <tr> <td>601-650 kg</td> <td>65 mL</td> <td>15</td> <td>38</td> <td>76</td> <td>92</td> <td>153</td> <td>230</td> <td>307</td> </tr> <tr> <td>Over 650 kg</td> <td>1 mL/10 kg</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> </tr> </tbody> </table>	SHEEP:		PACK TREATS							LIVEWEIGHT	DOSE	1L	2.5L	5L	6L	10L	15L	20L	15 kg or less	Do not treat								16-20 kg	2 mL	500	1250	2500	3000	5000	7500	10000	21-30 kg	3 mL	333	833	1666	2000	3333	5000	6666	31-40 kg	4 mL	250	625	1250	1500	2500	3750	5000	41-50 kg	5 mL	200	500	1000	1200	2000	3000	4000	51-60 kg	6 mL	166	416	833	1000	1666	2500	3333	61-70 kg	7 mL	142	357	714	857	1428	2142	2857	71-80 kg	8 mL	125	312	625	750	1250	1875	2500	Over 80 kg	1 mL/10 kg								CATTLE:		PACK TREATS							LIVEWEIGHT	DOSE	1L	2.5L	5L	6L	10L	15L	20L	100 kg or less	Do not treat								101-150 kg	15 mL	66	166	333	400	666	1000	1333	151-200 kg	20 mL	50	125	250	300	500	7500	1000	201-250 kg	25 mL	40	100	200	240	400	600	800	251-300 kg	30 mL	33	83	166	200	333	500	666	301-350 kg	35 mL	28	71	142	171	285	428	571	351-400 kg	40 mL	25	62	125	150	250	375	500	401-450 kg	45 mL	22	55	111	133	222	333	444	451-500 kg	50 mL	20	50	100	120	200	300	400	501-550 kg	55 mL	18	45	90	109	181	272	363	551-600 kg	60 mL	16	41	83	100	166	250	333	601-650 kg	65 mL	15	38	76	92	153	230	307	Over 650 kg	1 mL/10 kg							
SHEEP:		PACK TREATS																																																																																																																																																																																																																																									
LIVEWEIGHT	DOSE	1L	2.5L	5L	6L	10L	15L	20L																																																																																																																																																																																																																																			
15 kg or less	Do not treat																																																																																																																																																																																																																																										
16-20 kg	2 mL	500	1250	2500	3000	5000	7500	10000																																																																																																																																																																																																																																			
21-30 kg	3 mL	333	833	1666	2000	3333	5000	6666																																																																																																																																																																																																																																			
31-40 kg	4 mL	250	625	1250	1500	2500	3750	5000																																																																																																																																																																																																																																			
41-50 kg	5 mL	200	500	1000	1200	2000	3000	4000																																																																																																																																																																																																																																			
51-60 kg	6 mL	166	416	833	1000	1666	2500	3333																																																																																																																																																																																																																																			
61-70 kg	7 mL	142	357	714	857	1428	2142	2857																																																																																																																																																																																																																																			
71-80 kg	8 mL	125	312	625	750	1250	1875	2500																																																																																																																																																																																																																																			
Over 80 kg	1 mL/10 kg																																																																																																																																																																																																																																										
CATTLE:		PACK TREATS																																																																																																																																																																																																																																									
LIVEWEIGHT	DOSE	1L	2.5L	5L	6L	10L	15L	20L																																																																																																																																																																																																																																			
100 kg or less	Do not treat																																																																																																																																																																																																																																										
101-150 kg	15 mL	66	166	333	400	666	1000	1333																																																																																																																																																																																																																																			
151-200 kg	20 mL	50	125	250	300	500	7500	1000																																																																																																																																																																																																																																			
201-250 kg	25 mL	40	100	200	240	400	600	800																																																																																																																																																																																																																																			
251-300 kg	30 mL	33	83	166	200	333	500	666																																																																																																																																																																																																																																			
301-350 kg	35 mL	28	71	142	171	285	428	571																																																																																																																																																																																																																																			
351-400 kg	40 mL	25	62	125	150	250	375	500																																																																																																																																																																																																																																			
401-450 kg	45 mL	22	55	111	133	222	333	444																																																																																																																																																																																																																																			
451-500 kg	50 mL	20	50	100	120	200	300	400																																																																																																																																																																																																																																			
501-550 kg	55 mL	18	45	90	109	181	272	363																																																																																																																																																																																																																																			
551-600 kg	60 mL	16	41	83	100	166	250	333																																																																																																																																																																																																																																			
601-650 kg	65 mL	15	38	76	92	153	230	307																																																																																																																																																																																																																																			
Over 650 kg	1 mL/10 kg																																																																																																																																																																																																																																										
<p>General Directions:</p>	<p><b>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</b></p>																																																																																																																																																																																																																																										
<p>Withholding Period/s:</p>	<p><b>MEAT (CATTLE, SHEEP): DO NOT USE less than 21 days before slaughter for human consumption.</b></p> <p><b>MILK: CATTLE: DO NOT USE in lactating cows or within 28 days of calving where milk or milk products will be used for human consumption. If cows calve earlier than 28 days after treatment, milk may contain residues. This milk MUST NOT be used for human consumption, or supplied for processing for at least 28 days following treatment. Calves fed this milk should not be slaughtered for human consumption within 21 days.</b></p> <p><b>SHEEP: DO NOT USE in ewes which are producing or may in the future</b></p>																																																																																																																																																																																																																																										

	<b>produce milk that may be used or processed for human consumption.</b>
Trade Advice:	<b>TRADE ADVICE</b> <b>EXPORT SLAUGHTER INTERVAL (ESI):</b> CATTLE: DO NOT USE less than 21 days before slaughter for export. SHEEP: DO NOT USE less than 28 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the manufacturer on 1800 226 511 or the APVMA website <a href="http://www.apvma.gov.au/residues/">www.apvma.gov.au/residues/</a>
Safety Directions:	<b>Harmful if inhaled. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing.</b>
First Aid:	<b>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126, New Zealand 0800 764 766. If skin contact occurs, remove contaminated clothing and wash skin thoroughly.</b>
Additional User Safety:	Additional information is in the Material Safety Data Sheet. <b>For further information contact COOPERS Customer Service on 1 800 226 511.</b>
Environmental Statements:	<b>PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT</b> Very toxic to aquatic life. DO NOT contaminate dams, rivers, streams or other waterways with the chemical or used containers.
Disposal: (1L) Disposal: (2L and above)	<b>Dispose</b> of empty container by wrapping with paper and putting in garbage. This container can be recycled if it is clean, dry, free of visible residues and has the <b>drumMUSTER</b> logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any <b>drumMUSTER</b> collection or similar container management program site. The cap should not be replaced but may be taken separately.
Storage:	<b>Store</b> below 30°C (Room Temperature) in the closed original container in a well ventilated area. Protect from light.
Batch and Expiry	<b>SEE BELOW FOR BATCH AND EXPIRY</b> <b>(B) EXPIRY</b>
Name & address:	<b>COOPERS ANIMAL HEALTH</b> Intervet Australia Pty Limited (trading as MSD Animal Health) 91-105 Harpin Street, Bendigo East VIC 3550

The following is for APVMA use only:

APVMA approval no.	67327/61112
--------------------	-------------