



Company Name: SPECIALIST SALES PTY LTD
Product Name: CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE
APVMA Approval No: 80714/101481

Label Name:	CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE
-------------	---

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	5 g/L IVERMECTIN
-------------------------	------------------

Claims:	<p>For the treatment and control of Ivermectin-sensitive internal and external parasites of beef and dairy cattle.</p> <p>For the treatment and control of Ivermectin sensitive:</p> <p>Gastrointestinal nematodes:</p> <ul style="list-style-type: none">• Adult and immature (L4): Small Brown Stomach Worm (including inhibited stages) (<i>Ostertagia ostertagi</i>), Barber's Pole Worm (<i>Haemonchus placei</i>), Stomach Hair Worm (<i>Trichostrongylus axei</i>), Intestinal Hair Worm (<i>T. colubriformis</i>), Small Intestinal Worm (<i>Cooperia</i> spp.), Nodule Worm (<i>Oesophagostomum radiatum</i>); and• Adult stages of <i>Cooperia oncophora</i>, <i>C.punctata</i>, <i>Oesophagostomum venulosum</i>, Whip Worm (<i>Trichuris</i> spp.), Intestinal Threadworm (<i>Strongyloides papillosus</i>); and• Immature stages (L4) of Thin Necked Intestinal Worm (<i>Nematodirus</i> spp.). <p>Lungworms (adult and immature) <i>Dictyocaulus viviparus</i>. Eyeworm <i>Thelazia</i> spp. (adult stages). Sucking and Biting Lice: <i>Linognathus vituli</i>, <i>Haematopinus eurysternus</i>, <i>Bovicola bovis</i>, <i>Solenopotes capillatus</i>. Mites: <i>Chorioptes bovis</i>, <i>Sarcoptes scabiei</i> var <i>bovis</i>. Buffalo Flies <i>Haematobia irritans exigua</i> Cattle Ticks: (<i>Rhipicephalus</i> [<i>Boophilus</i>] <i>microplus</i>) including organophosphate, synthetic pyrethroid- and amidine-resistant strains. Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control programme. CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE should be alternated with conventional dips or pour-ons when the programme requires repeated treatments.</p> <p>PERSISTENT ACTIVITY: CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE applied at the recommended dose of 1mL per 10kg of liveweight controls infections of Stomach Hair Worm (<i>Trichostrongylus axei</i>) and Small Intestinal Worm (<i>Cooperia</i> spp.) for up to 14 days after treatment; Small Brown Stomach Worm (<i>Ostertagia</i> spp.) and Nodule Worm</p>
---------	---

(Oesophagostomum radiatum) for up to 21 days after treatment; Lungworm (Dictyocaulus viviparus) for up to 28 days after treatment; Buffalo Fly (Haematobia irritans exigua) for up to 21 days after treatment.

Net Contents: 1L [5L, 10L, 20L, 25L]

Directions for Use: DIRECTIONS FOR USE:

Restrains: Restraints
DO NOT USE in pregnant cows less than 7 days before calving.

Contraindications:

Precautions:

Side Effects:

Dosage and Administration:

Dosage and Administration:

This product is formulated only for external application to cattle; do not use in other species. The formulation should be applied along the topline in a narrow strip extending from the withers to the tailhead. The dose is 1mL for each 10kg of liveweight.

Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers etc.). When there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid overdosing. Do not underdose. Cattle should be weighed with either scales or with a weighband.

[Refer to the attached DOSAGE TABLES]

Heavier animals (for example, mature bulls) should receive an additional 5mL for each 50 kg over 650 kg.

Treatment should not be repeated in less than 42 days.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

This section contains file attachment.

File Name: eLabel_CattlePro PO Dosage Tables_20 Jan 15.pdf

File Size: 38972 bytes

General Directions:

General Directions:

For best results CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE should be part of a total parasite control programme including internal and external parasites. Consult your veterinarian for a complete parasite control programme.

Administration

Efficacy of CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE is not adversely affected when the hide is wet or if rainfall occurs shortly after treatment. However, it is not considered good agricultural practice to treat animals if they are wet or rain is likely.

[5 Litre Collapsible Pack:]

Use dosing equipment compatible with CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE. Follow manufacturer's directions for use and care of equipment. Other dosing equipment may be incompatible, resulting in locking, incorrect dosing and leakage. Connect the pour-on applicator to the pack as follows:

- Attach the open end of the draw-off tubing to the dosing gun and attach draw-off tubing to cap with stem.
- Replace the shipping cap with the supplied cap that has the stem for attaching the draw-off tubing and tighten this cap.
- Gently prime the pour-on applicator, checking for leaks.

Use only in well-ventilated areas or outdoors.

Close container when not in use.

Cloudiness may result when CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE is stored at temperatures below 0°C. Allowing to warm at room temperature will restore the normal appearance without affecting efficacy.

The antiparasitic activity of ivermectin will be impaired if the formulation is applied to areas of skin with mange scabs or lesions, or with dermatoses or adherent materials, e.g. caked mud or manure.

This product is for application to the skin surface only, do not give orally or parenterally. CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE is not recommended for use in species other than cattle.

NOTE TO USER:

The colour of CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE may fade when exposed to light. This loss of colour does not reflect loss of potency of ivermectin. However, prolonged exposure (i.e. weeks) to light can result in a gradual decline of ivermectin potency in the formulation.

SAFETY TO CATTLE:

Field studies have demonstrated a wide safety margin. The topically applied formulation is expected to be well tolerated by breeding animals.

Withholding Periods:	WITHHOLDING PERIOD: MEAT: DO NOT USE less than 28 days before slaughter for human consumption. MILK: Zero (0) days Bobby calves: DO NOT SLAUGHTER for 4 days bobby calves born to cows treated during pregnancy.
-----------------------------	--

Trade Advice:	TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): DO NOT SLAUGHTER for export for 28 days after treatment. Before using this product, confirm the current ESI from Specialist Sales on 1800 780 317 or the APVMA website (www.apvma.gov.au/residues).
----------------------	---

Safety Directions:	SAFETY DIRECTIONS: Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. If product gets on skin, immediately wash the area with soap and water. If product gets in eyes, wash out immediately with water. Wash hands after use.
---------------------------	--

First Aid Instructions:	FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126.
--------------------------------	--

First Aid Warnings:	
----------------------------	--

Additional User Safety:	ADDITIONAL USER SAFETY INFORMATION: For further information, refer to the Safety Data Sheet.
--------------------------------	--

Environmental Statements:	ENVIRONMENTAL PROTECTION Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container. When used according to label directions CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE is not likely to have an unintended effect that is harmful to the environment or have an overall adverse effect on dung beetle or earthworm populations. Increased mortality and impaired development of dung beetle larvae may be expected, but for a limited period after treatment. There may also be short-term indirect effect on the rate of earthworm colonisation of dung pats.
----------------------------------	--

Disposal:	DISPOSAL: [1L] Dispose of container by wrapping with paper and putting in garbage. [Other pack sizes] Triple-rinse container into the medicated pour-on. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
------------------	---

Storage:	STORAGE: Store below 30°C (room temperature). Protect from light.
-----------------	---

CATTLEPRO POUR-ON FOR BEEF AND DAIRY CATTLE – DOSAGE TABLES:

(1L pack)

Liveweight (kg)	Dose (mL)	This pack treats (head)	Liveweight (kg)	Dose (mL)	This pack treats (head)
< 100	1mL/10kg	≥100	351 – 400	40	25
101 – 150	15	66	401 – 450	45	22
151 – 200	20	50	451 – 500	50	20
201 – 250	25	40	501 – 550	55	18
251 – 300	30	33	551 – 600	60	16
301 – 350	35	28	601 – 650	65	15

(5L pack)

Liveweight (kg)	Dose (mL)	This pack treats (head)	Liveweight (kg)	Dose (mL)	This pack treats (head)
< 100	1mL/10kg	≥500	351 – 400	40	125
101 – 150	15	333	401 – 450	45	111
151 – 200	20	250	451 – 500	50	100
201 – 250	25	200	501 – 550	55	90
251 – 300	30	166	551 – 600	60	83
301 – 350	35	142	601 – 650	65	76

(10 L pack)

Liveweight (kg)	Dose (mL)	This pack treats (head)	Liveweight (kg)	Dose (mL)	This pack treats (head)
< 100	1mL/10kg	≥1000	351 – 400	40	250
101 – 150	15	666	401 – 450	45	222
151 – 200	20	500	451 – 500	50	200
201 – 250	25	400	501 – 550	55	181
251 – 300	30	333	551 – 600	60	166
301 – 350	35	285	601 – 650	65	153

(20L pack)

Liveweight (kg)	Dose (mL)	This pack treats (head)	Liveweight (kg)	Dose (mL)	This pack treats (head)
< 100	1mL/10kg	≥2000	351 – 400	40	500
101 – 150	15	1333	401 – 450	45	444
151 – 200	20	1000	451 – 500	50	400
201 – 250	25	800	501 – 550	55	363
251 – 300	30	666	551 – 600	60	333
301 – 350	35	571	601 – 650	65	307

(25L pack)

Liveweight (kg)	Dose (mL)	This pack treats (head)	Liveweight (kg)	Dose (mL)	This pack treats (head)
< 100	1mL/10kg	≥2500	351 – 400	40	625
101 – 150	15	1666	401 – 450	45	555
151 – 200	20	1250	451 – 500	50	500
201 – 250	25	1000	501 – 550	55	454
251 – 300	30	833	551 – 600	60	416
301 – 350	35	714	601 – 650	65	385