

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

# CYDECTIN® POUR-ON

FOR CATTLE & RED DEER

ACTIVE CONSTITUENT: 5 g/L MOXIDECTIN  
SOLVENT: 150 g/L HYDROCARBON LIQUID

For the treatment and control of moxidectin sensitive internal and external parasites of cattle and for the treatment and control of lungworm and gastrointestinal roundworms of red deer.

CONTENTS

## 15 Litres

CYPO15WLF10 1P0964-1



**Virbac**  
ANIMAL HEALTH



**CYDECTIN® POUR-ON FOR CATTLE AND RED DEER** comes ready to use as a clear solution containing 5 g/L moxidectin. Moxidectin is a second generation macrocyclic lactone endectocide. It is effective against parasites sensitive to this family. Resistance can occur to any anthelmintic. **(See product leaflet for additional information)**

### DIRECTIONS FOR USE:

#### Dosage and Administration

##### Cattle and Red Deer

CYDECTIN® POUR-ON FOR CATTLE AND RED DEER should be applied along the top of the back from the withers to the base of the tail. Do not retreat deer within 60 days. Volume to use: 1 mL per 10 kg liveweight. This will administer the correct dose of 0.5 mg/kg liveweight.

CYDECTIN® POUR-ON can be used in lactating dairy cattle, but **DO NOT TREAT LACTATING CATTLE AT LESS THAN 21 DAY INTERVALS.**

The following table may be used as a guide for the dose volume to use.

Liveweight (kg)	Dose (mL)	Doses Per Pack (15L)	Liveweight (kg)	Dose (mL)	Doses Per Pack (15L)
31 - 40	4	3750	176 - 200	20	750
41 - 50	5	3000	201 - 250	25	600
51 - 60	6	2500	251 - 300	30	500
61 - 70	7	2142	301 - 350	35	428
71 - 80	8	1875	351 - 400	40	375
81 - 90	9	1666	401 - 450	45	333
91 - 100	10	1500	451 - 500	50	300
101 - 125	12.5	1200	501 - 550	55	272
126 - 150	15	1000	551 - 600	60	250
151 - 175	17.5	857	601 - 650	65	230

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc.). Cattle in excess of 650 kg should be dosed at 1 mL per 10 kg liveweight.

Do not underdose. Check accuracy of dose delivery of applicator regularly.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

#### Broad Spectrum

CYDECTIN® POUR-ON FOR CATTLE AND RED DEER is efficacious against roundworms, lungworm, cattle tick, lice and mites which infect cattle, and lungworm and roundworms of red deer.

#### Persistent Protection Period

CYDECTIN® POUR-ON FOR CATTLE AND RED DEER prevents re-infection with the following parasite species:

Parasite species	Persistent protection period
<i>Ostertagia ostertagi</i> ( <i>O. lyrata</i> )	42 days
<i>Haemonchus placei</i>	28 days
<i>Trichostrongylus axei</i>	28 days
<i>Dictyocaulus viviparus</i>	42 days
<i>Oesophagostomum radiatum</i>	42 days
<i>Bunostomum phlebotomum</i>	42 days
<i>Nematodirus helvetianus</i>	21 days

The persistent activity of CYDECTIN® POUR-ON FOR CATTLE AND RED DEER prevents the development of viable cattle tick (*Boophilus microplus*) for 21 days after treatment.

The persistent activity of CYDECTIN® POUR-ON FOR CATTLE AND RED DEER

can be used as an aid in the control of buffalo fly (*Haematobia irritans exigua*) for up to 14 days after treatment.

#### Rainfast

The efficacy of CYDECTIN® POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However, the efficacy against tick and buffalo fly may be reduced if very heavy rain occurs following treatment. Treatment of animals under these circumstances is not good agricultural practice.

#### Weatherproof

The efficacy of CYDECTIN® POUR-ON is not adversely affected if applied to cattle under extreme seasonal conditions such as very high UV-light and temperatures in Summer and very low temperatures in Winter.

#### Safety to Target animals

CYDECTIN® POUR-ON FOR CATTLE AND RED DEER has a wide margin of safety when used at the recommended dose rate in all ages and breeds of cattle including pregnant cows and breeding bulls.

#### Concurrent Use

CYDECTIN® POUR-ON FOR CATTLE AND RED DEER is safe to use concurrently with other treatments in cattle including mineral supplements, fluke treatments and vaccines.

**Do not apply at the same time as any other pour-on product.**

#### Effect on Dung Beetles

When applied as directed, the levels of CYDECTIN® POUR-ON in the faeces of treated cattle are not likely to have any significant adverse effect on the following dung beetles: *Onthophagus gazella*, *O. taurus*, *Euoniticellus intermedius* and *E. fulvus*. Effects on other dung beetle species have not been fully evaluated.

#### WITHHOLDING PERIODS:

MEAT: CATTLE - NIL

DEER - **DO NOT USE less than 7 days before slaughter for human consumption**

MILK: NIL

The withholding period stated on this label applies only to meat destined for the Australian market or for export to USA. Some export markets apply different standards that may require an export slaughter interval that is longer than the withholding period. If necessary, details of overseas standards should be obtained prior to use of this product.

#### SAFETY DIRECTIONS:

Poisonous if swallowed. May irritate the nose and throat and will irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale. Wash hands after use. Additional information is listed in the Material Safety Data Sheet.

#### FIRST AID:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

#### Parasite Control Programs

Effective parasite control for maximum productivity is an essential part of farm management. Follow recommended parasite management practices to reduce the potential for resistance to develop.

#### DISPOSAL:

This container can be recycled if it is clean, dry, free of visible residues and has the drumMUSTER logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any drumMUSTER collection or similar container management program site. The cap should not be replaced but may be taken separately.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Moxidectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

#### STORAGE:

Store below 30°C (Room temperature) and protect from light.



**Virbac (AUSTRALIA) PTY. LIMITED**  
361 Horsley Road, Milperra NSW 2214

Freecall Customer Service Line:  
1800 009 847

[www.virbac.com.au](http://www.virbac.com.au)

[B]:

EXP:

APVMA Approval No. 45970/15L/1103

CYPO15WLB10 1P0965-1

© Registered Trademark of

Virbac (Australia) Pty. Limited



9 315216 501977