

TICKPRO™

POUR-ON TICK DEVELOPMENT INHIBITOR



DIRECTIONS FOR USE:

RESTRAINT:

DO NOT USE on cattle which are producing or may in the future produce milk for human consumption.

CONTRAINDICATIONS:

This product is contraindicated for use on pregnant cows within 6 weeks of calving to avoid handling stress.

DOSAGE AND ADMINISTRATION:

Animals and Pest	State	Body-weight (kg)	Dose (mL)	No. treated per 5L pack	Critical comments
Beef cattle after weaning. Cattle Tick (<i>Rhipicephalus microplus</i>)	QLD, NSW, WA & NT only	101 – 150	9	555	<ol style="list-style-type: none"> Retreat when you first see ticks but not before at least 6 weeks after the previous treatment. Maximum number of treatments per annual tick season: 3. Apply in two bands, about 7cm wide, one on each side of the spine between the rump and the shoulders with an approved applicator. DO NOT use any other applicator. The weight of cattle should be determined by either scales or a weigh-band. Dose individuals according to the heaviest animal in the mob. DO NOT under dose. There is no need to treat calves as long as they suckle treated cows. Calves will receive adequate protection from their mother. A reduced length of protection may occur in cows which are suckling calves. TICKPRO POUR-ON should not be applied to show cattle as a light scurfiness may be visible for some time after treatment. TICKPRO POUR-ON is a tick development inhibitor and kills ticks progressively. It can take 2 to 3 weeks for cattle to be visibly free of ticks. No additional treatment is necessary during that time, even on heavily infested animals. The development inhibiting effect of TICKPRO POUR-ON commences within 3 days of treatment. Not suitable for cleaning cattle moving into 'gazetted' cattle tick areas.
		151 – 200	12	416	
		201 – 250	15	333	
		251 – 300	18	277	
		301 – 350	21	238	
		351 – 400	24	208	
		401 – 450	27	185	
		451 – 500	30	166	
		501 – 550	33	151	
		551 – 600	36	138	
601 – 650	39	128			
Over 650		3 mL per 50 kg			



NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

TICKPRO™

POUR-ON TICK DEVELOPMENT INHIBITOR

ACTIVE CONSTITUENT: 25g/L FLUAZURON
Also Contains: (Solvents) 100 g/L 1-dodecyl-2-pyrrolidone
100 g/L 1-octyl-2-pyrrolidone
415 g/L 1-methyl-2-pyrrolidone



For strategic control of the Cattle Tick, *Rhipicephalus (Boophilus) microplus* (including all known acaricide resistant strains), on Beef Cattle.

TICKPRO™

POUR-ON TICK DEVELOPMENT INHIBITOR



GENERAL DIRECTIONS:

TICKPRO POUR-ON Treated Areas – Introduction of New Cattle

Tick infested cattle introduced onto properties where a TICKPRO POUR-ON treatment regime is in place should be treated with TICKPRO POUR-ON prior to release.

Handling of Treated Animals

Wear protective clothing recommended under Safety Directions, if handling treated cattle within 24 hours of application.

For local tick control recommendations contact your local department of agriculture.

Cleaning Instructions

After use wash TICKPRO POUR-ON applicator with detergent and hot water. Dispose of the rinsates in a disposal pit away from desirable plants and their roots and watercourses.

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 42 days before slaughter for human consumption.

Calves which have suckled on treated cows must not be slaughtered less than 4 months after the last treatment of these cows.

MILK: DO NOT USE on cattle which are producing or may in the future produce milk that may be used or processed for human consumption.

TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from Specialist Sales on 1800 780 317 or the APVMA website (www.apvma.gov.au/residues).

SAFETY DIRECTIONS

Harmful if swallowed. Will damage eyes and skin. Avoid contact with eyes and skin. Repeated exposure may cause allergic disorders. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow length PVC gloves and a face shield. If clothing becomes contaminated with product remove clothing immediately. If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking and smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID: If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126. If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL USER SAFETY INFORMATION: For further information, refer to the Safety Data Sheet.

ENVIRONMENTAL PROTECTION

DO NOT contaminate fish ponds, dams or rivers with the product or used containers. This product is very toxic to aquatic invertebrates.

DISPOSAL:

Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury empty containers in a local authority landfill. If no landfill is available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE:

Store below 30°C (Room Temperature). Store in original container tightly closed in a dry, cool place. Store out of direct sunlight.

APVMA Approval No.: 81640/103523

Exp:

B:

Packed for:
Specialist Sales Pty Ltd
ACN 103 415 603
19 Clifford St, Toowoomba QLD 4350
Ph: 1800 780 317



AHD/TPSL1

File Name	TICKPro Pour-on_5L.pdf
Date	4.12.2015
Full-Size Printed Label Dimensions	470 x 230 mm
Approximate Scale (%) at which the pdf will print on A4 (or A3)	A4 : 51% A3 : 76%