

**POISON**

**KEEP OUT OF REACH OF CHILDREN**  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING



# Conquest Maca 600 Herbicide

**ACTIVE CONSTITUENT:** 600 g/L TRICLOPYR present as the butoxyethyl ester

**GROUP I HERBICIDE**

**For the control of a range of woody weeds and melons as specified in the  
Directions for Use table.**

**IMPORTANT: READ THE ATTACHED LEAFLET  
BEFORE USING THIS PRODUCT.**

**Contents:** 1 L, 5 L, 10L, 20 L, 200 L

**Conquest Agrochemicals Pty Ltd.**

ABN 84 098 814 932  
Suite 10, 41 Walters Drive  
Herdsman Business Park  
Osborne Park WA 6017

Telephone (08) 9244 1268  
Facsimile: (08) 9446 4696

## **STORAGE AND DISPOSAL**

Store in closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. **DO NOT** store near food, feedstuffs, fertilisers or seed. **DO NOT** re-use containers.

Triple rinse or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## **SMALL SPILL MANAGEMENT**

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb the wash liquid as above for disposal.

## **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. **DO NOT** inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

## **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (Ph: 13 1126). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## **MATERIAL SAFETY DATA SHEET**

Additional information is listed on the Material Safety Data Sheet which is available from the supplier.

## **CONDITIONS OF SALE**

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

<p>IN EMERGENCY DIAL</p> <p><b>000,</b></p> <p><b>POLICE OR FIRE BRIGADE</b></p>	<p>FOR SPECIALIST ADVICE IN AN EMERGENCY CALL</p> <p><b>1800 033 111</b></p> <p><b>(24 HOURS)</b></p>
----------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------

## **Conquest Agrochemicals Pty Ltd.**

ABN 84 098 814 932  
Suite 10, 41 Walters Drive  
Herdsman Business Park  
Osborne Park WA 6017  
Telephone (08) 9244 1268  
Facsimile: (08) 9446 4696

Batch No:

DOM:

APVMA approval No: 58112/1/0104

**POISON**

**KEEP OUT OF REACH OF CHILDREN**  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

# Conquest Maca 600 Herbicide

**ACTIVE CONSTITUENT:** 600 g/L TRICLOPYR present as the butoxyethyl ester

**GROUP I HERBICIDE**

**For the control of a range of woody weeds and melons as specified in the  
Directions for Use table.**

**IMPORTANT: READ THIS LEAFLET  
BEFORE USING THIS PRODUCT.**

**Conquest Agrochemicals Pty Ltd.**

ABN 84 098 814 932  
Suite 10, 41 Walters Drive  
Herdsman Business Park  
Osborne Park WA 6017

Telephone (08) 9244 1268  
Facsimile: (08) 9446 4696

**DIRECTIONS FOR USE****RESTRAINTS:**

- **DO NOT** apply to weeds which may be stressed (not actively growing) due to prolonged periods of extreme cold, moisture stress (water-logged or drought affected), poor nutrition, presence of disease, or previous herbicide treatment as reduced levels of control may result.
- **DO NOT** spray if rain is likely within one hour or if foliage is wet from rain and dew. However, when tankmixed with Conquest Glyphosate 450 Herbicide, this time extends to 6 hours.
- **DO NOT** burn off, cut or clear blackberry or other woody weeds for six months after spraying.

**1. WOODY WEED SITUATIONS****Table A: High Volume Spraying**See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /100 L water</b>	<b>CRITICAL COMMENTS</b>
African boxthorn	Less than 2m tall	Tas only	170 mL	
Angophora and Banksia regrowth	1 to 2 m tall	All States	400 or 560 mL	Use the higher rate on larger regrowth. Ensure the weed has dense foliage.
Blackberry	Active growth during late spring to early autumn		170 mL	Where herbicides other than Group I herbicides have been used, allow two seasons regrowth to occur before spraying with Conquest Maca 600. Any subsequent regrowth and seedlings should be sprayed after hardening off.
In association with: St John's wort	During flowering (Nov-Jan)	NSW, Vic and Tas only		Apply as a thorough foliage spray.
Brigalow	1 to 2 m tall	NSW and Qld only		Use at least 1000 L of water /ha.
Brooms: English	Spring to mid-summer prior to pod formation	All States		
Camphor laurel	Seedlings up to 3 m tall			
Capeweed	Rosette	Tas only	80 mL	
Common prickly pear	Active growth	All States	3 L	
English ivy	Active growth during late spring to late summer	Vic only	1 L + 1 L glyphosate (360g/L)	<b>DO NOT</b> treat ivy growing up trees or on other plants as death of the host may result. This mixture is not selective to grasses.
Eucalyptus	Seedlings and regrowth from small lignotubers, 1 to 2 m tall	Qld, SA, WA, & NT only	400 mL	Add Decol® S35B Surfactant at 250 mL to 500 mL/100 L of water for best results.
		NSW, Tas and Vic only	560 mL	
Fennel	1 to 2 m tall	Tas only	170 mL	Some regrowth may be expected the following season, which can be sprayed after hardening off.
Green cestrum		NSW, Qld and Vic only		
Groundsel bush ( <i>Baccharis halimifolia</i> )	Seedlings, 1 to 2 m tall	All States	160 mL	
	2 to 3 m tall		320 mL	
Gorse	Spring to mid summer		170 mL or 340 mL	Add a 100% concentrate non-ionic wetting agent at rate of 125-mL/100L water. Retreatment of regrowth may be necessary. Use higher water rate on older hardened off plants.
Horehound	Rosette	Tas only	170 mL	
Saffron thistle	Up to bud stage		80 mL	
Tiger pear	Active growth	All States	3 L	
Wattles, including Silver wattle Black wattle	Seedlings, 1 to 2 m tall		160 mL	
	2 to 3 m tall		320 mL	



**Table B: Aerial Application**  
**Helicopter NSW, SA, Tas, Vic and WA only**  
**Helicopter or fixed wing aircraft (Qld only)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /ha</b>	<b>CRITICAL COMMENTS</b>
Blackberry	Late spring to autumn	All States	4.8 L	AVOID overspray/drift onto waterways.

**Table C: Controlled Droplet Application (C.D.A.)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /1L water</b>	<b>CRITICAL COMMENTS</b>
Blackberry	Late spring to autumn	All States	170 mL	

**Table D: Low Volume High Concentrate Application Techniques**  
**(Gas Powered Gun, Sprinkler Sprayer)**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /10 L water</b>	<b>CRITICAL COMMENTS</b>
Blackberry	Late spring to autumn	All States	280 mL	
Eucalypt seedlings	1 to 2 m tall		400 mL	

**Table E: Basal Bark and Cut Stump Treatment**

See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /60 L diesel</b>	<b>CRITICAL COMMENTS</b>
African boxthorn	<b>Basal bark:</b> Plants up to 5 cm basal diameter  <b>Cut stump:</b> Plants up to and in excess of basal bark sizes	All States	2 L	
Australian blackthorn			1 L	
Broom		Tas only	1.25 L	
Brown salwood		All States	0.5 L	
Bitter bark		NSW and Qld only	1 L	
Castor oil		All States		
Chinese apple				
Dawson gum		Qld only	2 L	
Eucalypts (except Dawson gum)		All States	1 L	
False sandalwood				
Green wattle				
Lantana				
Needlewood				
Olive		SA only	2 L	
		NSW only	4 L	

**Table E: Basal Bark and Cut Stump Treatment (Continued)**See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>AGRICULTURAL NON-CROP AREAS, COMMERCIAL AND INDUSTRIAL AREAS, FORESTS, PASTURES AND RIGHTS-OF-WAY.</b>					
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /60 L diesel</b>	<b>CRITICAL COMMENTS</b>	
Paperbark teatree Rubbervine Silver wattle	Plants up to and in excess of basal bark sizes	All States	1 L		
Sweet briar Yellow-wood			2 L		
Camphor laurel	<b>Basal bark:</b> Plants up to 10 cm basal diameter	NSW and Qld only	1L		
Common prickly pear		All States	0.8 L		
Groundsel bush ( <i>Baccharis halimifolia</i> )	<b>Cut stump</b> Plants up to and in excess of basal bark sizes	All States	0.5 L		Treat from early summer rains to end of April when regrowth is apparent.
Prickly acacia			5 L		Treatment may be carried out at any time of the year.
Privet			0.8 L		Apply as a thorough foliage spray.
Smooth tree pear Tiger pear			1 L		
Tree of heaven					
<b>FENCELINES AND FIRE TRAILS only.</b>					
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /60 L diesel</b>	<b>CRITICAL COMMENTS</b>	
Broadleaf hopbush, Narrowleaf hopbush, Turpentine bush	<b>Basal bark:</b> Plants up to 10 cm basal diameter	NSW only	1 L		

**2. CROPPING/FALLOW SITUATIONS**See **GENERAL INSTRUCTIONS - APPLICATION** section for application method details.

<b>FALLOW, STUBBLE, FIREBREAKS.</b>				
<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /ha</b>	<b>CRITICAL COMMENTS</b>
Camel (Afghan, Bitter) melon ( <i>Citrullus lanatus</i> )	Up to 20 cm diameter	NSW, Qld, SA, Vic and WA only	120 mL ①	There are some strains of melon that are not controlled. Contact your Conquest Agrochemicals representative for more information. ①Add a crop oil such as Uptake* Spraying Oil at 500 mL/100 L water or D-C-Trate® Crop Oil at 1 L/100 L water. <b>DO NOT</b> use oils when tank mixing with Conquest Glyphosate 450 Herbicide. See <b>COMPATIBILITY</b> section. This mixture is not selective to grasses. When using Conquest Maca 600 and Conquest Glyphosate 450 Herbicide in aerial application, observance of a buffer zone of 150 metres to protect native tree species is required.
	Runners from 20 to 40 cm diameter		160 mL ①	
Prickly paddy melon ( <i>Cucumis myriocarpus</i> )	Up to 20 cm diameter		80 mL ①	
	Runners from 20 to 40 cm diameter		160 mL ①	

**SORGHUM****(Apply between 4 to 6 leaf stage, when secondary roots have developed)**

<b>WEEDS CONTROLLED</b>	<b>WEED GROWTH STAGE</b>	<b>STATE</b>	<b>RATE /ha</b>	<b>CRITICAL COMMENTS</b>
Prickly paddy melon	Up to 20 cm diameter	NSW and Qld only	80 mL	<b>DO NOT</b> add crop oils, as severe crop damage may occur. Fusing of sorghum prop roots may be observed and this may be worse under stress (e.g. root disease). Should only be mixed with Starane* 200 Herbicide and atrazine (500 g/L flowable only) for increased weed spectrum.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**IN TASMANIA FOR BLACKBERRY: DO NOT** treat bushes carrying mature or near mature fruit.

**FOR NATIVE VEGETATION:** Use of Conquest Maca 600 on native vegetation must be done in accordance with STATE and/or LOCAL legislation.

**WITHHOLDING PERIODS:**

**PASTURE, SORGHUM: NOT REQUIRED WHEN USED AS DIRECTED.**

**GENERAL INSTRUCTIONS****RESISTANT WEEDS WARNING**

<b>GROUP</b>	<b>I</b>	<b>HERBICIDE</b>
--------------	----------	------------------

Conquest Maca 600 Herbicide is a member of the Pyridines group of herbicides. The product has the disrupters of plant cell growth mode of action. For weed resistance management, the product is a Group I herbicide. Some naturally-occurring weed biotypes resistant to the product and other disrupters of plant cell growth herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other disrupters of plant cell growth herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Agrochemicals Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Strategies to minimise the risk of herbicide resistance are available. Contact your farm chemical supplier, consultant, local Department of Agriculture, or local Conquest Agrochemicals representative.

**MIXING**

Half fill the spray unit with water and add the required amount of Conquest Maca 600. Add the remaining water with the agitator running. If required, then add crop oils or wetters (surfactants). Maintain mechanical or by-pass agitation in the spray tank during spraying. Only mix sufficient solution for immediate daily use and avoid storing.

**Basal Bark and Cut Stump Application:** Quarter fill the spray unit or mixing container with diesel and add the required amount of Conquest Maca 600. Add the remaining diesel and shake or agitate thoroughly to mix the contents. Periodically shake or agitate to stop product settling out. Only mix sufficient solution for immediate daily use and avoid storing.

## APPLICATION

### SHAKE WELL BEFORE USE

#### 1. WOODY WEED SITUATIONS

Weeds need to be actively growing for herbicides to have optimum effect. Delay treatment until all regrowth has had time to grow to one metre high in situations, which have been bulldozed, slashed, burnt, ploughed or areas having a previous chemical treatment.

#### A. High Volume Spraying

Thorough coverage of foliage and stems to the point of runoff is essential, however, avoid excess spraying, which is wasteful of chemical.

##### Hand Gun

Apply the recommended mix to give full coverage of leaves and stems through a No. 6 to 8 tip at 700 to 1500 kPa.

A spray volume of 3,000 to 4,000 L per infested hectare (30 to 40 L/100 m<sup>2</sup>) should be used on the weed infestation.

##### Knapsack

Used for smaller infestations, where penetration of the canopy is not essential.

A spray volume of 3 to 4 L/10 m<sup>2</sup> of infested area should be used.

#### B. Aerial Application

Apply in 100 to 200 L water/ha. Use a calibrated aircraft to apply in half overlap passes. Nozzle configurations should produce droplets of not less than 250 to 350 micron diameter (e.g. D8/45).

The potential for damage from drift can be greatly reduced by avoiding unsuitable spraying conditions and using spray pressure and nozzles to minimise the production of small droplets.

**DO NOT** spray when wind exceeds 10 km/hr and/or air temperature reaches 30°C.

Human flagging is not authorised unless protected by engineering controls such as vehicles with cabs.

#### C. Controlled Droplet Application (C.D.A.)

Results similar to high volume spraying can be obtained using Micron Herbi<sup>®</sup> or similar equipment. Select a nozzle to give a flow rate of 2 mL/sec and sweeping action of approximately 1 m/sec to ensure a droplet density of 20/cm<sup>2</sup>. Use a marking agent as recommended by the equipment manufacturers, to check spray coverage. Also, consult directions provided with C.D.A. unit.

#### D. Low Volume High Concentrate Application Techniques

Good control will be achieved, similar to high volume application, where bush size enables good coverage of the bush. Use a marking agent, as recommended by the equipment manufacturers, to check spray coverage.

**Gas powered gun:** Apply 50 mL shots to obtain uniform coverage of 4 to 5 m<sup>2</sup> of surface area of bush. This relates to 20 droplets/cm<sup>2</sup> of leaf surface.

**Sprinkler sprayer:** This technique involves using a micro sprinkler, which is connected to a hollow fibreglass rod attached to a pressure knapsack sprayer. Use at low pressures (50 to 200 kPa) and apply with a slow sweeping action over the top of the plants ensuring even coverage on the leaves.

## **E. Basal Bark and Cut Stump Treatment**

Mix Conquest Maca 600 in diesel. The use of diesel as a herbicide carrier may affect the rubber seals in some sprayers. To avoid this, use sprayers, which use Viton<sup>®</sup> seals and fittings. When using Conquest Maca 600 with diesel, nitrile gloves should be worn instead of rubber gloves.

### **Basal Bark Method**

**DO NOT** apply to wet stems as this can repel the diesel mixture.

Apply only with hand-directed equipment such as a pressure sprayer or a paintbrush.

Spray equipment should be used at low pressures, up to 200 kPa, to avoid excessive splashing or drift.

Species with old, rough bark require more thorough wetting than smooth barked species.

Liberally spray or paint the bark around the stem from ground level up to 30 cm high, wetting thoroughly to the point of runoff (unless otherwise stated).

### **Cut Stump Method**

Stems should be cut less than 15 cm above the ground.

**Immediately** apply Conquest Maca 600/diesel mixture **liberally** to the **freshly** cut stump by spray or painting the cut surface and sides of the stem.

## **2. CROPPING/FALLOW SITUATIONS**

### **A. Boom Application**

Application of Conquest Maca 600 in a minimum spray volume of 50 L/ha is recommended. Flat fan nozzles are recommended, using pressures in the range of 200-300 kPa. Boom height must be set to ensure double overlap of nozzle patterns.

### **B. Aerial Application**

**DO NOT** allow Conquest Maca 600 to physically drift onto desirable plants.

Aircraft may be used to apply Conquest Maca 600 in fallow situations, when ground application equipment cannot be used due to prolonged wet conditions.

A minimum spray volume of 35 L/ha should be used with flat fan and CP nozzle configurations.

**DO NOT** apply Conquest Maca 600 by aircraft when wind exceeds 10 km/hr and/or air temperature reaches 30°C.

Droplets with an average size (or Volume Mean Diameter) of 250 to 350 micron diameter are recommended.

Human flagging is not authorised unless protected by engineering controls such as vehicles with cabs.

## **CLEANING SPRAY EQUIPMENT**

### **WATER-BASED CLEANING**

#### **Rinsing**

After using Conquest Maca 600, empty the spray unit completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain the spray unit and clean any filters in the tank, pump, lines, hoses and nozzles.

After cleaning the spray unit as above, quarter fill with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

## Decontamination

Before spraying cotton and other sensitive crops, with equipment that has been used to apply Conquest Maca 600, see PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS section.

Wash the tank and rinse the system as above. Then quarter fill the tank and add an alkali detergent (e.g. liquid SURF<sup>®</sup>, OMO<sup>®</sup>, DRIVE<sup>®</sup> at 500 mL/100 L of water or the powder equivalent at 500 g/100 L of water) and circulate throughout the system for at least 15 minutes.

Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain.

**Rinse water should be discharged onto a designated disposal area or, if this is unavailable, onto unused land away from desirable plants and watercourses.**

## DIESEL-BASED CLEANING

### Rinsing

After using Conquest Maca 600, empty the spray unit completely and drain the whole system. Thoroughly wash inside the spray unit using a pressure hose. Drain the spray unit and clean any filters in the tank, pump, lines, hoses and nozzles.

On completion of spraying, use a degreaser such as Caltex Kwik-D-Grease<sup>®</sup> to remove traces of diesel from the sprayer. Rinse tank and spray through the nozzles with water to remove degreaser.

### Decontamination

After the above, quarter fill the tank with clean water and add an alkali detergent (e.g. liquid SURF<sup>®</sup>, OMO<sup>®</sup>, OMOMATIC<sup>®</sup>, DRIVE<sup>®</sup> at 50 mL/10 L of water or the powder equivalent at 50 g/10 L of water). Shake or operate spray to circulate the washing solution throughout the sprayer and spray the solution through the nozzle. Rinse well with clean water to remove detergent.

To clean brushes and container, spray liberally with degreaser. Hose off thoroughly with clean water and repeat using detergents (see above).

**DO NOT** use this equipment for any other purpose.

## COMPATIBILITY

### FALLOW SITUATIONS:

- Conquest Maca 600 is compatible with the following products:
  - Lontrel\* Herbicide
  - Starane\* 200 Herbicide
  - Conquest Glyphosate 450 Herbicide
  - Roundup<sup>®</sup> CT Xtra Broadacre Herbicide
  - Touchdown<sup>®</sup> Broadacre
  - 2,4-D amine or 2,4-D ester
  - glyphosate
  - Lorsban\* 500 EC Insecticide
- When mixing with Conquest Glyphosate 450 Herbicide in fallow, refer to the Conquest Glyphosate 450 Herbicide label for use rate and adjuvants recommended. **DO NOT** use Uptake\* Spraying Oil or D-C-Trate<sup>®</sup> Crop Oil.

### SORGHUM:

- Conquest Maca 600 is compatible with:
  - Starane 200 Herbicide
  - atrazine (500 g/L flowable product only)

PVC gloves are not recommended for Conquest Maca 600 Herbicide, therefore, when tank mixing with products that need to be handled with PVC gloves, workers should wear nitrile gloves.

**MINIMUM RECROPPING PERIODS**

Before using Conquest Maca 600 in tank mixes with other herbicides, check the plant-back information on all product labels. The time between spraying and planting will be determined by the product with the longest plant-back period.


Observe the following recropping periods for Conquest Maca 600:

- |                                   |         |
|-----------------------------------|---------|
| • wheat, barley, sorghum, maize   | 7 days  |
| • chickpeas, soybeans, sunflowers | 7 days  |
| • cotton                          | 14 days |

**PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS**

Crops susceptible to Conquest Maca 600 include, but are not limited to: peas, lupins, lucerne, navy beans, soybeans and other legumes; cotton, fruit, hops, ornamentals, shade trees and *Pinus* spp, potatoes, safflower, sugarbeet, sunflower, tobacco, tomatoes, vegetables and vines.

Conquest Maca 600 is damaging to susceptible crops during both growing and dormant periods. Grasses are normally unaffected and establish quickly after treatment.

**DO NOT** allow physical spray drift onto waterways, native vegetation and susceptible crops. When using Conquest Maca 600 and Conquest Glyphosate 450 Herbicide  aerial application in fallow situations, observance of a buffer zone of 150 metres to protect native tree species is required.

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

**PROTECTION OF LIVESTOCK**

Poisonous plants may become more palatable after spraying and stock should be kept away from these plants until they have died down.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

This product is highly toxic to fish and other aquatic organisms. **DO NOT** contaminate streams, rivers or waterways with the chemical or used containers.

**STORAGE AND DISPOSAL**

Store in closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. **DO NOT** store near food, feedstuffs, fertilisers or seed. **DO NOT** re-use containers.

Triple rinse or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**SMALL SPILL MANAGEMENT**

Wear protective equipment (see SAFETY DIRECTIONS). Apply absorbent material such as earth, sand, clay granules or cat litter to the spill. Sweep up material for disposal when absorption is completed and contain in a refuse vessel for disposal (see STORAGE AND DISPOSAL section). If necessary wash the spill area with an alkali detergent and water and absorb the wash liquid as above for disposal.

## **SAFETY DIRECTIONS**

Avoid contact with eyes and skin. DO NOT inhale spray mist. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

## **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (Ph: 13 1126). If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

## **MATERIAL SAFETY DATA SHEET**

Additional information is listed on the Material Safety Data Sheet which is available from the supplier.

## **CONDITIONS OF SALE**

Conquest Agrochemicals Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

<p>IN EMERGENCY DIAL</p> <p><b>000,</b></p> <p><b>POLICE OR FIRE BRIGADE</b></p>	<p>FOR SPECIALIST ADVICE IN AN EMERGENCY CALL</p> <p><b>1800 033 111</b></p> <p><b>(24 HOURS)</b></p>
----------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------

## **Conquest Agrochemicals Pty Ltd.**

ABN 84 098 814 932

Suite 10, 41 Walters Drive

Herdsmen Business Park

Osborne Park WA 6017

Telephone (08) 9244 1268

Facsimile: (08) 9446 4696

APVMA approval No: 58112/0104