



Company Name: JUROX PTY LIMITED
Product Name: COBALEX 2000 B12 INJECTION

APVMA Approval No: 67608/102086



Label Name:	COBALEX 2000 B12 INJECTION
Signal Headings:	FOR ANIMAL TREATMENT ONLY
Constituent Statements:	2 mg/mL HYDROXOCOBALAMIN (as the acetate)
Claims:	For the treatment and control of vitamin B12 deficiency in sheep and cattle.
Net Contents:	500mL
Directions for Use:	
Restraints:	
Contraindications:	This product is contraindicated for use unless cobalt deficiency is confirmed. This product is contraindicated for use in sheep that are dehydrated or unwell.
Precautions:	
Side Effects:	
Dosage and Administration:	If stored below 30°C (room temperature) and protected from light, contents may be used for 60 days after initial broaching of the vial. Discard any unused portion after 60 days. LAMBS: 0.5 mL at marking or weaning SHEEP: 1.0 - 1.5 mL pre lambing CALVES: 2 - 3 mL from 2 months of age COWS: 4 - 6 mL pre calving Give dose by subcutaneous injection only.

The frequency of dosing should be related to the severity of the deficiencies. For instance, for sheep in extremely deficient areas, repeat dose every 2 - 3 months. In less deficient areas, repeat dose every 5 - 6 months or as found necessary. Consult your animal health adviser for advice specific to your flock or herd.

General Directions:

Caution: avoid carcass damage

1. Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. Use shortest needle possible, certainly not exceeding 15 mm.
5. As far as possible, avoid injection of animals during wet weather or under dusty conditions.
6. This product should be injected only under the skin.
7. If possible, inject high on the neck behind the ear.

Withholding Periods:

WITHHOLDING PERIOD: Zero (0) days

Trade Advice:

TRADE ADVICE - Export Slaughter Interval: Zero (0) days

Safety Directions:**First Aid Instructions:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 1126.

First Aid Warnings:**Additional User Safety:****Environmental Statements:****Disposal:**

Dispose of container by wrapping in paper and putting in garbage. Discarded needles should immediately be placed in a designated and appropriately labelled 'sharps' container.

Storage:

Store below 30°C (room temperature). Protect from light. Do not freeze. If stored below 30°C (room temperature) and protected from light, contents may be used for 60 days after initial broaching of the vial. Discard any unused portion after 60 days.