

Virbac (Australia) Pty Limited	Page 1 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE
APVMA:	66660 / XXXXX
Version Date:	19 April 2013
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L

Text above this line or in italics between parallel lines is not included in the label copy.

IMMEDIATE CONTAINER – Front Panel

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CYDECTIN[®]
PLUS FLUKE POUR-ON FOR CATTLE

ACTIVE CONSTITUENTS: 5 g/L MOXIDECTIN
200 g/L TRICLABENDAZOLE

For the treatment and control of moxidectin sensitive strains of gastrointestinal nematodes and triclabendazole sensitive strains of liver fluke (*Fasciola hepatica*) of cattle.



RLP
Approved



CONTENTS: 500 mL / 1 LITRE / 1.1 LITRES / 2 LITRES / 2.2 LITRES / 5 LITRES / 5.5 LITRES / 7.5 LITRES / 10 LITRES / 11 LITRES / 15 LITRES / 20 LITRES



IMMEDIATE CONTAINER – Back Panel

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE contains moxidectin, a second generation macrocyclic lactone endectocide that is effective against parasites sensitive to this family and triclabendazole, which is effective against liver fluke. **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** does not cause irritation to the skin or hide damage when used as directed.

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE is effective in the treatment and control of the following parasites affecting cattle:

Triclabendazole susceptible Liver fluke (late immature (6 to 8 weeks) and adult):

Fasciola hepatica

Moxidectin susceptible Gastrointestinal (adult and larval L4):

Small Brown Stomach Worm (*Ostertagia ostertagi*), Stomach Hair Worm (*Trichostrongylus axei*) and Small Intestinal Worm (*Cooperia* spp.).

Moxidectin susceptible Gastrointestinal (adult): Barbers Pole Worm (*Haemonchus placei*), Nodule worm (*Oesophgostomum* spp.), Thin-necked Intestinal Worm (*Nematodirus* spp.) and Hookworm (*Bunostomum* spp.).

Virbac (Australia) Pty Limited		Page 2 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

PROTECTION PERIOD

A single treatment at the recommended dose rate prevents re-infection with moxidectin susceptible Barber's Pole Worm (*Haemonchus contortus*) and Small Brown Stomach Worm (*Ostertagia ostertagi*) for 14 days.

Parasite Control Programs

Effective parasite control for maximum productivity is an essential part of farm management. Follow recommended parasite management practices to reduce the potential for resistance to develop. Resistance may develop to any chemical.

DIRECTIONS FOR USE:

RESTRAINT:

DO NOT USE on dairy cattle which are producing or may in the future produce milk for human consumption or processing.

DOSE RATE

0.5 mg moxidectin and 20 mg triclabendazole/kg liveweight (1 mL per 10 kg liveweight).

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE should be applied along the top of the back from the withers to the base of the tail. The product is ready to use. Administer the product according to the dosing table below.

Dosing Table

(500 mL Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 500 mL	Liveweight (kg)	Dose Volume (mL)	Doses per pack 500 mL
50-55	5.5	90	181-190	19	26
56-60	6	83	191-200	20	25
61-65	6.5	76	201-220	22	22
66-70	7	71	221-240	24	20
71-75	7.5	66	241-260	26	19
76-80	8	62	261-280	28	17
81-85	8.5	58	281-300	30	16
86-90	9	55	301-320	32	15
91-95	9.5	52	321-340	34	14
96-100	10	50	341-360	36	13
101-110	11	45	361-380	38	13
111-120	12	41	381-420	42	11
121-130	13	38	421-460	46	10
131-140	14	35	461-500	50	10
141-150	15	33	501-550	55	9
151-160	16	31	551-600	60	8
161-170	17	29	601-650	65	7
171-180	18	27			

Virbac (Australia) Pty Limited		Page 3 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(1 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 1 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 1 L
50-55	5.5	181	181-190	19	52
56-60	6	166	191-200	20	50
61-65	6.5	153	201-220	22	45
66-70	7	142	221-240	24	41
71-75	7.5	133	241-260	26	38
76-80	8	125	261-280	28	35
81-85	8.5	117	281-300	30	33
86-90	9	111	301-320	32	31
91-95	9.5	105	321-340	34	29
96-100	10	100	341-360	36	27
101-110	11	90	361-380	38	26
111-120	12	83	381-420	42	23
121-130	13	76	421-460	46	21
131-140	14	71	461-500	50	20
141-150	15	66	501-550	55	18
151-160	16	62	551-600	60	16
161-170	17	58	601-650	65	15
171-180	18	55			

(1.1 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 1.1 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 1.1 L
50-55	5.5	200	181-190	19	57
56-60	6	183	191-200	20	55
61-65	6.5	169	201-220	22	50
66-70	7	157	221-240	24	45
71-75	7.5	146	241-260	26	42
76-80	8	137	261-280	28	39
81-85	8.5	129	281-300	30	36
86-90	9	122	301-320	32	34
91-95	9.5	115	321-340	34	32
96-100	10	110	341-360	36	30
101-110	11	100	361-380	38	28
111-120	12	91	381-420	42	26
121-130	13	84	421-460	46	23
131-140	14	78	461-500	50	22
141-150	15	73	501-550	55	20
151-160	16	68	551-600	60	18
161-170	17	64	601-650	65	16
171-180	18	61			

Virbac (Australia) Pty Limited		Page 4 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(2 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 2 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 2 L
50-55	5.5	363	181-190	19	105
56-60	6	333	191-200	20	100
61-65	6.5	307	201-220	22	90
66-70	7	285	221-240	24	83
71-75	7.5	266	241-260	26	76
76-80	8	250	261-280	28	71
81-85	8.5	235	281-300	30	66
86-90	9	222	301-320	32	62
91-95	9.5	210	321-340	34	58
96-100	10	200	341-360	36	55
101-110	11	181	361-380	38	52
111-120	12	166	381-420	42	47
121-130	13	153	421-460	46	43
131-140	14	142	461-500	50	40
141-150	15	133	501-550	55	36
151-160	16	125	551-600	60	33
161-170	17	117	601-650	65	30
171-180	18	111			

(2.2 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 2.2 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 2.2 L
50-55	5.5	400	181-190	19	115
56-60	6	366	191-200	20	110
61-65	6.5	338	201-220	22	100
66-70	7	314	221-240	24	91
71-75	7.5	293	241-260	26	84
76-80	8	275	261-280	28	78
81-85	8.5	258	281-300	30	73
86-90	9	244	301-320	32	68
91-95	9.5	231	321-340	34	64
96-100	10	220	341-360	36	61
101-110	11	200	361-380	38	57
111-120	12	183	381-420	42	52
121-130	13	169	421-460	46	47
131-140	14	157	461-500	50	44
141-150	15	146	501-550	55	40
151-160	16	137	551-600	60	36
161-170	17	129	601-650	65	33
171-180	18	122			

Virbac (Australia) Pty Limited		Page 5 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(5 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 5 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 5 L
50-55	5.5	909	181-190	19	263
56-60	6	833	191-200	20	250
61-65	6.5	769	201-220	22	227
66-70	7	714	221-240	24	208
71-75	7.5	666	241-260	26	192
76-80	8	625	261-280	28	178
81-85	8.5	588	281-300	30	166
86-90	9	555	301-320	32	156
91-95	9.5	526	321-340	34	147
96-100	10	500	341-360	36	138
101-110	11	454	361-380	38	131
111-120	12	416	381-420	42	119
121-130	13	384	421-460	46	108
131-140	14	357	461-500	50	100
141-150	15	333	501-550	55	90
151-160	16	312	551-600	60	83
161-170	17	294	601-650	65	76
171-180	18	277			

(5.5 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 5.5 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 5.5 L
50-55	5.5	1000	181-190	19	289
56-60	6	916	191-200	20	275
61-65	6.5	846	201-220	22	250
66-70	7	785	221-240	24	229
71-75	7.5	733	241-260	26	211
76-80	8	687	261-280	28	196
81-85	8.5	647	281-300	30	183
86-90	9	611	301-320	32	171
91-95	9.5	578	321-340	34	161
96-100	10	550	341-360	36	152
101-110	11	500	361-380	38	144
111-120	12	458	381-420	42	130
121-130	13	423	421-460	46	119
131-140	14	392	461-500	50	110
141-150	15	366	501-550	55	100
151-160	16	343	551-600	60	91
161-170	17	323	601-650	65	84
171-180	18	305			

Virbac (Australia) Pty Limited		Page 6 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(7.5 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 7.5 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 7.5 L
50-55	5.5	1363	181-190	19	394
56-60	6	1250	191-200	20	375
61-65	6.5	1153	201-220	22	340
66-70	7	1071	221-240	24	312
71-75	7.5	1000	241-260	26	288
76-80	8	937	261-280	28	267
81-85	8.5	882	281-300	30	250
86-90	9	833	301-320	32	234
91-95	9.5	789	321-340	34	220
96-100	10	750	341-360	36	208
101-110	11	681	361-380	38	197
111-120	12	625	381-420	42	178
121-130	13	576	421-460	46	163
131-140	14	535	461-500	50	150
141-150	15	500	501-550	55	136
151-160	16	468	551-600	60	125
161-170	17	441	601-650	65	115
171-180	18	416			

(10 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 10 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 10 L
50-55	5.5	1818	181-190	19	526
56-60	6	1666	191-200	20	500
61-65	6.5	1538	201-220	22	454
66-70	7	1428	221-240	24	416
71-75	7.5	1333	241-260	26	384
76-80	8	1250	261-280	28	357
81-85	8.5	1176	281-300	30	333
86-90	9	1111	301-320	32	312
91-95	9.5	1052	321-340	34	294
96-100	10	1000	341-360	36	277
101-110	11	909	361-380	38	263
111-120	12	833	381-420	42	238
121-130	13	769	421-460	46	217
131-140	14	714	461-500	50	200
141-150	15	666	501-550	55	181
151-160	16	625	551-600	60	166
161-170	17	588	601-650	65	153
171-180	18	555			

Virbac (Australia) Pty Limited	Page 7 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE
APVMA:	66660 / XXXXX
Version Date:	19 April 2013
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L

Text above this line or in italics between parallel lines is not included in the label copy.

(11 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 11 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 11 L
50-55	5.5	2000	181-190	19	578
56-60	6	1833	191-200	20	550
61-65	6.5	1692	201-220	22	500
66-70	7	1571	221-240	24	458
71-75	7.5	1466	241-260	26	423
76-80	8	1375	261-280	28	392
81-85	8.5	1294	281-300	30	366
86-90	9	1222	301-320	32	343
91-95	9.5	1157	321-340	34	323
96-100	10	1100	341-360	36	305
101-110	11	1000	361-380	38	289
111-120	12	916	381-420	42	261
121-130	13	846	421-460	46	239
131-140	14	785	461-500	50	220
141-150	15	733	501-550	55	200
151-160	16	687	551-600	60	183
161-170	17	647	601-650	65	169
171-180	18	611			

(15 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 15 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 15 L
50-55	5.5	2727	181-190	19	789
56-60	6	2500	191-200	20	750
61-65	6.5	2307	201-220	22	681
66-70	7	2142	221-240	24	625
71-75	7.5	2000	241-260	26	576
76-80	8	1875	261-280	28	535
81-85	8.5	1764	281-300	30	500
86-90	9	1666	301-320	32	468
91-95	9.5	1578	321-340	34	441
96-100	10	1500	341-360	36	416
101-110	11	1363	361-380	38	394
111-120	12	1250	381-420	42	357
121-130	13	1153	421-460	46	326
131-140	14	1071	461-500	50	300
141-150	15	1000	501-550	55	272
151-160	16	937	551-600	60	250
161-170	17	882	601-650	65	230
171-180	18	833			

Virbac (Australia) Pty Limited		Page 8 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(20 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 20 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 20 L
50-55	5.5	3636	181-190	19	1052
56-60	6	3333	191-200	20	1000
61-65	6.5	3076	201-220	22	909
66-70	7	2857	221-240	24	833
71-75	7.5	2666	241-260	26	769
76-80	8	2500	261-280	28	714
81-85	8.5	2352	281-300	30	666
86-90	9	2222	301-320	32	625
91-95	9.5	2105	321-340	34	588
96-100	10	2000	341-360	36	555
101-110	11	1818	361-380	38	526
111-120	12	1666	381-420	42	476
121-130	13	1538	421-460	46	434
131-140	14	1428	461-500	50	400
141-150	15	1333	501-550	55	363
151-160	16	1250	551-600	60	333
161-170	17	1176	601-650	65	307
171-180	18	1111			

Cattle in excess of 650 kg should be dosed at 1 mL per 10 kg liveweight. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc.). Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Do not underdose. Check accuracy of dose delivery of applicator regularly.

RE-TREATMENT INTERVAL: DO NOT retreat animals with this product for 160 days after last treatment.

Rainfast

The efficacy of **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application.

Ease of Use

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE comes ready to use.

Non Irritant

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE has not been observed to cause irritation to the skin or hide damage when used as directed.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 50 days before slaughter for human consumption.

MILK: DO NOT USE on dairy cattle which are producing or may in the future produce milk for human consumption or processing.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 160 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before

Virbac (Australia) Pty Limited	Page 9 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE
APVMA:	66660 / XXXXX
Version Date:	19 April 2013
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L

Text above this line or in italics between parallel lines is not included in the label copy.

using this product, confirm the current ESI from either Virbac (Australia) Pty Ltd 1800 009 847 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Do not inhale. Will irritate the eyes. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves and face shield or goggles. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

Additional information is listed in the Material Safety Data Sheet.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

500 mL, 1 L Pack

DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

ALL OTHER PACK SIZES

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced but may be taken separately.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Moxidectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

STORAGE

Store below 30°C (Room temperature) and protect from light.

APVMA Approval No.: 66660/59567

BN:
EXP:



VIRBAC (AUSTRALIA) PTY. LIMITED
361 Horsley Road, Milperra NSW 2214
Freecall Customer Service Line: 1800 009 847

Virbac (Australia) Pty Limited		Page 10 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

www.virbac.com.au

[®] Registered trademark of Virbac (Australia) Pty Limited.

Virbac (Australia) Pty Limited	Page 11 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE
APVMA:	66660 / XXXXX
Version Date:	19 April 2013
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L

Text above this line or in italics between parallel lines is not included in the label copy.

OPTIONAL CARTON – Front Panel

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CYDECTIN®
PLUS FLUKE POUR-ON FOR CATTLE

ACTIVE CONSTITUENTS: 5 g/L MOXIDECTIN
200 g/L TRICLABENDAZOLE

For the treatment and control of moxidectin sensitive strains of gastrointestinal nematodes and triclabendazole sensitive strains of liver fluke (*Fasciola hepatica*) of cattle.

CONTENTS: 500 mL / 1 LITRE / 1.1 LITRES / 2 LITRES / 2.2 LITRES / 5 LITRES / 5.5 LITRES / 7.5 LITRES / 10 LITRES / 11 LITRES / 15 LITRES / 20 LITRES



OPTIONAL CARTON – Back Panel

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE contains moxidectin, a second generation macrocyclic lactone endectocide that is effective against parasites sensitive to this family and triclabendazole, which is effective against liver fluke. **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** does not cause irritation to the skin or hide damage when used as directed.

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE is effective in the treatment and control of the following parasites affecting cattle:

Triclabendazole susceptible Liver fluke (late immature (6 to 8 weeks) and adult):

Fasciola hepatica

Moxidectin susceptible Gastrointestinal (adult and larval L4):

Small Brown Stomach Worm (*Ostertagia ostertagi*), Stomach Hair Worm (*Trichostrongylus axei*) and Small Intestinal Worm (*Cooperia* spp.).

Moxidectin susceptible Gastrointestinal (adult): Barbers Pole Worm (*Haemonchus placei*), Nodule worm (*Oesophgostomum* spp.), Thin-necked Intestinal Worm (*Nematodirus* spp.) and Hookworm (*Bunostomum* spp.).

Virbac (Australia) Pty Limited		Page 12 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

PROTECTION PERIOD

A single treatment at the recommended dose rate prevents re-infection with moxidectin susceptible Barber's Pole Worm (*Haemonchus contortus*) and Small Brown Stomach Worm (*Ostertagia ostertagi*) for 14 days.

Parasite Control Programs

Effective parasite control for maximum productivity is an essential part of farm management. Follow recommended parasite management practices to reduce the potential for resistance to develop. Resistance may develop to any chemical.

DIRECTIONS FOR USE:

RESTRAINT:

DO NOT USE on dairy cattle which are producing or may in the future produce milk for human consumption or processing.

DOSE RATE

0.5 mg moxidectin and 20 mg triclabendazole/kg liveweight (1 mL per 10 kg liveweight).

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE should be applied along the top of the back from the withers to the base of the tail. The product is ready to use. Administer the product according to the dosing table below.

Dosing Table

(500 mL Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 500 mL	Liveweight (kg)	Dose Volume (mL)	Doses per pack 500 mL
50-55	5.5	90	181-190	19	26
56-60	6	83	191-200	20	25
61-65	6.5	76	201-220	22	22
66-70	7	71	221-240	24	20
71-75	7.5	66	241-260	26	19
76-80	8	62	261-280	28	17
81-85	8.5	58	281-300	30	16
86-90	9	55	301-320	32	15
91-95	9.5	52	321-340	34	14
96-100	10	50	341-360	36	13
101-110	11	45	361-380	38	13
111-120	12	41	381-420	42	11
121-130	13	38	421-460	46	10
131-140	14	35	461-500	50	10
141-150	15	33	501-550	55	9
151-160	16	31	551-600	60	8
161-170	17	29	601-650	65	7
171-180	18	27			

Virbac (Australia) Pty Limited		Page 13 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(1 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 1 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 1 L
50-55	5.5	181	181-190	19	52
56-60	6	166	191-200	20	50
61-65	6.5	153	201-220	22	45
66-70	7	142	221-240	24	41
71-75	7.5	133	241-260	26	38
76-80	8	125	261-280	28	35
81-85	8.5	117	281-300	30	33
86-90	9	111	301-320	32	31
91-95	9.5	105	321-340	34	29
96-100	10	100	341-360	36	27
101-110	11	90	361-380	38	26
111-120	12	83	381-420	42	23
121-130	13	76	421-460	46	21
131-140	14	71	461-500	50	20
141-150	15	66	501-550	55	18
151-160	16	62	551-600	60	16
161-170	17	58	601-650	65	15
171-180	18	55			

(1.1 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 1.1 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 1.1 L
50-55	5.5	200	181-190	19	57
56-60	6	183	191-200	20	55
61-65	6.5	169	201-220	22	50
66-70	7	157	221-240	24	45
71-75	7.5	146	241-260	26	42
76-80	8	137	261-280	28	39
81-85	8.5	129	281-300	30	36
86-90	9	122	301-320	32	34
91-95	9.5	115	321-340	34	32
96-100	10	110	341-360	36	30
101-110	11	100	361-380	38	28
111-120	12	91	381-420	42	26
121-130	13	84	421-460	46	23
131-140	14	78	461-500	50	22
141-150	15	73	501-550	55	20
151-160	16	68	551-600	60	18
161-170	17	64	601-650	65	16
171-180	18	61			

Virbac (Australia) Pty Limited		Page 14 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(2 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 2 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 2 L
50-55	5.5	363	181-190	19	105
56-60	6	333	191-200	20	100
61-65	6.5	307	201-220	22	90
66-70	7	285	221-240	24	83
71-75	7.5	266	241-260	26	76
76-80	8	250	261-280	28	71
81-85	8.5	235	281-300	30	66
86-90	9	222	301-320	32	62
91-95	9.5	210	321-340	34	58
96-100	10	200	341-360	36	55
101-110	11	181	361-380	38	52
111-120	12	166	381-420	42	47
121-130	13	153	421-460	46	43
131-140	14	142	461-500	50	40
141-150	15	133	501-550	55	36
151-160	16	125	551-600	60	33
161-170	17	117	601-650	65	30
171-180	18	111			

(2.2 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 2.2 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 2.2 L
50-55	5.5	400	181-190	19	115
56-60	6	366	191-200	20	110
61-65	6.5	338	201-220	22	100
66-70	7	314	221-240	24	91
71-75	7.5	293	241-260	26	84
76-80	8	275	261-280	28	78
81-85	8.5	258	281-300	30	73
86-90	9	244	301-320	32	68
91-95	9.5	231	321-340	34	64
96-100	10	220	341-360	36	61
101-110	11	200	361-380	38	57
111-120	12	183	381-420	42	52
121-130	13	169	421-460	46	47
131-140	14	157	461-500	50	44
141-150	15	146	501-550	55	40
151-160	16	137	551-600	60	36
161-170	17	129	601-650	65	33
171-180	18	122			

Virbac (Australia) Pty Limited		Page 15 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(5 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 5 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 5 L
50-55	5.5	909	181-190	19	263
56-60	6	833	191-200	20	250
61-65	6.5	769	201-220	22	227
66-70	7	714	221-240	24	208
71-75	7.5	666	241-260	26	192
76-80	8	625	261-280	28	178
81-85	8.5	588	281-300	30	166
86-90	9	555	301-320	32	156
91-95	9.5	526	321-340	34	147
96-100	10	500	341-360	36	138
101-110	11	454	361-380	38	131
111-120	12	416	381-420	42	119
121-130	13	384	421-460	46	108
131-140	14	357	461-500	50	100
141-150	15	333	501-550	55	90
151-160	16	312	551-600	60	83
161-170	17	294	601-650	65	76
171-180	18	277			

(5.5 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 5.5 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 5.5 L
50-55	5.5	1000	181-190	19	289
56-60	6	916	191-200	20	275
61-65	6.5	846	201-220	22	250
66-70	7	785	221-240	24	229
71-75	7.5	733	241-260	26	211
76-80	8	687	261-280	28	196
81-85	8.5	647	281-300	30	183
86-90	9	611	301-320	32	171
91-95	9.5	578	321-340	34	161
96-100	10	550	341-360	36	152
101-110	11	500	361-380	38	144
111-120	12	458	381-420	42	130
121-130	13	423	421-460	46	119
131-140	14	392	461-500	50	110
141-150	15	366	501-550	55	100
151-160	16	343	551-600	60	91
161-170	17	323	601-650	65	84
171-180	18	305			

Virbac (Australia) Pty Limited		Page 16 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(7.5 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 7.5 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 7.5 L
50-55	5.5	1363	181-190	19	394
56-60	6	1250	191-200	20	375
61-65	6.5	1153	201-220	22	340
66-70	7	1071	221-240	24	312
71-75	7.5	1000	241-260	26	288
76-80	8	937	261-280	28	267
81-85	8.5	882	281-300	30	250
86-90	9	833	301-320	32	234
91-95	9.5	789	321-340	34	220
96-100	10	750	341-360	36	208
101-110	11	681	361-380	38	197
111-120	12	625	381-420	42	178
121-130	13	576	421-460	46	163
131-140	14	535	461-500	50	150
141-150	15	500	501-550	55	136
151-160	16	468	551-600	60	125
161-170	17	441	601-650	65	115
171-180	18	416			

(10 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 10 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 10 L
50-55	5.5	1818	181-190	19	526
56-60	6	1666	191-200	20	500
61-65	6.5	1538	201-220	22	454
66-70	7	1428	221-240	24	416
71-75	7.5	1333	241-260	26	384
76-80	8	1250	261-280	28	357
81-85	8.5	1176	281-300	30	333
86-90	9	1111	301-320	32	312
91-95	9.5	1052	321-340	34	294
96-100	10	1000	341-360	36	277
101-110	11	909	361-380	38	263
111-120	12	833	381-420	42	238
121-130	13	769	421-460	46	217
131-140	14	714	461-500	50	200
141-150	15	666	501-550	55	181
151-160	16	625	551-600	60	166
161-170	17	588	601-650	65	153
171-180	18	555			

Virbac (Australia) Pty Limited		Page 17 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(11 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 11 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 11 L
50-55	5.5	2000	181-190	19	578
56-60	6	1833	191-200	20	550
61-65	6.5	1692	201-220	22	500
66-70	7	1571	221-240	24	458
71-75	7.5	1466	241-260	26	423
76-80	8	1375	261-280	28	392
81-85	8.5	1294	281-300	30	366
86-90	9	1222	301-320	32	343
91-95	9.5	1157	321-340	34	323
96-100	10	1100	341-360	36	305
101-110	11	1000	361-380	38	289
111-120	12	916	381-420	42	261
121-130	13	846	421-460	46	239
131-140	14	785	461-500	50	220
141-150	15	733	501-550	55	200
151-160	16	687	551-600	60	183
161-170	17	647	601-650	65	169
171-180	18	611			

(15 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 15 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 15 L
50-55	5.5	2727	181-190	19	789
56-60	6	2500	191-200	20	750
61-65	6.5	2307	201-220	22	681
66-70	7	2142	221-240	24	625
71-75	7.5	2000	241-260	26	576
76-80	8	1875	261-280	28	535
81-85	8.5	1764	281-300	30	500
86-90	9	1666	301-320	32	468
91-95	9.5	1578	321-340	34	441
96-100	10	1500	341-360	36	416
101-110	11	1363	361-380	38	394
111-120	12	1250	381-420	42	357
121-130	13	1153	421-460	46	326
131-140	14	1071	461-500	50	300
141-150	15	1000	501-550	55	272
151-160	16	937	551-600	60	250
161-170	17	882	601-650	65	230
171-180	18	833			

Virbac (Australia) Pty Limited		Page 18 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

(20 L Pack)

Liveweight (kg)	Dose Volume (mL)	Doses per pack 20 L	Liveweight (kg)	Dose Volume (mL)	Doses per pack 20 L
50-55	5.5	3636	181-190	19	1052
56-60	6	3333	191-200	20	1000
61-65	6.5	3076	201-220	22	909
66-70	7	2857	221-240	24	833
71-75	7.5	2666	241-260	26	769
76-80	8	2500	261-280	28	714
81-85	8.5	2352	281-300	30	666
86-90	9	2222	301-320	32	625
91-95	9.5	2105	321-340	34	588
96-100	10	2000	341-360	36	555
101-110	11	1818	361-380	38	526
111-120	12	1666	381-420	42	476
121-130	13	1538	421-460	46	434
131-140	14	1428	461-500	50	400
141-150	15	1333	501-550	55	363
151-160	16	1250	551-600	60	333
161-170	17	1176	601-650	65	307
171-180	18	1111			

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc.). Cattle in excess of 650 kg should be dosed at 1 mL per 10 kg liveweight. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Do not underdose. Check accuracy of dose delivery of applicator regularly.

RE-TREATMENT INTERVAL: DO NOT retreat animals with this product for 160 days after last treatment.

Rainfast

The efficacy of **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application.

Ease of Use

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE comes ready to use.

Non Irritant

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE has not been observed to cause irritation to the skin or hide damage when used as directed.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 50 days before slaughter for human consumption.

MILK: DO NOT USE on dairy cattle which are producing or may in the future produce milk for human consumption or processing.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 160 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before

Virbac (Australia) Pty Limited		Page 19 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

using this product, confirm the current ESI from either Virbac (Australia) Pty Ltd 1800 009 847 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Do not inhale. Will irritate the eyes. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves and face shield or goggles. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

Additional information is listed in the Material Safety Data Sheet

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

500 mL, 1 L Pack

DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

ALL OTHER PACK SIZES

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced but may be taken separately.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Moxidectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

STORAGE

Store below 30°C (Room temperature) and protect from light.

APVMA Approval No.: 66660/59567

BN:
EXP:



VIRBAC (AUSTRALIA) PTY. LIMITED
 361 Horsley Road, Milperra NSW 2214
 Freecall Customer Service Line: 1800 009 847
www.virbac.com.au

Virbac (Australia) Pty Limited	Page 20 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE
APVMA:	66660 / XXXXX
Version Date:	19 April 2013
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L

Text above this line or in italics between parallel lines is not included in the label copy.

[®] Registered trademark of Virbac (Australia) Pty Limited.

Virbac (Australia) Pty Limited	Page 21 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE
APVMA:	66660 / XXXXX
Version Date:	19 April 2013
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L

Text above this line or in italics between parallel lines is not included in the label copy.

OPTIONAL COMPOSITE LEAFLET For 500mL, 1L, 1.1L, 2L 2.2L and 5L packs

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CYDECTIN[®] **PLUS FLUKE POUR-ON FOR CATTLE**

For the treatment and control of moxidectin sensitive strains of gastrointestinal nematodes and triclabendazole sensitive strains of liver fluke (*Fasciola hepatica*) of cattle.

ACTIVE CONSTITUENTS: 5 g/L MOXIDECTIN
200 g/L TRICLABENDAZOLE

INTRODUCTION

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE is an endectocide with added triclabendazole with efficacy in treating and controlling internal parasites in cattle including immature and adult liver fluke.

Research on and development of **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** has produced a product with a broad spectrum of activity and wide safety margin.

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE comes ready to use as a clear solution containing 5 g/L moxidectin and 200 g/L triclabendazole. When given at a dose volume of 1 mL/10 kg liveweight, **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** delivers the recommended dosage of 0.5 mg moxidectin/kg and 20 mg triclabendazole/kg liveweight. **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** does not cause irritation to the skin or hide damage when used as directed.

INDICATIONS

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE is effective in the treatment and control of the following parasites affecting cattle:

Triclabendazole susceptible Liver fluke (late immature (6 to 8 week) and adult):
Fasciola hepatica

Moxidectin susceptible Gastrointestinal (adult and larval L4)

Small Brown Stomach Worm (*Ostertagia ostertagi*)
Stomach Hair Worm (*Trichostrongylus axei*)
Small Intestinal Worm (*Cooperia* spp.)

Moxidectin susceptible Gastrointestinal (adult)

Barber's Pole Worm (*Haemonchus placei*)
Nodule Worm (*Oesophagostomum* spp.)
Thin-necked Intestinal Worm (*Nematodirus* spp.)
Hookworm (*Bunostomum* spp.)

PROTECTION PERIOD

Virbac (Australia) Pty Limited		Page 22 of 24
Product Name:	CYDECTIN [®] PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

A single treatment at the recommended dose rate prevents re-infection with moxidectin susceptible Barber's Pole Worm (*Haemonchus contortus*) and Small Brown Stomach Worm (*Ostertagia ostertagi*) for 14 days.

DIRECTIONS FOR USE:

RESTRAINT:

DO NOT USE on dairy cattle which are producing or may in the future produce milk for human consumption or processing.

DOSE RATE

0.5 mg moxidectin and 20 mg triclabendazole/kg liveweight (1 mL per 10 kg liveweight).

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE should be applied along the top of the back from the withers to the base of the tail. The product is ready to use. Administer the product according to the dosing table below.

Dosing Table

Liveweight (kg)	Dose Volume (mL)	Doses Per Pack					
		500 mL	1 L	1.1 L	2L	2.2 L	5 L
50-56	5.5	90	181	200	363	400	909
56-60	6	83	166	183	333	366	833
61-65	6.5	76	153	169	307	338	769
66-70	7	71	142	157	285	314	714
71-75	7.5	66	133	146	266	293	666
76-80	8	62	125	137	250	275	625
81-85	8.5	58	117	129	235	258	588
86-90	9	55	111	122	222	244	555
91-95	9.5	52	105	115	210	231	526
96-100	10	50	100	110	200	220	500
101-110	11	45	90	100	181	200	454
111-120	12	41	83	91	166	183	416
121-130	13	38	76	84	153	169	384
131-140	14	35	71	78	142	157	357
141-150	15	33	66	73	133	146	333
151-160	16	31	62	68	125	137	312
161-170	17	29	58	64	117	129	294
171-180	18	27	55	61	111	122	277
181-190	19	26	52	57	105	115	263
191-200	20	25	50	55	100	110	250
201-220	22	22	45	50	90	100	227
221-240	24	20	41	45	83	91	208
241-260	26	19	38	42	76	84	192
261-280	28	17	35	39	71	78	178
281-300	30	16	33	36	66	73	166
301-320	32	15	31	34	62	68	156
321-340	34	14	29	32	58	64	147
341-360	36	13	27	30	55	61	138
361-380	38	13	26	28	52	57	131

Virbac (Australia) Pty Limited		Page 23 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

381-420	42	11	23	26	47	52	119
421-460	46	10	21	23	43	47	108
461-500	50	10	20	22	40	44	100
501-550	55	9	18	20	36	40	90
551-600	60	8	16	18	33	36	83
601-650	65	7	15	16	30	33	76

Cattle in excess of 650 kg should be dosed at 1 mL per 10 kg liveweight. A representative sample of animals should be weighed prior to treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc.). Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Do not underdose. Check accuracy of dose delivery of applicator regularly.

RE-TREATMENT INTERVAL: DO NOT retreat animals with this product for 160 days after last treatment.

Rainfast

The efficacy of **CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE** is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application.

Ease of Use

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE comes ready to use.

Non Irritant

CYDECTIN PLUS FLUKE POUR-ON FOR CATTLE has not been observed to cause irritation to the skin or hide damage when used as directed.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 50 days before slaughter for human consumption.

MILK: DO NOT USE on dairy cattle which are producing or may in the future produce milk for human consumption or processing.

TRADE ADVICE:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 160 days before

slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from either Virbac (Australia) Pty Ltd 1800 009 847 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

Harmful if inhaled or swallowed. Do not inhale. Will irritate the eyes. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow length chemical resistant gloves and face shield or goggles. If clothing becomes contaminated with product remove clothing immediately. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands thoroughly with soap and water. After each day's use wash gloves, face shield or goggles and contaminated clothing.

Additional information is listed in the Material Safety Data Sheet

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

Virbac (Australia) Pty Limited		Page 24 of 24
Product Name:	CYDECTIN® PLUS FLUKE POUR-ON FOR CATTLE	
APVMA:	66660 / XXXXX	
Version Date:	19 April 2013	
Pack sizes:	500 mL, 1 L, 1.1L, 2 L, 2.2 L, 5 L, 5.5 L, 7.5 L, 10 L, 11L, 15 L, 20 L	

Text above this line or in italics between parallel lines is not included in the label copy.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management program site. The cap should not be replaced but may be taken separately. For small containers less than or equal to 1 L dispose of empty container by wrapping with paper and putting in garbage.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Moxidectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

APVMA Approval No.: 66660/59567



VIRBAC (AUSTRALIA) PTY. LIMITED
361 Horsley Road, Milperra NSW 2214
Freecall Customer Service Line: 1800 009 847
www.virbac.com.au

® Registered trademark of Virbac (Australia) Pty Limited.