

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

SURESHOT™ GOLD

POUR-ON CATTLE WORMER

ACTIVE CONSTITUENT: 125g/L LEVAMISOLE

Also contains: 373.5 g/L ETHYLENE GLYCOL MONOBUTYL ETHER ACETATE (solvent)



20L

For the treatment of
Mature and Immature
Roundworms including
Lungworm in Cattle.

NOT TO BE USED AS AN
ORAL DRENCH OR
INJECTION.

SURESHOT GOLD

POUR-ON CATTLE WORMER

CLAIMS

SURESHOT GOLD Cattle Wormer contains levamisole, a member of the 'clear' imidazothiazole family of drenches and is effective in cattle against levamisole-sensitive strains of the following species: Mature and Immature Barber's pole worm (*Haemonchus placei*), Stomach hair worm (*Trichostrongylus axei*), Black scour worm (*Trichostrongylus* spp.), Small intestinal worm (*Cooperia* spp.), Thin necked intestinal worm (*Nematodirus* spp.), Hookworm (*Bunostomum* spp.), Strongyloides, Nodule and large bowel worms (*Oesophagostomum* spp.), Large lungworm (*Dictyoaulus* spp.), and mature (adult only) Small brown stomach worm (*Ostertagia* spp.) in cattle under 12 months of age.

DIRECTIONS FOR USE

Restraints:

DO NOT treat dogs or horses with this product.
DO NOT use this product as an oral drench or as an injection.

Precautions:

Cattle may show temporary signs of stimulation and friskiness when released from the yards soon after treatment. It is desirable to hold pregnant cows and heifers in the yards or a small paddock for 2-3 hours after treatment. Other slight side-effects of brief duration such as salivation and shivering may be seen. Avoid treatment of animals which are wet or during wet conditions. Some animals may show varying degrees of skin reactivity following treatment with this formulation. This should be considered particularly before treating show and stud animals.

Exercise care when treating weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded before drenching.

Recommended dose should not be exceeded except under veterinary supervision.

Dosage and Administration:

Dose rate 5 mL per 50 kg.

The following table may be used as a guide.

ANIMAL WEIGHT (kg)	DOSE (mL)	CRITICAL COMMENTS
50	5	Apply evenly along the backline using the special applicator. It is recommended that a broad stripe application about 75mm wide be used.
100	10	
150	15	
200	20	
250	25	DO NOT use oral drenching guns or other applicators.
300	30	NOTE: The liveweight should be estimated carefully so that the dose can be calculated accordingly. A few representative animals should be weighed or a weight band used.
350	35	
400	40	
450	45	
500	50	
550	55	
600	60	
650	65	

Cattle in excess of 650kg should be dosed at 5mL per 50kg. Concurrent use with other pour-on products requires a separate application site on the backline.

GENERAL DIRECTIONS

When *Ostertagia* are the problem in cattle over 12 months of age, a Macrocytic Lactone based product (avermectins and milbemycins) is recommended. Internal parasites may develop resistance to drenches (anthelmintics). If drench (anthelmintic) failure with this drench is suspected, seek veterinary confirmation and notify the manufacturer.

General Instructions: After use clean applicator by washing thoroughly with warm water containing detergent followed by clean cold water.

WITHOLDING PERIODS:

MEAT: DO NOT USE less than 3 days before slaughter for human consumption.

MILK: ZERO (0) days

TRADE ADVICE - EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export. THE ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from The Hunter River Company on (02) 9499 7370, or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

This product is poisonous if absorbed by skin contact or inhaled. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and face shield. If clothing becomes contaminated with product remove clothing immediately. If product gets in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor of Poisons Information Centre. Phone Australia 131126. If skin contact occurs remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL USER SAFETY INFORMATION

Additional information is listed in the Safety Data Sheet.

DISPOSAL

Triple or (preferably) pressure rinse container. Do not dispose of undiluted chemicals on site. Return clean container for recycling where this is an option or for disposal at a landfill authorised to accept that waste. If neither of these options are available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Empty containers should not be burned.

Containers should be taken after cleaning to an approved collection centre.

STORAGE

Store below 30°C (Room Temperature) in the closed, original container in a well ventilated area. Do not store for prolonged periods in direct sunlight.

APVMA Approval No.:70119/62552

Combustible liquid. Harmful if swallowed.

IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Call a POISON CENTRE if you feel unwell.

IF INHALED: Remove victim to fresh air and keep at rest in a position comfortable for breathing.

Keep away from heat, sparks, open flames and hot surfaces. - NO smoking.

Use only outdoors or in a well-ventilated area.

In case of fire, use carbon dioxide, dry chemical, foam, water fog.

Batch:

Expiry:



The Hunter River Company
7/20 St Johns Avenue,
Gordon NSW 2072
Ph: (02) 9499 7370



THE HUNTER RIVER CO.