



## APPENDIX 2

### TEMPLATE FOR RELEVANT LABEL PARTICULARS (RLPs) (Veterinary Products)

Select appropriate:

New Product (include all applicable RLPs) OR

Variation (highlight instructions that are being varied). Approval no. of label being varied:.....

Signal heading:	<b>POISON</b> <b>KEEP OUT OF REACH OF CHILDREN</b> <b>READ SAFETY DIRECTIONS BEFORE OPENING OR USING</b> <b>FOR ANIMAL TREATMENT ONLY</b>						
Product name:	<b>SURESHOT GOLD POUR-ON CATTLE WORMER</b>						
Active constituent/s:	Active constituent: 125g/L LEVAMISOLE Solvent: 373.5 g/L ETHYLENE GLYCOL MONOBUTYL ETHER ACETATE						
Statement of claims:	For the treatment of Mature and Immature Roundworms including Lungworm in Cattle						
Net contents:	1L, 2L, 5L, 20L						
Directions for Use Heading:	DIRECTIONS FOR USE						
Restrictions:	<b>DO NOT treat dogs or horses with this product.</b> <b>DO NOT use this product as an oral drench or as an injection.</b>						
Contraindications:							
Precautions:	Cattle may show temporary signs of stimulation and friskiness when released from the yards soon after treatment. It is desirable to hold pregnant cows and heifers in the yards or a small paddock for 2-3 hours after treatment. Other slight side-effects of brief duration such as salivation and shivering may be seen. Avoid treatment of animals which are wet or during wet conditions. Some animals may show varying degrees of skin reactivity following treatment with this formulation. This should be considered particularly before treating show and stud animals.  Exercise care when treating weak, pregnant and young animals to avoid unnecessary stress.  Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded before drenching.  Recommended dose should not be exceeded except under veterinary supervision.						
Side effects:							
Dosage & administration:	Dose rate 5mL per 50 Kg. The following table may be used as a guide. <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>ANIMAL WEIGHT (kg)</th> <th>DOSE (mL)</th> <th>CRITICAL COMMENTS</th> </tr> </thead> <tbody> <tr> <td>50</td> <td>5</td> <td>Apply evenly along the backline using the special</td> </tr> </tbody> </table>	ANIMAL WEIGHT (kg)	DOSE (mL)	CRITICAL COMMENTS	50	5	Apply evenly along the backline using the special
ANIMAL WEIGHT (kg)	DOSE (mL)	CRITICAL COMMENTS					
50	5	Apply evenly along the backline using the special					

	<table border="1"> <tr><td>100</td><td>10</td></tr> <tr><td>150</td><td>15</td></tr> <tr><td>200</td><td>20</td></tr> <tr><td>250</td><td>25</td></tr> <tr><td>300</td><td>30</td></tr> <tr><td>350</td><td>35</td></tr> <tr><td>400</td><td>40</td></tr> <tr><td>450</td><td>45</td></tr> <tr><td>500</td><td>50</td></tr> <tr><td>550</td><td>55</td></tr> <tr><td>600</td><td>60</td></tr> <tr><td>650</td><td>65</td></tr> </table> <p>applicator. It is recommended that a broad stripe application about 75mm wide be used.</p> <p>DO NOT use oral drenching guns or other applicators.</p> <p>NOTE: The live-weight should be estimated carefully so that the dose can be calculated accordingly.</p> <p>A few representative animals should be weighed or a weight band used.</p>	100	10	150	15	200	20	250	25	300	30	350	35	400	40	450	45	500	50	550	55	600	60	650	65
100	10																								
150	15																								
200	20																								
250	25																								
300	30																								
350	35																								
400	40																								
450	45																								
500	50																								
550	55																								
600	60																								
650	65																								
	<p>Cattle in excess of 650 kg should be dosed at 5 mL per 50 kg.</p> <p>Concurrent use with other pour-on products requires a separate application site on the backline.</p>																								
General directions:	<p><b>CLAIMS</b></p> <p>SURESHOT GOLD Cattle Wormer contains levamisole, a member of the 'clear' imidazothiazole family of drenches and is effective in cattle against levamisole-sensitive strains of the following species: Mature and Immature Barber's pole worm (<i>Haemonchus placei</i>), Stomach hair worm (<i>Trichostrongylus axei</i>), Black scour worm (<i>Trichostrongylus</i> spp.), Small intestinal worm (<i>Cooperia</i> spp.), Thin necked intestinal worm (<i>Nematodirus</i> spp.), Hookworm (<i>Bunostomum</i> spp.), Strongyloides, Nodule and large bowel worms (<i>Oesophagostomum</i> spp.), Large lungworm (<i>Dictyocaulus</i> spp.), and mature (adult only) Small brown stomach worm (<i>Ostertagia</i> spp.) in cattle under 12 months of age.</p> <p>When <i>Ostertagia</i> are the problem in cattle over 12 months of age, a Macrocytic Lactone based product (ivermectins and milbemycins) is recommended. Internal parasites may develop resistance to drenches (anthelmintics). If drench (anthelmintic) failure with this drench is suspected, seek veterinary confirmation and notify the manufacturer.</p> <p><b>General Instructions</b></p> <p>After use clean applicator by washing thoroughly with warm water containing detergent followed by clean cold water.</p>																								
Withholding Period/s:	<p><b>WITHOLDING PERIODS:</b></p> <p><b>MEAT: DO NOT USE less than 3 days before slaughter for human consumption.</b></p> <p><b>MILK: ZERO (0) days</b></p>																								
Trade Advice:	<p><b>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from The Hunter River Company on (02) 9499 7370, or the APVMA website (<a href="http://www.apvma.gov.au/residues/">www.apvma.gov.au/residues/</a>).</b></p>																								

18 Wormald Street, Symonston ACT 2609

PO Box 6182, Kingston ACT 2604 Australia

Tel: +61 2 6210 4700 Fax: +61 2 6210 4874

[www.apvma.gov.au](http://www.apvma.gov.au) ABN 19 495 043 447

Safety Directions:	This product is poisonous if absorbed by skin contact or inhaled. May irritate the eyes and skin. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist, washable hat, elbow length PVC gloves and face shield. If clothing becomes contaminated with product remove clothing immediately. If product gets in eyes wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves, face shield and contaminated clothing.
First Aid:	If poisoning occurs contact a doctor of Poisons Information Centre. Phone Australia 131 126. If skin contact occurs remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.
Additional user safety:	Additional information is listed in the Material Safety Data Sheet.
Environmental statements:	
Disposal:	[1L] Dispose of empty container by wrapping with paper and putting in garbage. [2L, 5L and 20L] Triple or (preferably) pressure rinse container. DO NOT dispose of undiluted chemicals on site. Return clean container for recycling where this is an option or for disposal at a landfill authorised to accept that waste. If neither of these options are available bury the container below 500mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, vegetation and roots. Empty containers should not be burned. Containers should be taken after cleaning to an approved collection centre.
Storage:	Store below 30°C (Room Temperature) in the closed, original container in a well ventilated area. Do not store for prolonged periods in direct sunlight.
Name & address:	The Hunter River Company Pty Limited 7/20 St Johns Avenue Gordon NSW 2072

The following is for APVMA use only:

APVMA approval no.	<b>APVMA No: 70119/62552</b>
--------------------	------------------------------