

## CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

Y·TEX®

# BRUTE®

## Pour-On for Cattle

Active Constituent:

87 g/L PERMETHRIN (25:75 cis:trans)

**FOR CONTROL OF BUFFALO FLIES\*  
AND CONTROL OF  
CATTLE LICE ON BEEF CATTLE**

\*Pyrethroid-susceptible strains

*Ready to Use*

*Rain-resistant, low-irritant formulation*

**1 litre**



**RLP  
APPROVED**

Infopest

✓  
Verified

**FILE NAME:** 1 Litre Front - 56165\_50287\_1L\_Front\_MPL\_V1.pdf

**SIZE:** 150mm x 120mm

**SCALE:** A4 100%

**DATE:** V06, 3060

## DIRECTIONS FOR USE

**Restrictions:** DO NOT USE on lactating or pregnant cows where milk or milk products may be used for human consumption. DO NOT TREAT animals with this product more than 3 times in any given 12 month period. Allow at least 21 days between treatments. More frequent use of this product is likely to result in residues above the MRL.

**Precaution:** The safety of this product in calves has not been established.

### DOSAGE AND ADMINISTRATION:

ANIMAL & TARGET PEST	DOSAGE	ADMINISTRATION
<b>Cattle, Ready-to-Use Pour-On Application:</b>  <b>Buffalo Flies</b> <i>(Haematobia irritans exigua)</i>  <b>Cattle Lice, biting and sucking species</b> <i>(Bovicola bovis, Linognathus vituli, Haematopinus eurysternus)</i>	1 mL / 30 kg body weight per animal up to 15 mL maximum. Reapply as needed but not more often than once every 21 days.	<b>SHAKE WELL BEFORE USE</b> Apply by pouring from the calibrated chamber of the dispenser; <b>For Buffalo Fly control:</b> Apply along the back-line, from the shoulders to the tail butt, when fly numbers exceed acceptable levels. <b>For Cattle Lice control:</b> Apply along the backline, from the head poll down the neck and over the shoulders. For optimum residual control, reapply after 3 weeks.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.**

### WITHOLDING PERIODS:

**MEAT:** DO NOT USE less than 7 days before slaughter for human consumption.

**MILK:** DO NOT USE on lactating or pregnant cows where milk or milk products may be used for human consumption.

### TRADE ADVICE

#### EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than 14 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the registrant on (07) 3277 7070 or the APVMA website ([www.apvma.gov.au/residues/ESI.shtml](http://www.apvma.gov.au/residues/ESI.shtml)).

**General Directions:** BRUTE Pour-on for Cattle will give effective control of Buffalo Flies and Cattle Lice when used as directed. The level and duration of control may be reduced due to high fly pressure. Treat all animals in the group.

**Application:** Shake well before using. Loosen cap and squeeze bottle to fill measuring chamber with required dosage. Remove cap and pour along the animal's backline. Apply 1 mL / 30 kg body weight up to a maximum of 15 mL per animal. The maximum number of applications is 3 per season.

**SAFETY DIRECTIONS:** May irritate the eyes. Avoid contact with eyes. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13-1126.

**Material Safety Data Sheet:** If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy, write to GPO Box 2566, Canberra, ACT 2601.

#### Protection of Wildlife, Fish, Crustaceans and the Environment:

This product is highly toxic to fish. DO NOT contaminate streams, rivers or waterways with the insecticide or used containers.

**DISPOSAL:** Dispose of empty container by wrapping with paper and putting in garbage.

**NOTICE OF WARRANTY:** Manufacturer's Warranty and Exclusion of Liability: This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the direction for use given on the label. No responsibility is accepted in respect of this product, save those non-excludable conditions implied by any Federal and State legislation or law of a territory.

**STORAGE:** Store below 30°C (room temperature). Store upright in the original container tightly closed, in a dry place. DO NOT store in direct sunlight. FLY002 4019

BATCH NUMBER

EXPIRY DATE



9 322349 003216

Distributed by: FLYCAM Pty. Ltd., A.B.N.50310625216, 7/36 Achievement Crescent, Acacia Ridge, QLD.4110 APVMA No. 56165/1L/0610

**FILE NAME:** 1 Litre Rear - 56165\_50287\_1L\_Rear\_MPL\_V2.pdf

**SIZE:** 150mm x 120mm

**SCALE:** A4 100%

**DATE:** V06, 3060

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

**Y•TEX®**

# BRUTTE®

## Pour-On for Cattle

Active Constituent: 87 g/L PERMETHRIN (25:75 cis:trans)

**FOR CONTROL OF BUFFALO FLIES\*  
AND CONTROL OF  
CATTLE LICE ON BEEF CATTLE**

\*Pyrethroid-susceptible strains



*Ready to Use*

*Rain-resistant, low-irritant formulation*

**5 litres**

FLY002\_4005

**FILE NAME:** 5 Litre Front - 56165\_50287\_5L\_Front\_MPL\_V1.pdf

**SIZE:** 172mm x 155mm

**SCALE:** A4 100%

**DATE:** V06, 3060

## DIRECTIONS FOR USE

**Restraints:** DO NOT USE on lactating or pregnant cows where milk or milk products may be used for human consumption. DO NOT TREAT animals with this product more than 3 times in any given 12 month period. Allow at least 21 days between treatments. More frequent use of this product is likely to result in residues above the MRL.

**Precaution:** The safety of this product in calves has not been established.

### DOSAGE AND ADMINISTRATION:

ANIMAL & TARGET PEST	DOSAGE	ADMINISTRATION
<b>Cattle, Ready-to-Use Pour-On Application:</b> <b>Buffalo Flies</b> <i>(Haematobia irritans exigua)</i> <b>Cattle Lice, biting and sucking species</b> <i>(Bovicola bovis, Linognathus vituli, Haematopinus eurysternus)</i>	<b>1 mL / 30 kg body weight per animal up to 15 mL maximum.</b> Reapply as needed but not more often than once every 21 days.	<b>SHAKE WELL BEFORE USE</b> Apply with the Brute applicator gun; <b>For Buffalo Fly control:</b> Apply along the back-line, from the shoulders to the tail butt, when fly numbers exceed acceptable levels. <b>For Cattle Lice control:</b> Apply along the backline, from the head poll down the neck and over the shoulders. For optimum residual control, reapply after 3 weeks.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS:

**MEAT:** DO NOT USE less than 7 days before slaughter for human consumption.

**MILK:** DO NOT USE on lactating or pregnant cows where milk or milk products may be used for human consumption.

### TRADE ADVICE

#### EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than 14 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the registrant on (07) 3277 7070 or the APVMA website ([www.apvma.gov.au/residues/ESI.shtml](http://www.apvma.gov.au/residues/ESI.shtml)).

**General Directions:** BRUTE Pour-on for Cattle will give effective control of Buffalo Flies and Cattle Lice when used as directed. The level and duration of control may be reduced due to high fly pressure. Treat all animals in the group.

**Application:** Shake well before using. Adjust the Brute Applicator gun setting to dispense the correct dosage for the animal's weight. Apply along the animal's backline while squeezing the handle of the applicator gun. Apply 1 mL / 30 kg body weight up to a maximum of 15 mL per animal. The maximum number of applications is 3 per season.

**SAFETY DIRECTIONS:** May irritate the eyes. Avoid contact with eyes. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13-1126.

**Material Safety Data Sheet:** If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy, write to GPO Box 2566, Canberra, ACT 2601.

#### Protection of Wildlife, Fish, Crustaceans and the Environment:

This product is highly toxic to fish. DO NOT contaminate streams, rivers or waterways with the insecticide or used containers.

**DISPOSAL:** Triple rinse container before disposal. Dispose of rinsate in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Do not dispose of undiluted chemicals on site. Break, crush or puncture and bury container in a local authority landfill. If not available, bury the containers below 500 mm in the disposal pit referred to above. Empty containers and product should not be burned.

**NOTICE OF WARRANTY:** Manufacturer's Warranty and Exclusion of Liability: This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the direction for use given on the label. No responsibility is accepted in respect of this product, save those non-excludable conditions implied by any Federal and State legislation or law of a territory.

**STORAGE:** Store below 30°C (room temperature). Store upright in the original container tightly closed, in a dry place. DO NOT store in direct sunlight.

FLY002 4020

BATCH NUMBER

EXPIRY DATE



Distributed by: FLYCAM Pty. Ltd., A.B.N. 50 310 625 216, 7/36 Achievement Crescent, Acacia Ridge, QLD. 4110

APVMA No. 56165/5L/0610

**FILE NAME:** 5 Litre Rear - 56165\_50287\_5L\_Rear\_MPL\_V1.pdf

**SIZE:** 172mm x 155mm

**SCALE:** A4 100%

**DATE:** V06, 3060

# CAUTION

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

**Y•TEX**<sup>®</sup>

# BRUTTE<sup>®</sup>

## Pour-On for Cattle

Active Constituent: 87 g/L PERMETHRIN (25:75 cis:trans)

**FOR CONTROL OF BUFFALO FLIES\*  
AND CONTROL OF  
CATTLE LICE ON BEEF CATTLE**

\*Pyrethroid-susceptible strains



*Ready to Use*

*Rain-resistant, low-irritant formulation*

**10 litres**

FLY002 4016

**FILE NAME:** 10 Litre Front - 56165\_50287\_10L\_Front\_MPL\_V1.pdf

**SIZE:** 172mm x 155mm

**SCALE:** A4 100%

**DATE:** V06, 3060

## DIRECTIONS FOR USE

**Restraints:** DO NOT USE on lactating or pregnant cows where milk or milk products may be used for human consumption. DO NOT TREAT animals with this product more than 3 times in any given 12 month period. Allow at least 21 days between treatments. More frequent use of this product is likely to result in residues above the MRL.

**Precaution:** The safety of this product in calves has not been established.

### DOSAGE AND ADMINISTRATION:

ANIMAL & TARGET PEST	DOSAGE	ADMINISTRATION
<b>Cattle, Ready-to-Use Pour-On Application:</b> <b>Buffalo Flies</b> <i>(Haematobia irritans exigua)</i> <b>Cattle Lice, biting and sucking species</b> <i>(Bovicola bovis, Linognathus vituli, Haematopinus eurysternus)</i>	<b>1 mL / 30 kg body weight per animal up to 15 mL maximum.</b> Reapply as needed but not more often than once every 21 days.	<b>SHAKE WELL BEFORE USE</b> Apply with the Brute applicator gun; <b>For Buffalo Fly control:</b> Apply along the back-line, from the shoulders to the tail butt, when fly numbers exceed acceptable levels. <b>For Cattle Lice control:</b> Apply along the backline, from the head poll down the neck and over the shoulders. For optimum residual control, reapply after 3 weeks.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER CONTRARY TO THIS LABEL, UNLESS AUTHORIZED UNDER APPROPRIATE LEGISLATION.**

### WITHHOLDING PERIODS:

**MEAT:** DO NOT USE less than 7 days before slaughter for human consumption.

**MILK:** DO NOT USE on lactating or pregnant cows where milk or milk products may be used for human consumption.

### TRADE ADVICE

#### EXPORT SLAUGHTER INTERVAL (ESI):

DO NOT USE less than 14 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from the registrant on (07) 3277 7070 or the APVMA website ([www.apvma.gov.au/residues/ESI.shtml](http://www.apvma.gov.au/residues/ESI.shtml)).

**General Directions:** BRUTE Pour-on for Cattle will give effective control of Buffalo Flies and Cattle Lice when used as directed. The level and duration of control may be reduced due to high fly pressure. Treat all animals in the group.

**Application:** Shake well before using. Adjust the Brute Applicator gun setting to dispense the correct dosage for the animal's weight. Apply along the animal's backline while squeezing the handle of the applicator gun. Apply 1 mL / 30 kg body weight up to a maximum of 15 mL per animal. The maximum number of applications is 3 per season.

**SAFETY DIRECTIONS:** May irritate the eyes. Avoid contact with eyes. Wash hands after use.

**FIRST AID:** If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 13-1126.

**Material Safety Data Sheet:** If additional hazard information is required, refer to the Material Safety Data Sheet. For a copy, write to GPO Box 2566, Canberra, ACT 2601.

#### Protection of Wildlife, Fish, Crustaceans and the Environment:

This product is highly toxic to fish. DO NOT contaminate streams, rivers or waterways with the insecticide or used containers.

**DISPOSAL:** Triple rinse container before disposal. Dispose of rinsate in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Do not dispose of undiluted chemicals on site. Break, crush or puncture and bury container in a local authority landfill. If not available, bury the containers below 500 mm in the disposal pit referred to above. Empty containers and product should not be burned.

**NOTICE OF WARRANTY:** Manufacturer's Warranty and Exclusion of Liability: This product as supplied is of a high grade and believed to be suitable for any purpose for which it is expressly recommended and must be used in accordance with the direction for use given on the label. No responsibility is accepted in respect of this product, save those non-excludable conditions implied by any Federal and State legislation or law of a territory.

**STORAGE:** Store below 30°C (room temperature). Store upright in the original container tightly closed, in a dry place. DO NOT store in direct sunlight.

FLY002 4017

BATCH NUMBER

EXPIRY DATE



Distributed by: FLYCAM Pty. Ltd., A.B.N. 50 310 625 216, 7/36 Achievement Crescent, Acacia Ridge, QLD. 4110

APVMA No. 56165/10L/0610

**FILE NAME:** 10 Litre Rear - 56165\_50287\_10L\_Rear\_MPL\_V1.pdf

**SIZE:** 172mm x 155mm

**SCALE:** A4 100%

**DATE:** V03, 3060