

LAB0248
IMMEDIATE CONTAINER
FRONT PANEL

Prepared: 13/12/2011

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CATTLEGUARD®

POUR-ON FOR CATTLE AND RED DEER

MOXIDECTIN 5g/L
Hydrocarbon Liquid 150g/L



For the treatment and control of moxidectin sensitive internal and external parasites of cattle and for the treatment and control of lungworm and gastrointestinal roundworms of red deer.

5 litre [10, 15, 20, 25 L]

[Pfizer logo]

BACK PANEL

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER

Comes ready to use as a clear solution containing 5g/L moxidectin. Moxidectin is a second generation macrocyclic lactone endectocide.

It is effective against parasites sensitive to this family. Resistance can occur to any anthelmintic.

(See product leaflet for additional information).

DIRECTIONS FOR USE**Dosage and Administration****Cattle and Red Deer**

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER should be applied along the top of the back from the withers to the base of the tail. Do not retreat deer within 60 days. Volume to use is 1mL per 10 kg liveweight. This will administer the correct dose of 0.5mg/kg liveweight.

CATTLEGUARD POUR-ON can be used in lactating dairy cattle, but **DO NOT TREAT LACTATING CATTLE AT LESS THAN 21 DAY INTERVALS.**

The following table may be used as a guide for the dose volume to use. [*Only relevant packsize data shown on label*].

Liveweight (kg)	Dose (mL)	Doses per Pack 5 L	Doses per Pack 10 L	Doses per Pack 15 L	Doses per Pack 20 L	Doses per Pack 25 L
31-40	4	1250	2500	3750	5000	6250
41-50	5	1000	2000	3000	4000	5000
51-60	6	833	1666	2500	3333	4166
61-70	7	714	1428	2142	2857	3571
71-80	8	625	1250	1875	2500	3125
81-90	9	555	1111	1666	2222	2777
91-100	10	500	1000	1500	2000	2500
101-125	12.5	400	800	1200	1600	2000
126-150	15	333	666	1000	1333	1666
151-175	17.5	285	571	857	1142	1428
176-200	20	250	500	750	1000	1250
201-250	25	200	400	600	800	1000
251-300	30	166	333	500	666	833
301-350	35	142	285	428	571	714
351-400	40	125	250	375	500	625
401-450	45	111	222	333	444	555
451-500	50	100	200	300	400	500
501-550	55	90	181	272	363	454
551-600	60	83	166	250	333	416
601-650	65	76	153	230	307	384

Cattle in excess of 650 kg should be dosed at 1mL per 10 kg liveweight.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc.).

LAB0248

Prepared: 13/12/2011

Do not underdose. Check accuracy of dose delivery of applicator regularly.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Broad Spectrum

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is efficacious against roundworms, lungworm, cattle tick, lice and mites which infect cattle, and lungworm and roundworms of red deer.

Persistent Protection Period

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER prevents re-infection with the following parasite species:

Parasite species	Persistent protection period
<i>Ostertagia ostertagi (O.lyrata)</i>	42 days
<i>Haemonchus placei</i>	28 days
<i>Trichostrongylus axei</i>	28 days
<i>Dictyocaulus viviparus</i>	42 days
<i>Oesophagostomum radiatum</i>	42 days
<i>Bunostomum phlebotomum</i>	42 days
<i>Nematodirus helvetianus</i>	21 days

The persistent activity of Cattleguard Pour-On for Cattle and Red Deer prevents the development of viable cattle tick (*Rhipicephalus microplus*) for 21 days after treatment.

The persistent activity of Cattleguard Pour-On for Cattle and Red Deer can be used as an aid in the control of buffalo fly (*Haematobia irritans exigua*) for up to 14 days after treatment.

Rainfast

The efficacy of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However the efficacy against tick and buffalo fly may be reduced if very heavy rain occurs following treatment. Treatment of animals under these circumstances is not good agricultural practice.

Weatherproof

The efficacy of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied to cattle under extreme seasonal conditions such as very high UV-light and temperatures in summer and very low temperatures in winter.

Safety to Target Animals

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER has a wide margin of safety when used at the recommended dose rate in all ages and breeds of cattle including pregnant cows and breeding bulls.

Concurrent Use

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is safe to use concurrently with other treatments in cattle including mineral supplements, fluke treatments and vaccines.

Do not apply at the same time as any other pour-on product.

LAB0248

Prepared: 13/12/2011

Effect on Dung Beetles

When applied as directed, the levels of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER in the faeces of treated cattle are not likely to have any significant adverse effect on the following dung beetles: *Onthophagus gazella*, *O. taurus*, *Euoniticellus intermedius* and *E.fulvus*. Effects on other dung beetle species have not been fully evaluated.

WITHHOLDING PERIOD

MEAT: CATTLE – 0 days

DEER – DO NOT USE LESS THAN 7 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

MILK: 0 days.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): 0 days

The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI with the manufacturer on 1800 814 883 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use. Do not inhale. May irritate the nose and throat.

Additional information is listed in the material safety data sheet.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

Parasite Control Programs

Effective parasite control for maximum productivity is an essential part of farm management. Follow recommended parasite management practices to reduce the potential for resistance to develop.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMuster* collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If recycling is not available, dispose of containers at a local authority landfill. If neither of these options are available, bury the containers below 500 mm in a disposal pit specifically marked and set out for this purpose, clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

LAB0248

Prepared: 13/12/2011

Made in Australia

Pfizer Animal Health

A division of Pfizer Australia Pty Ltd

38-42 Wharf Road

West Ryde. NSW 2114

©2011

EAN

APVMA Approval Number 66931/54712

STORE below 30°C (Room Temperature) and protect from light.

BATCH NO.

EXPIRY DATE

AL_____

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CATTLEGUARD®

POUR-ON FOR CATTLE AND RED DEER

MOXIDECTIN 5g/L
Hydrocarbon Liquid 150g/L

Introduction

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER is an endectocide with efficacy in treating and controlling internal and external parasites in cattle and lungworm and roundworms in red deer.

Product Description

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER comes ready to use as a clear solution containing 5 g/L moxidectin. When given at a dose volume of 1mL/10kg liveweight, CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER delivers the recommended dosage of 0.5mg moxidectin/kg liveweight. CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER does not cause irritation to the skin or hide damage when used as directed.

Spectrum of Activity

Cattle

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER is effective in the treatment and control of the following parasites affecting cattle:

Gastrointestinal (adult and larval (L4) nematodes)

Haemonchus placei

Ostertagia ostertagi (*O. lyrata*) small brown stomach worm including inhibited larvae)

Trichostrongylus axei (stomach hair worm)

Trichostrongylus colubriformis (black scour worm)

Trichostrongylus longispicularis

Cooperia oncophora (*C. surnabada*/*C. mcmasteri*)
(small intestinal worm)

Cooperia pectinata

Nematodirus helvetianus

Bunostomum phlebotomum (hookworm)

Oesophagostomum radiatum (nodule worm)

Trichuris spp

Lungworm

Dictyocaulus viviparus

LAB0248

Prepared: 13/12/2011

Cattle Tick*Rhipicephalus (Boophilus) microplus***Lice***Linognathus vituli**Haematopinus eurysternus**Solenopotes capillatus**Bovicola bovis***Mange Mites***Chorioptes bovis*

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER is effective in the treatment and aid in control of:

Buffalo Fly*Haematobia irritans exigua***Persistent activity**

In cattle, following a single application at the recommended dose rate, CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER prevents re-infection with the following parasite species:

Parasite species	Persistent protection period
<i>Ostertagia ostertagi (O.lyrata)</i>	42 days
<i>Haemonchus placei</i>	28 days
<i>Trichostrongylus axei</i>	28 days
<i>Dictyocaulus viviparus</i>	42 days
<i>Oesophagostomum radiatum</i>	42 days
<i>Bunostomum phlebotomum</i>	42 days
<i>Nematodirus helvetianus</i>	21 days

Cattle Tick (*Rhipicephalus (Boophilus) microplus*) Control

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER is effective for treatment and control of cattle tick on cattle, including strains resistant to organophosphates, synthetic pyrethroids and amidines.

The persistent activity of CATTLEGUARD® POUR-ON prevents the development of viable cattle tick (*Rhipicephalus (Boophilus) microplus*) for 21 days after treatment. For strategic programs treatment at 21 day intervals is recommended.

Some engorged females continue to drop for up to three days after the first treatment, and these may lay some viable eggs. This should be taken into account when planning a strategic dipping program.

Buffalo Fly (*Haematobia irritans exigua*) Control

The persistent activity of CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER can be used as an aid in the control of buffalo fly (*Haematobia irritans exigua*) for up to 14 days after treatment.

LAB0248

Prepared: 13/12/2011

Deer

In addition, CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER can be used to treat and control the following parasites of red deer.

Lungworm*Dictyocaulus viviparus***Gastrointestinal***Haemonchus* spp.*Ostertagia* spp*Trichostrongylus* spp*Oesophagostomum* spp**Persistent Activity against Deer Parasites**

Parasite species	Persistent protection period
<i>Ostertagia</i> spp	42 days
<i>Oesophagostomum</i> spp	42 days
<i>Dictyocaulus viviparus</i>	42 days

DIRECTIONS FOR USE**Dosage and Administration****Cattle and Red Deer**

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER should be applied along the top of the back from the withers to the base of the tail. Do not retreat deer within 60 days. Volume to use is 1mL per 10 kg liveweight. This will administer the correct dose of 0.5mg/kg liveweight.

CATTLEGUARD POUR-ON can be used in lactating dairy cattle, but **DO NOT TREAT LACTATING CATTLE AT LESS THAN 21 DAY INTERVALS.**

The following table may be used as a guide for the dose volume to use.

Liveweight	Dose	Doses per Pack	Doses per Pack	Doses per Pack	Doses per Pack	Doses per Pack
(kg)	(mL)	5 L	10 L	15 L	20 L	25 L
31-40	4	1250	2500	3750	5000	6250
41-50	5	1000	2000	3000	4000	5000
51-60	6	833	1666	2500	3333	4166
61-70	7	714	1428	2142	2857	3571
71-80	8	625	1250	1875	2500	3125
81-90	9	555	1111	1666	2222	2777
91-100	10	500	1000	1500	2000	2500
101-125	12.5	400	800	1200	1600	2000
126-150	15	333	666	1000	1333	1666
151-175	17.5	285	571	857	1142	1428
176-200	20	250	500	750	1000	1250
201-250	25	200	400	600	800	1000
251-300	30	166	333	500	666	833
301-350	35	142	285	428	571	714

LAB0248

Prepared: 13/12/2011

351-400	40	125	250	375	500	625
401-450	45	111	222	333	444	555
451-500	50	100	200	300	400	500
501-550	55	90	181	272	363	454
551-600	60	83	166	250	333	416
601-650	65	76	153	230	307	384

Cattle in excess of 650 kg should be dosed at 1mL per 10 kg liveweight.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group (bulls, cows, steers, calves, etc.).

Do not underdose. Check accuracy of dose delivery of applicator regularly.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Rainfast

The efficacy of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However the efficacy against tick and buffalo fly may be reduced if very heavy rain occurs following treatment. Treatment of animals under these circumstances is not good agricultural practice.

Weatherproof

The efficacy of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied to cattle under extreme seasonal conditions such as very high UV-light and temperatures in summer and very low temperatures in winter.

Non Irritant

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER does not cause irritation to the skin or hide damage when used as directed.

Ease of Use

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER comes ready to use. A number of applicators have been tested with CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER and several have proved suitable, including the Phillips 45mL applicator.

Broad Spectrum

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is efficacious against roundworms, lungworm, cattle tick, lice and mites which infect cattle, and lungworm and roundworms of red deer.

Safety to Target Animals

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER has a wide margin of safety when used at the recommended dose rate in all ages and breeds of cattle including pregnant cows and breeding bulls.

LAB0248

Prepared: 13/12/2011

Concurrent Use

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is safe to use concurrently with other treatments in cattle including mineral supplements, fluke treatments and vaccines.

Do not apply at the same time as any other pour-on product.

Effect on Dung Beetles

When applied as directed, the levels of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER in the faeces of treated cattle are not likely to have any significant adverse effect on the following dung beetles: *Onthophagus gazella*, *O. taurus*, *Euoniticellus intermedius* and *E.fulvus*. Effects on other dung beetle species have not been fully evaluated.

WITHHOLDING PERIOD

MEAT: CATTLE – 0 days

DEER – DO NOT USE LESS THAN 7 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

MILK: 0 days

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): 0 days

The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI with the manufacturer on 1800 814 883 or the APVMA website (www.apvma.gov.au/residues/).

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use. Do not inhale. May irritate the nose and throat.

Additional information is listed in the material safety data sheet.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone *Australia* 13 11 26.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMuster* collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If recycling is not available, dispose of containers at a local authority landfill. If neither of these options are available, bury the containers below 500 mm in a disposal pit specifically marked and set out for this purpose, clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

LAB0248

Prepared: 13/12/2011

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

Parasite Control Programs

Effective parasite control for maximum productivity is an essential part of farm management. Follow recommended parasite management practices to reduce the potential for resistance to develop.

Made in Australia

Pfizer Animal Health

A division of Pfizer Australia Pty Ltd

38-42 Wharf Road

West Ryde. NSW 2114

©2011

APVMA Approval Number 66931/54712

STORE below 30°C (Room Temperature) and protect from light.

AL_____

LAB0248
CARTON LABEL
MAIN PANEL

Prepared: 13/12/2011

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

CATTLEGUARD®

POUR-ON FOR CATTLE AND RED DEER

MOXIDECTIN 5g/L
Hydrocarbon Liquid 150g/L

For the treatment and control of moxidectin sensitive internal and external parasites of cattle and for the treatment and control of lungworm and gastrointestinal roundworms of red deer.

5 litre [10, 15, 20, 25 L]

[Pfizer logo]

LAB0248
 CARTON LABEL
 BACK PANEL

Prepared: 13/12/2011

CATTLEGUARD® POUR-ON FOR CATTLE AND RED DEER

Comes ready to use as a clear solution containing 5g/L moxidectin. Moxidectin is a second generation macrocyclic lactone endectocide.

It is effective against parasites sensitive to this family. Resistance can occur to any anthelmintic.

(See product leaflet for additional information).

DIRECTIONS FOR USE

Dosage and Administration

Cattle and Red Deer

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER should be applied along the top of the back from the withers to the base of the tail. Do not retreat deer within 60 days. Volume to use is 1mL per 10 kg liveweight. This will administer the correct dose of 0.5mg/kg liveweight.

CATTLEGUARD POUR-ON can be used in lactating dairy cattle, but **DO NOT TREAT LACTATING CATTLE AT LESS THAN 21 DAY INTERVALS.**

The following table may be used as a guide for the dose volume to use. [*Only relevant packsize data shown on label*].

Liveweight	Dose	Doses per Pack	Doses per Pack	Doses per Pack	Doses per Pack	Doses per Pack
(kg)	(mL)	5 L	10 L	15 L	20 L	25 L
31-40	4	1250	2500	3750	5000	6250
41-50	5	1000	2000	3000	4000	5000
51-60	6	833	1666	2500	3333	4166
61-70	7	714	1428	2142	2857	3571
71-80	8	625	1250	1875	2500	3125
81-90	9	555	1111	1666	2222	2777
91-100	10	500	1000	1500	2000	2500
101-125	12.5	400	800	1200	1600	2000
126-150	15	333	666	1000	1333	1666
151-175	17.5	285	571	857	1142	1428
176-200	20	250	500	750	1000	1250
201-250	25	200	400	600	800	1000
251-300	30	166	333	500	666	833
301-350	35	142	285	428	571	714
351-400	40	125	250	375	500	625
401-450	45	111	222	333	444	555
451-500	50	100	200	300	400	500
501-550	55	90	181	272	363	454
551-600	60	83	166	250	333	416
601-650	65	76	153	230	307	384

Cattle in excess of 650 kg should be dosed at 1mL per 10 kg liveweight.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest animal in each group

CARTON LABEL
BACK PANEL

(bulls, cows, steers, calves, etc.).

Do not underdose. Check accuracy of dose delivery of applicator regularly. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Broad Spectrum

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is efficacious against roundworms, lungworm, cattle tick, lice and mites which infect cattle, and lungworm and roundworms of red deer.

Persistent Protection Period

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER prevents re-infection with the following parasite species:

Parasite species	Persistent protection period
<i>Ostertagia ostertagi (O.lyrata)</i>	42 days
<i>Haemonchus placei</i>	28 days
<i>Trichostrongylus axei</i>	28 days
<i>Dictyocaulus viviparus</i>	42 days
<i>Oesophagostomum radiatum</i>	42 days
<i>Bunostomum phlebotomum</i>	42 days
<i>Nematodirus helvetianus</i>	21 days

The persistent activity of Cattleguard Pour-On for Cattle and Red Deer prevents the development of visible cattle tick (*Rhipicephalus microplus*) for 21 days after treatment.

The persistent activity of Cattleguard Pour-On for Cattle and Red Deer can be used as an aid in the control of buffalo fly (*Haematobia irritans exigua*) for up to 14 days after treatment.

Rainfast

The efficacy of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied when the hide is wet, or if rain occurs shortly after application. However the efficacy against tick and buffalo fly may be reduced if very heavy rain occurs following treatment. Treatment of animals under these circumstances is not good agricultural practice.

Weatherproof

The efficacy of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is not adversely affected if applied to cattle under extreme seasonal conditions such as very high UV-light and temperatures in summer and very low temperatures in winter.

Safety to Target Animals

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER has a wide margin of safety when used at the recommended dose rate in all ages and breeds of cattle including pregnant cows and breeding bulls.

LAB0248
CARTON LABEL
BACK PANEL

Prepared: 13/12/2011

Concurrent Use

CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER is safe to use concurrently with other treatments in cattle including mineral supplements, fluke treatments and vaccines.

Do not apply at the same time as any other pour-on product.

Effect on Dung Beetles

When applied as directed, the levels of CATTLEGUARD POUR-ON FOR CATTLE AND RED DEER in the faeces of treated cattle are not likely to have any significant adverse effect on the following dung beetles: *Onthophagus gazella*, *O. taurus*, *Euoniticellus intermedius* and *E. fulvus*. Effects on other dung beetle species have not been fully evaluated.

WITHHOLDING PERIOD

MEAT: CATTLE – 0 days

DEER – DO NOT USE LESS THAN 7 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.

MILK: 0 days.

EXPORT SLAUGHTER INTERVAL (ESI): 0 days

The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI with the manufacturer on 1800 814 883 or the APVMA website (www.apvma.gov.au/residues/).

LAB0248
CARTON LABEL
LEFT SIDE PANEL

Prepared: 13/12/2011

SAFETY DIRECTIONS

Poisonous if swallowed. Will irritate eyes and skin. Avoid contact with eyes and skin. Wash hands after use. Do not inhale. May irritate the nose and throat.

Additional information is listed in the material safety data sheet.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 13 11 26.*

Parasite Control Programs

Effective parasite control for maximum productivity is an essential part of farm management. Follow recommended parasite management practices to reduce the potential for resistance to develop.

DISPOSAL

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to state legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMuster* collection or similar container management program site. The cap should not be replaced, but may be taken separately.

If recycling is not available, dispose of containers at a local authority landfill. If neither of these options are available, bury the containers below 500 mm in a disposal pit specifically marked and set out for this purpose, clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT

Extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

LAB0248
CARTON LABEL
RIGHT SIDE PANEL

Prepared: 13/12/2011

CATTLEGUARD®

Made in Australia
Pfizer Animal Health
A division of Pfizer Australia Pty Ltd
38-42 Wharf Road
West Ryde. NSW 2114
©2011

EAN

APVMA Approval Number 66931/54712

STORE below 30°C (Room Temperature) and protect from light.

BATCH NO.

EXPIRY DATE

AL_____

LAB0248
CARTON LABEL
LID

Prepared: 13/12/2011

CATTLEGUARD®