

**CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY**



BAYCOX[®]



COCCIDIOCIDE SOLUTION

Active Constituent: 25 g/L TOLTRAZURIL

For the treatment and control of coccidiosis in chickens caused by Eimeria species.

100mL

[chicken pictogram]

For chickens

Baycox Coccidiocide Solution

Baycox is added to the drinking water of chickens for the treatment and control of coccidiosis caused by *Eimeria* species including: *Eimeria tenella*, *Eimeria necatrix*, *Eimeria acervulina* and *Eimeria maxima*.

Restraint

DO NOT treat replacement pullets more than twice.

Directions for Use

Baycox is diluted at the rate of 3 litres per 1000 litres of drinking water (3:1000). It is administered on 2 consecutive days with an 8 hour treatment period on both days. Ensure full lighting conditions during treatment. Do not withhold feed during treatment. Add the required amount of Baycox to the amount of water consumed in an 8 hour period and stir thoroughly. Allow for varying water consumption during hot weather. Treat the birds for 8 hours each day for 2 days (2x8 hour treatments). No other source of drinking water should be available to birds during the 8 hour treatment period.

WITHHOLDING PERIOD

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

EGGS: DO NOT USE in laying hens. DO NOT USE in replacement pullets within 8 weeks of laying onset, where eggs are to be used or processed for human consumption. Eggs laid during the withholding period MUST NOT BE USED or processed for human consumption.

General Instructions

DO NOT USE GALVANISED OR RUSTY IRON WATER TANKS AND RETICULATION SYSTEMS.

Baycox mixes with all types of water. However in areas where the drinking water is acidic, the water should be tested with pH test strips and brought to a minimum pH of 8.5 using sodium carbonate (washing soda).

To avoid excessive foaming, add Baycox to the water after filling the supply tank.

Broiler Treatment Programme: To use Baycox in a clinical outbreak obtain an accurate diagnosis. Begin treatment with Baycox as soon as coccidiosis is confirmed. A treatment period of 2 days is sufficient. If required, treatment may be repeated after 5 days.

Broiler Preventive Programme: To use Baycox as a coccidiosis preventive in broiler production, give Baycox as above on days 9 and 10, 16 and 17, and 23 and 24 of life. Do not use a coccidiostat in the feed. An additional treatment may be necessary after 35 days of age but not within 14 days of slaughter.

Layer Birds: Baycox may be used in replacement pullets and replacement layer breeder birds via the drinking water for an 8 hour period each day for two consecutive days. Treatment may be repeated after 5 days if required.

Protection of wildlife, fish, crustaceans and the environment

Do not discharge waste water containing Baycox into dams, streams or watercourses.

Disposal and Protection of Others

Dispose of empty container by wrapping with paper and putting in garbage.

This container can be recycled if it is clean, dry, free of visible residues and has the **drumMUSTER** logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State Legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any **drumMUSTER** collection or similar container management program site. The cap should not be replaced but may be taken separately.

Safety Directions

Harmful if swallowed. Will irritate the eyes. Avoid contact with eyes.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Additional information is listed in the Material Safety Data Sheet.

Limitation of Liability

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

BAYCOX® is a registered trademark of Bayer AG, Leverkusen, Germany.
© Bayer Australia Ltd 2010

Manufactured in Germany

APVMA Approval No 45626/56401

Storage

Store below 30°C (room temperature) in the closed, original container, as cool as possible.

Batch: Expiry:

Bayer

875 Pacific Highway
PYMBLE NSW 2073
ACN 000 138 714

APN Barcode

Back panel

Issue Date: 4/2/2013

Page: 4 of 4

The information above this line is not included in the label

CUSTOMER INFORMATION LINE

1800 678 368 TOLL FREE from anywhere in Australia

9.00 am to 4.00 pm (Eastern Time) Monday to Friday

www.bayeranimal.com.au