

Version 1.3 28/07/08
Size: 120 x 175 mm

MPL_Minbal 4 in 1_350mL

FOR ANIMAL TREATMENT ONLY

MINBAL 4 IN 1
BALANCED MINERAL INJECTION

ACTIVE CONSTITUENTS:	g/350mL	g/L
CALCIUM (as Ca Gluconate)	9.6	27.5*
MAGNESIUM (as Magnesium Hypophosphite)	1.5	4.7
PHOSPHORUS (as Hypophosphite)	4.3	12.2
ALSO CONTAINS GLUCOSE	63.7	182.0
*Equivalent to 330g/L Calcium borogluconate		

For the treatment of Milk fever, Grass Tetany and associated metabolic conditions in Cattle & Sheep.

Contents: 350mL

Directions for Use:

Use of this product intravenously may be hazardous to the animal and advice from a registered veterinary surgeon should be sought

Caution: Avoid carcass damage.

1. Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. This product should be injected only under the skin.

When treating hypomagnesaemia (grass/lactating tetany) an initial treatment with a magnesium salt solution should be given as well as this product.

DOSAGE: (By subcutaneous injections)

Cattle: Small cows: 350mL
Heavier Cows: 350mL - 700mL
Sheep: Small breeds 40-70kGS: 60-100mL
Heavier breeds 70-100Kgs: 100-150mL

Discard un-used portion within 24 hours of first broaching pack.

If possible warm to body temperature prior to injection. Snap tip off plug, ensuring the solution can flow freely and connect the sterile needle provided. Disinfect the injection site and insert the needle under the skin of the neck or over the ribs. Gently squeeze the bag to inject the contents. Large volumes should be distributed over more than one injection site. If complete recovery does not occur within 3-4 hours repeat injection and seek veterinary advice.

WITHHOLDING PERIODS: NIL

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): 0 days

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 13 11 26.

Disposal: Dispose of unwanted packaging by wrapping in paper and placing in garbage. Discarded needles should be placed in a designated and appropriately labelled sharps container.

Storage: Store below 30°C (Room Temperature). Protect from light.

Rudduck Australia
A division of Sykes Vet International Pty Ltd
27 Healey Road
Dandenong VIC 3174



APVMA: 38151/350ML/1007
Batch
Expiry:

Version 1.3 28/07/08
Size: 115 x 235 mm

MPL_Minbal 4 in 1_500mL

FOR ANIMAL TREATMENT ONLY

MINBAL 4 IN 1
BALANCED MINERAL INJECTION

ACTIVE CONSTITUENTS:	g/500mL	g/L
CALCIUM (as Ca Gluconate)	13.8	27.5*
MAGNESIUM (as Magnesium Hypophosphite)	2.4	4.7
PHOSPHORUS (as Hypophosphite)	6.1	12.2
ALSO CONTAINS GLUCOSE	90.8	182.0
*Equivalent to 330g/L Calcium borogluconate		

For the treatment of Milk fever, Grass Tetany and associated metabolic conditions in Cattle & Sheep.

Contents: 500mL

Directions for Use:

Use of this product intravenously may be hazardous to the animal and advice from a registered veterinary surgeon should be sought

Caution: Avoid carcass damage.

1. Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. This product should be injected only under the skin.

When treating hypomagnesaemia (grass/lactating tetany) an initial treatment with a magnesium salt solution should be given as well as this product.

DOSAGE: (By subcutaneous injections)
Cattle: Small cows: 350mL
Heavier Cows: 350mL - 700mL
Sheep: Small breeds: 40-70kGS: 60-100mL
Heavier breeds 70-100Kgs: 100-150mL

Discard un-used portion within 24 hours of first broaching pack.

If possible warm to body temperature prior to injection. Snap tip off plug, ensuring the solution can flow freely and connect the sterile needle provided. Disinfect the injection site and insert the needle under the skin of the neck or over the ribs. Gently squeeze the bag to inject the contents. Large volumes should be distributed over more than one injection site. If complete recovery does not occur within 3-4 hours repeat injection and seek veterinary advice.

WITHHOLDING PERIODS: NIL

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): 0 days

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone Australia 13 11 26.

Disposal: Dispose of unwanted packaging by wrapping in paper and placing in garbage. Discarded needles should be placed in a designated and appropriately labelled sharps container.

Storage: Store below 30°C (Room Temperature). Protect from light.

Rudduck Australia
A division of Sykes Vet International Pty Ltd
27 Healey Road
Dandenong VIC 3174

APVMA: 38151/500ML/1007
Batch
Expiry:

