

FOR ANIMAL TREATMENT ONLY



CETRIGEN[®]

Antibacterial Wound Aerosol

ACTIVE CONSTITUENT: Each 100 g contains:

0.1 g CETRIMIDE

4 g DIETHYL TOLUAMIDE

3 g DI-N-PROPYL ISOCINCHOMERONATE

With purple dye and Insect Repellents
**An antibacterial aerosol for use on open wounds in cattle, sheep,
pigs, horses, poultry and all other animals**

100 g

VIRBAC (AUSTRALIA) PTY. LIMITED
15 Pritchard Place, Peakhurst NSW 2210

DIRECTIONS FOR USE:

Spray lightly onto affected area, holding can 15 - 20 cm away. Avoid excessive run-off from wound site. Keep away from eyes and mucous membranes.

WITHHOLDING PERIOD:

MEAT: DO NOT USE less than 28 days before slaughter for human consumption.

MILK: DO NOT USE on female cattle or sheep which are producing, or may in the future produce, milk for human consumption.

EGGS: DO NOT USE on laying birds where eggs or egg products are to be used for human consumption.

WARNING

THIS PRODUCT CONTAINS DIETHYLTOLUAMIDE, WHICH MAY BE DANGEROUS WHEN USED IN LARGE AMOUNTS OR FOR A LONG PERIOD. INTENTIONAL MISUSE BY DELIBERATELY CONCENTRATING AND INHALING CONTENTS CAN BE HARMFUL OR FATAL.

PRESSURISED DISPENSER. PROTECT FROM SUNLIGHT AND DO NOT EXPOSE TO TEMPERATURES EXCEEDING 50C. DO NOT PIERCE OR BURN, EVEN AFTER USE.

EXTREMELY FLAMABLE.

DO NOT SPRAY ON A NAKED FLAME OR ANY INCANDESCENT MATERIAL.

KEEP AWAY FROM SOURCES OF IGNITION – NO SMOKING.

KEEP OUT OF REACH OF CHILDREN.

DISPOSAL:

Aerosols should be completely emptied before disposal. Place empty used can in household rubbish.

STORAGE:

–

Store below 30°C (Room Temperature).

® Registered Trade Mark of Virbac.

[B]

CODE:

EXP.

NRA: 38731 / 0902