

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY



BAYTICOL®

CATTLE DIP AND SPRAY

Active Constituent: 75g/L FLUMETHRIN
Solvent: 745g/L LIQUID HYDROCARBON

For the control of synthetic pyrethroid susceptible strains of
cattle tick (*Boophilus microplus*) including chlorinated
hydrocarbon-organophosphorus-amidine-resistant strains,
Paralysis Tick (*Ixodes holocyclus*) and New Zealand Cattle Tick
(*Haemaphysalis longicornis*) on cattle, horses;
and Brown Dog Tick (*Rhipicephalus sanguineus*), New Zealand Cattle Tick
(*Haemaphysalis longicornis*) ticks on dogs.

250 mL
[500 mL]
[1 Litre]
[5 Litres]

DrumMuster Logo (on 5 litre pack)

BAYTICOL® CATTLE DIP AND SPRAY

DIRECTIONS FOR USE (In QLD, NSW, WA, NT only)

Precautions and Contraindications

Sick or exhausted cattle, horses and dogs should not be dipped.

Exercise normal care when handling stock. Dip young animals separately. Individual horses may be sensitive to pyrethroids - hose down thoroughly if reaction is observed.

PEST	METHOD	RATE*		CRITICAL COMMENTS
Cattle tick (<i>Boophilus microplus</i>) Paralysis tick (<i>Ixodes holocyclus</i>)	Plunge Dip	Initial charge 1:1000 (1 litre per 1000 litres of water) = 75mg flumethrin/litre of wash	Replenishment 1:1000 (1 litre per 1000 litres of ADDED water)	Do not allow dip level to fall by more than one quarter of its capacity before replenishment.
	New Zealand Cattle Tick (<i>Haemaphysalis longicornis</i>) on cattle and horses	Recirculating Spray-race	Initial charge 1:1000 (1 litre per 1000 litres of water) = 75mg flumethrin/litre of wash	Replenishment Method A 1:1000 (250ml per 250 litres of ADDED water)
			Method B 1:1000 (1 litre per 1000 litres of ADDED water)	Replenish after initial 1000 litre fall in sump volume; then after 500 litre fall; subsequently after every 250 litre fall. No more than 750 adults should be treated before the sump is emptied and recharged.
	Constant replenishment Spray-race	Initial Charge 1:1000 (1 litre per 1000 litres of water) = 75mg flumethrin/litre of wash.	Replenishment 1:1000 (1 litre per 1000 litres of replenishment water)	No more than 750 adults should be treated before the sump is emptied and recharged.
	Hand sprays or Non-recirculation Spray-races	Initial Charge 1:1000 (1 litre per 1000 litres of water) = 75mg flumethrin/litre of wash	Replenishment Not applicable	Use a minimum of 8-10 litres of wash per adult to ensure saturation.

<p>Brown Dog Tick (<i>Rhipicephalus sanguineus</i>) and New Zealand Cattle Tick (<i>Haemaphysalis longicornis</i>) on dogs</p> <p>This product does not control Paralysis Ticks (<i>Ixodes holocyclus</i>) on dogs</p>	<p>Rinse</p>	<p>Dilution rate: 3 ml Bayticol per 5 litres of water</p>	<p>Replenishment Not applicable</p>	<p>The dog must be thoroughly wet all over with the rinse. Do this by using a sponge to rub the wash into all parts of the coat. Do not wash out. Allow to dry in the coat. Repeat fortnightly for continuous control. Ticks commonly attach to the skin around the head and neck and between the toes. Pay particular attention to ensure that these areas are thoroughly treated. When tick pressure is particularly heavy owners are advised to visually inspect the animal for ticks at frequent intervals to ensure application has been thorough. This product does not control Paralysis Ticks. It is recommended that in paralysis tick areas dogs are thoroughly searched each day for attached ticks and that any ticks found are removed.</p>
--	--------------	--	--	---

* Reinforcement of dips is not generally necessary. However, if through flooding or neglect, the active constituent concentration is found to be below 60 mg per litre, reinforce the dip back to the charging rate (75mg per litre).

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

GENERAL INSTRUCTIONS (Cattle, Horses)

Bayticol is effective against sensitive strains of cattle tick but resistance may develop to any tickicide. If lowered level of control is experienced, resistance testing of your ticks is recommended.

Bayticol should be used at intervals of not more than 19-21 days to control cattle tick (*Boophilus microplus*).

Control is not adversely affected by rainfall a few hours after treatment.

Bayticol controls Paralysis Tick (*Ixodes holocyclus*) and New Zealand cattle tick (*Haemaphysalis longicornis*) on cattle and horses. To control these ticks when active the treatment interval should be 10 days.

Bayticol is compatible with externally applied Buffalo Fly Sprays plus vaccination and normal management procedures such as castration and dehorning.

Bayticol may be used in water of any hardness. Softening agents should not be used.

DIP AND SPRAY MANAGEMENT (Cattle, Horses)

1. **VOLUME.** The volume of the dip or sump should be accurately measured. Relevant volume levels should be marked on the dip wall or a dip stick. Sumps should be maintained in near-full condition. For efficient tick control, replenish before volume falls below recommended levels. (See Table).

2. **MIXING.** Add required concentrate to twice its volume of water in a bucket, mix, then pour along the length of the dip, or across the surface of the sump. Stir thoroughly. In a plunge dip, use 20-25 cattle as stirrers. These must be dipped again. A spray-race sump should be stirred with a paddle and agitated vigorously with the pump. It is essential that the dip be stirred at the commencement of each day's dipping and after every replenishment.

3. **SAMPLING.** Stir dip or sump before sampling. In a plunge dip the sample should be taken from 1 metre below the surface at the jump end as the last animal is passing through the dip. In a spray-race, on completion allow drainings to return to sump, stir with paddle and pump and sample from below the surface immediately. Sample analysis is available through Bayer Australia Limited, P.O. Box 377, Cannon Hill, QLD 4170 Contact the company for suitable sample bottles.

WITHHOLDING PERIOD:**CATTLE – MEAT: NIL****MILK: NIL****HORSES - DO NOT USE less than 1 day before slaughter for human consumption.****TRADE ADVICE****EXPORT SLAUGHTER INTERVAL (ESI): 0 (zero) days****SAFETY DIRECTIONS**

Will damage the eyes. Will irritate the nose, throat and skin. Avoid contact with the skin and eyes. Do not inhale vapour. If product on skin, immediately wash area with soap and water. If product in eyes wash it out immediately with water. Wash hands after use. When opening the container and preparing the spray/dip, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length nitrile gloves, and goggles. When using the prepared spray/dip, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). After each day's use wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126.

If swallowed do NOT induce vomiting.

Additional information is in the Material Safety Data Sheet.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND ENVIRONMENT.

Very toxic to aquatic life. Dangerous to fish and aquatic organisms.. Do not contaminate dams, streams or waterways with product or used containers. Store in original container, tightly closed in a safe place under lock and key.

ADDITIONAL SAFETY INFORMATION

Avoid ingestion. Do not apply the product against the wind. Do not apply the product in the presence of unprotected persons. Avoid contact with mouth and clothing. Do not eat or drink while using the product. Do not allow children to contact freshly treated dogs while they are wet.

Back Panel (cont'd)	Issue Date: 20/2/2012	Page: 5 of 5
<i>The information above this line is not included in the label</i>		

DISPOSAL OF USED DIPWASH

Disposal of unwanted dipwash should be into a pit of adequate size on level ground and away from dams, streams and watercourses. Degradation of the active chemical in the dipwash may be accelerated by adding slaked lime (calcium hydroxide) at the rate of 4 kg per 1000 litres of wash.

Disposal of containers:

Triple or (preferably) pressure rinse empty containers, adding rinsate to the dip. DO NOT dispose of undiluted chemicals on site. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. DO NOT burn empty containers or product.

Limitation of Liability

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

TOXIC 6 diamond	PESTICIDE, LIQUID, TOXIC (Flumethrin)		
	UN No. 2902	PG III	Hazchem 2X
	Bayer Australia Limited emergency contact 1 800 033 111 Australia wide, 24 hours		

<p>CUSTOMER INFORMATION LINE 1800 678 368 TOLL FREE from anywhere in Australia 9.00 am to 4.00 pm (Eastern Time) Monday to Friday www.bayeranimal.com.au</p>

Bayer

875 Pacific Highway
PYMBLE NSW 2073
ACN 000 138 714

BAYTICOL® is a trademark of Bayer AG, Leverkusen, Germany.
MANUFACTURED IN GERMANY

Store below 30°C (Room Temperature).
Store in the closed, original container, as cool as possible. Store away from food and feed.

APVMA Approval No 41037/55190
See container for batch number and expiry date