



Product Name: TRI-SOLFEN TOPICAL ANAESTHETIC & ANTISEPTIC SOLUTION FOR PAIN RELIEF IN LAMBS AND CALVES
APVMA Approval No: 60099/117753



Label Name:	TRI-SOLFEN TOPICAL ANAESTHETIC & ANTISEPTIC SOLUTION FOR PAIN RELIEF IN LAMBS AND CALVES
-------------	--

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	40.6 g/L LIGNOCAINE (as hydrochloride) 4.2 g/L BUPIVACAINE (as hydrochloride) 24.8 mg/L ADRENALINE (as acid tartrate) 5.0 g/L CETRIMIDE
-------------------------	--

Claims:	TRI-SOLFEN is a local anaesthetic and antiseptic gel spray for pain relief: <ul style="list-style-type: none">• on lambs following Mulesing and tail docking,• on lambs and calves during and following castration, and• on calves after disbudding or dehorning.
---------	---

Net Contents:	10 L 15 L 1 L 2.5 L 20 L 22 L 5 L
---------------	---

Directions for Use:	
---------------------	--

Restrains:	DO NOT USE on lactating or pregnant cows and ewes where milk may be used or processed for human consumption.
------------	--

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	<p>Mulesing Total Dose Lambs (5-10 kg) 6 mL Lambs (11-15 kg) 8 mL Lambs (16-20 kg) 10 mL Lambs (over 20kg) 12 mL Tail docking (if not concurrently mulesed) Total Dose Lambs up to 10 kg 1.5 mL Lambs over 10 kg 2 mL Apply TRI-SOLFEN spray to the wound immediately after mulesing or tail docking. Ensure that entire wound area is evenly covered with spray, particularly wound edges. Do not spray to runoff.</p> <p>Castration Total Dose Lambs up to 10 kg 3 mL Lambs over 10 kg 4.5 mL Calves 30-100 kg 6 mL Calves over 100 kg 9 mL Refer to table for total TRI-SOLFEN dose per animal.</p> <ol style="list-style-type: none"> 1. Cut through skin of scrotum as per normal procedure and gently extrude the testis to expose the spermatic cord (without cutting or tearing it) 2. Insert nozzle through skin incision (prior to removal of testicle) and carefully advance applicator along the length of exposed cord as far as possible within the scrotum. 3. Eject one third (1/3) of total TRI-SOLFEN dose as the applicator is withdrawn from the scrotum to coat the spermatic cord. 4. Insert nozzle through the other scrotal incision and apply one third (1/3) of total TRI-SOLFEN dose to the spermatic cord (as described above) 5. After application of TRI-SOLFEN to the spermatic cords, the testicles can be extracted from the scrotum and removed as per normal procedure 6. Lastly, apply the final one third (1/3) of total TRI-SOLFEN dose to the edges of the scrotal skin wound(s). <p>Disbudding/Dehorning Method Dose per horn Hot Iron (Cautery) Disbudding 4 mL Calves < 3 months of age Scoop, cup or knife (excision) disbudding or dehorning 4 mL Calves</p> <ol style="list-style-type: none"> 1. Perform disbudding or dehorning for each horn according to acceptable procedures. 2. Immediately after disbudding or dehorning the first horn, apply TRI-SOLFEN with the recommended applicator to completely cover the wound including the surrounding burn wound (if using cautery) or the surrounding cut skin edge (if using excision). 3. Repeat step 2 for the second horn, also ensuring the wound and surrounding affected tissues are completely covered. 4. Do not spray to run-off. 5. Take care to prevent TRI-SOLFEN getting into the eyes of treated calves. Use of the special TRI-SOLFEN applicator is recommended.
----------------------------	--

General Directions:	In lambs, insecticidal preparations can be applied to the wound after application of TRISOLFEN to prevent fly strike.
---------------------	---

Withholding Periods:	<p>MEAT (CATTLE and SHEEP): Treated sheep and cattle must not be supplied or sold for slaughter for human consumption for at least 90 days following treatment with this product.</p> <p>MILK: DO NOT USE on lactating or pregnant cows and ewes where milk may be used or processed for human consumption.</p>
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 90 days before slaughter for export. Before using this product, confirm the current ESI from Bayer Australia Ltd (Animal Health) on 1800 678 368 before using the product or the APVMA website (www.apvma.gov.au/residues)
Safety Directions:	May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and using the product, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves. Wash hands after use. After each day's use, wash gloves and contaminated clothing. Do not rehandle treated animals until after the product has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length PVC gloves.
First Aid Instructions:	First Aid Instructions: If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126.
First Aid Warnings:	
Additional User Safety:	
Environmental Statements:	
Disposal:	Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted chemical according to State legislative requirements. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30°C (room temperature). Do not store at a low temperature. Protect from light.