

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

FOR SPECIALIST ADVICE IN AN EMERGENCY DIAL

1800 033 111

24 HOURS AUSTRALIA WIDE

CONQUEST
SNOKER

240 EC SELECTIVE HERBICIDE

ACTIVE CONSTITUENT: 240 g/L CLODINAFOP-PROPARGYL
60 g/L CLOQUINTOCET-MEXYL
SOLVENTS: 553 g/L LIQUID HYDROCARBONS
100 g/L N-METHYL-2-PYRROLIDONE

GROUP A HERBICIDE

Controls Wild Oats, Paradoxa Grass (Annual Phalaris), Canary Grass and Annual Ryegrass in Wheat. Highly active on Wild oat and can be mixed with many broadleaf herbicides

IMPORTANT: READ THIS BOOKLET THOROUGHLY BEFORE USING THIS PRODUCT.

 **CONQUEST**
CROP PROTECTION

 **CONQUEST**
CROP PROTECTION

CONQUEST CROP PROTECTION PTY. LTD
ABN 84 098 814 932
Level 1, 4 Collingwood St, Osborne Park, WA 6017
Telephone: 08 9347 0500, Fax: 08 9347 0551

Conquest Crop Protection Pty Ltd.

ABN 84 098 814 932

Level 1, 4 Collingwood Street
Osborne Park, WA 6017

Telephone (08) 9347 0500

Facsimile: (08) 9347 0551

APVMA Approval No: 62567/1007

DIRECTIONS FOR USE**RESTRAINTS:**

DO NOT apply if rain is likely to occur within two hours.

DO NOT apply to weeds or crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure.

DO NOT apply to weeds or crops immediately after a prolonged period of frosts.

DO NOT apply under poor growing conditions.

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE	WEED GROWTH STAGE	CRITICAL COMMENTS
Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Wild Oats	QLD, Nthn NSW only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 to 5 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT THESE RATES Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 leaf to mid tillering	
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 to 5 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing Conquest Snooker with certain broadleaf weed herbicides some reduction in grass weed control may occur.
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water.	2 leaf to mid tillering	Unless otherwise specified always use D-C Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 125 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
		Sthn NSW, Vic, SA, WA only	65 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.	2 to 5 leaf	MAY BE TANK MIXED WITH COMPATIBLE HERBICIDES Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing Conquest Snooker with certain broadleaf weed herbicides some reduction in grass weed control may occur. Refer to compatibility chart for specific rates for tank mixes. Unless otherwise specified always use D-C Trate, Uptake or Hasten when tank mixing broadleaf herbicides. Apply at 85 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			85 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake oil per 100 L of water. Or 100 mL of a 100% non-ionic surfactant per 100 L of water.	2 leaf to mid tillering	

CROP	WEEDS CONTROLLED	STATE	RATE PER HECTARE	WEED GROWTH STAGE	CRITICAL COMMENTS
continued Wheat 2 leaf to late jointing (Zadocks 12 – 37)	Paradoxa Grass (<i>Phalaris paradoxa</i>) Canary Grass (<i>Phalaris minor</i>)	Qld, NSW, Vic, SA only	85 mL + 1 L of D-C Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 3 leaf	NOT TO BE MIXED WITH BROADLEAF HERBICIDES AT BELOW 160 mL/ha Apply in mixtures for broadleaf weed control only when weeds are actively growing. When mixing Conquest Snooker with certain broadleaf weed herbicides some reduction in grass weed control may occur. Apply at 160 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.
			125 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 5 leaf	
			160 mL+ 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 leaf to early tillering	
	Annual Ryegrass	NSW, Vic, SA, WA only	160 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.	2 to 4 leaf	
	210 mL + 1 L of D-C-Trate oil or 500 mL Hasten or Uptake per 100 L of water.		2 leaf to early tillering	Apply at 210 mL/ha when weeds are not actively growing. Where low water volumes are used, DO NOT use less than the minimum oil volumes specified (See Minimum Oil Volumes). Cold water: Under cold water conditions (10°C or less), use Hasten instead of D-C-Trate or Uptake.	

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD

HARVEST GRAIN:

ANIMAL GRAZING/FEDING:

WITHHOLDING PERIOD NOT REQUIRED WHEN USED AS DIRECTED.

DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER APPLICATION.

GENERAL INSTRUCTIONS

Conquest Snooker 240 EC Selective Herbicide should be applied only when weeds are actively growing.

RESISTANT WEEDS WARNING

GROUP A HERBICIDE

Conquest Snooker 240 EC Selective Herbicide is a member of the aryloxyphenoxypropionate ("fop") group of herbicides. Conquest Snooker has the inhibition of fat (lipid) synthesis, (or inhibitors of acetyl CoA carboxylase) mode of action. For weed resistance management Conquest Snooker is a Group A herbicide.

Some naturally-occurring weed biotypes resistant to Conquest Snooker and other Group A herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Conquest Snooker or other Group A herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Conquest Agrochemicals accepts no liability for any losses that may result from the failure of Conquest Snooker to control resistant weeds.

Weed biotypes resistant to diclofop-methyl and other "grass specific" herbicides can also often be resistant to Conquest Snooker.

Before using Conquest Snooker on a population resistant to or suspected to be resistant to "grass specific" herbicides, have a resistance test conducted to ensure that it is susceptible to Conquest Snooker.

Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be made to prevent seed set of the surviving weeds.

DO NOT make more than one application of a Group A herbicide with the inhibition of fat (lipid) synthesis (or inhibitors of acetyl CoA carboxylase) mode of action to a crop in the same season.

If the user suspects that the target weed population is resistant to herbicides with this mode of action, Conquest Snooker or other Group A herbicides should not be used.

Strategies to minimise the risk of herbicide resistance are available. The above recommendations should be incorporated into an Integrated Weed Management (IWM) Program. Consult your farm chemical supplier, consultant, local Department of Agriculture or Primary Industries, or local Conquest Agrochemical representative for details.

MIXING

Fill the spray tank to one quarter full. Then add Conquest Snooker while adding water to make up the final spray volume. Pour Conquest Snooker into the stream of incoming water. Good agitation at the time Conquest Snooker is added to the tank will ensure good mixing. Add the required amount of D-C-Trate*, Hasten* or Uptake* oil just before the tank is full of water and with the agitators in motion.

Mixing order

Some products may react with other products if they are not mixed in the correct order. The general mixing order of products should be -

1. Water conditioners or buffers
2. Water dispersable granules (WG)
3. Wettable powders (WP)

4. Flowable or suspension concentrates (SC)
5. Emulsifiable concentrates (EC) (Conquest Snooker)
6. Water based or soluble concentrates
7. Oils and wetters - eg Hasten, BS 1000

It is important to ensure that each individual component of the tank mix is fully dissolved and in solution before the next product is added to the tank mix, otherwise mixing problems may occur.

Minimum Oil Volumes

The minimum amount of oil to be applied with Conquest Snooker will depend on the type of oil used. When applying Conquest Snooker using low water volumes, DO NOT use less than the minimum oil volumes specified below:

- For oils normally applied at 1% v/v (eg D-C-Trate), DO NOT apply less than 500 mL oil/ha.
- For oils normally applied at 0.5% v/v (eg Uptake, Hasten), DO NOT apply less than 250 mL oil/ha.

Cleaning Spray Equipment

After using Conquest Snooker, empty the tank completely and drain the whole system. Thoroughly wash inside the tank using a pressure hose, drain the tank and clean any filters in the tank, pump, line and nozzles.

To rinse: After cleaning the tank as above, quarter fill the tank with clean water and circulate through the pump, lines, hoses and nozzles. Drain and repeat the rinsing procedure twice.

To decontaminate: Before spraying cereals (except wheat), maize, sorghum or other sensitive crops, wash the tank and rinse the system as above. Drain and repeat the rinsing procedure. Then quarter fill the tank and add a liquid alkali detergent (eg SURF*, OMO*, DRIVE*) at 0.5L/100L of water and circulate throughout the system for at least fifteen minutes. Drain the whole system. Remove filters and nozzles and clean them separately. Finally flush the system with clean water and allow to drain. Dispose of all water used for cleaning in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots.

Application

Ground Application: Boom spray with flat fan nozzles. Ensure good spray coverage is obtained. Apply using 50 to 110 L water per hectare. The ideal droplet size is 200 to 300 microns VMD.

Aircraft Application: For best results use 20 to 30 L water per hectare and spray at 2m to 3m above the crop. Ideal droplet size is 250 to 350 microns VMD. For rotaryatomizers (micronairs) this can be achieved using blade angles of 55° to 65°. Where booms and nozzles are attached, flat fan nozzles with a spray angle of 65° to 90° should be used.

Avoid applying Conquest Snooker if wind speeds are greater than 5 m/s.

If spraying in calm conditions use larger droplets and reduce flying height.

Compatibility

When Conquest Snooker is applied with compatible herbicides always use a recommended oil. Only use a 100% non-ionic surfactant when mixing with diflufenican based products.

Grass Herbicides: Conquest Snooker is compatible with diclofop methyl, eg Nugrass*.

Broadleaf Herbicides: Some broadleaf weed herbicides are incompatible with Conquest Snooker because they reduce the effectiveness of Conquest Snooker or because in mixture they may cause crop yellowing. For the latest information on compatibility of Conquest Snooker with broadleaf herbicides refer to the Conquest Snooker Compatibility Chart available from your local reseller, FarmWorks Store or the Conquest web site, www.conquestag.com.au

Incompatible: The following herbicides are known to be incompatible with Conquest Snooker: Diuron, LOGRAN[®], Barrel* (bromoxynil + dicamba + MCPA) and all 2,4-D formulations.

Herbicides which are incompatible with Conquest Snooker must be applied separately. In this situation, apply Conquest Snooker first and then allow at least 10 days between its application and application of the broadleaf weed herbicide.

Insecticides: Conquest Snooker is compatible with SUPRACIDE[®].

PRECAUTIONS

Re-entry Period

DO NOT enter treated areas without protective clothing until the spray has dried.

PROTECTION OF CROPS, NATIVE AND OTHER NON- TARGET PLANTS

DO NOT apply on or near shrubs, trees, lawns or crops other than wheat. DO NOT drain or flush equipment on or near desirable trees or other plants, where their roots may extend, or in situations where by movement of soil or by seepage absorption of the herbicide may occur.

DO NOT apply under meteorological conditions or from spraying equipment which could be expected to cause spray to drift onto nearby susceptible plants, adjacent crops, crop lands or pastures. DO NOT allow spray to drift onto adjacent fallow land.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT apply to irrigation drains or channels.

DO NOT contaminate dams, streams, creeks and rivers with the chemical, spray mixture, used containers or equipment washings.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid inhaling spray mist. Avoid contact with the eyes and skin. When opening the container, preparing spray and using the product wear:

- cotton overalls buttoned to the neck and wrist (or equivalent clothing),
- a washable hat,
- elbow- length butyl rubber gloves and
- face shield or goggles.

If product gets on skin, immediately wash area with soap and water. If product gets in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26). If swallowed, do NOT induce vomiting. Give a glass of water. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

MATERIAL SAFETY DATA SHEET

Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.

CONDITIONS OF SALE

Conquest Crop Protection Pty. Ltd. shall not be liable for any loss, injury, damage or death whether consequential or otherwise, whatsoever or whosoever arising whether through negligence or otherwise, in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Conquest skill of judgement in the purchasing or using the same and every person using this product does so at his own risk absolutely. No representative of Conquest has the authority to add or alter these.

