



TECHNICAL NOTES



Tremacide® 120

Flukicide for Cattle and Sheep



Active Constituents

Triclabendazole 120 g/L

Actions

Triclabendazole is a member of the benzimidazole family of drenches. Tremacide® 120 is a ready-to-use, free flowing, aqueous suspension for the treatment and control of acute and chronic fasciolosis in cattle and sheep. Tremacide® 120 is palatable, and may be administered to young, pregnant, weak or debilitated animals.

When used at the recommended rate it combines high efficacy against all stages of susceptible fluke with a high safety margin.

Indications

Treatment and control of acute and chronic fasciolosis in cattle and sheep caused by susceptible early immature, immature and mature liver fluke (*Fasciola hepatica*).

Precautions

In areas where black disease is known to occur, adequate vaccination procedures should be maintained.

Resistance may develop to any drench. Ask a local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.

Withholding Periods

Meat: DO NOT USE less than 21 days before slaughter for human consumption.

Milk: DO NOT USE in lactating cows or ewes where milk or milk products may be used for human consumption. DO NOT USE less



www.jurox.com.au
Customer Service 1800 023 312
® Registered Trademark of Jurox Pty Limited

AUSTRALIA'S
ANIMAL HEALTH COMPANY





TECHNICAL NOTES



than 21 days before calving or lambing in cows or ewes where milk or milk products from treated animals may be used for human consumption. Where this product is accidentally given within this period, or should any cows or ewes give birth earlier than 21 days after treatment, milk will contain residues. This milk must not be used for human consumption or supplied for processing for at least 21 days following treatment. Calves fed this milk within 10 days of slaughter, or lambs fed this milk within 7 days of slaughter, must not be used for human consumption.

Trade Advice

Export Slaughter Interval (ESI)

DO NOT USE less than 56 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Limited on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues)

Dosage and Administration

Shake well before use.

Administer orally, and check accuracy of drenching equipment both before and during use. Maintain cleanliness of drenching equipment and wash out with water after use.

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (bulls, cows, steers; ewes, wethers, rams, lambs etc). Do not underdose. Where there is a large variation in size within the group, the dose should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Dose Rate

Cattle: 5 mL per 50 kg bodyweight

Cattle				
Body weight (kg)	Dose (mL)	5 L treats (head)	10 L treats (head)	20 L treats (head)
Up to 100	10	500	1000	2000
101-150	15	333	666	1333
151-200	20	250	500	1000
201-250	25	200	400	800
251-300	30	166	333	666
301-350	35	142	285	571
351-400	40	125	250	500
401-450	45	111	222	444
451-500	50	100	200	400
501-550	55	90	181	363
551-600	60	83	166	333
601-650	65	76	153	307

Cattle over 650 kg to be dosed at 5 mL per 50 kg bodyweight.

Sheep: 1 mL per 12 kg bodyweight

Sheep				
Body weight (kg)	Dose (mL)	5 L treats (head)	10 L treats (head)	20 L treats (head)
Up to 12	1	5000	10000	20000
13-24	2	2500	5000	10000
25-36	3	1666	3333	6666
37-48	4	1250	2500	5000
49-60	5	1000	2000	4000
61-72	6	833	1666	3333

Sheep over 72 kg to be dosed at 1 mL per 12 kg bodyweight





TECHNICAL NOTES



Safety Directions

Harmful if absorbed by skin contact or swallowed. Avoid contact with the skin. Wash hands after use.

First Aid

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

Presentation

Liquid: 5 L, 10 L, 20 L

Storage

Store below 30°C (room temperature). Store in the original container, tightly closed, in a cool, dry place.

Poisons Schedule

Nil

Registration Numbers

APVMA No. 52463

AU - Tech Note - Tremacide 120 / May 2019 / APPROVED



www.jurox.com.au
Customer Service 1800 023 312
® Registered Trademark of Jurox Pty Limited

AUSTRALIA'S
ANIMAL HEALTH COMPANY

