



Product Name: TREMACIDE 120 FLUKICIDE FOR CATTLE AND SHEEP
APVMA Approval No: 52463/117163

Label Name:	TREMACIDE 120 FLUKICIDE FOR CATTLE AND SHEEP
Signal Headings:	FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	120 g/L TRICLABENDAZOLE
Claims:	<p>For the treatment of susceptible early immature, immature and mature liver fluke (<i>Fasciola hepatica</i>) in cattle and sheep.</p> <p>TREMACIDE 120 contains triclabendazole, a member of the benzimidazole family of drenches. It is a ready-to-use, free flowing aqueous suspension for the treatment and control of acute and chronic fasciolosis of cattle and sheep. TREMACIDE 120 is palatable and may be administered to young, pregnant, weak or debilitated animals.</p> <p>When used at the recommended rate of 12 mg triclabendazole per kg bodyweight for cattle and 10 mg per kg bodyweight for sheep, TREMACIDE 120 combines high efficacy against all stages of susceptible fluke with a high safety margin.</p> <p>Resistance may develop to any drench. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a drench resistance test be conducted before any drench is used.</p>
Net Contents:	5 L, 10 L, 20 L
Directions for Use:	
Restrains:	DO NOT USE in lactating cows or ewes, or within 21 days of calving or lambing where milk or milk products may be used for human consumption.
Contraindications:	
Precautions:	In areas where Black Disease is known to occur, adequate vaccination procedures should be maintained.

Side Effects:	
---------------	--

Dosage and Administration:	<p>SHAKE WELL BEFORE USE</p> <p>Administer orally and check accuracy of drenching equipment both before and during use. Maintain cleanliness of drenching equipment and wash out with water after use.</p> <p>A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (bulls/cows/steers; ewes/wethers/rams/lambs, etc.) Do not underdose. Where there is a large variation in size within the group, the dose should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing.</p> <p>Dose rate: Cattle: 5 mL per 50 kg bodyweight. Sheep: 1 mL per 12 kg bodyweight.</p> <p>[Dose Volume Table for Cattle is Attached]</p> <p>Cattle over 650 kg to be dosed at 5 mL/50 kg.</p> <p>[Dose Volume Table for Sheep is Attached]</p> <p>Sheep over 72 kg to be dosed at 1 mL/12 kg.</p> <p>This section contains file attachment. File Name: 2018-08-13 Tremacide 120 Dose Volume Table for e-label.pdf File Size: 143626 bytes</p>
----------------------------	---

General Directions:	
---------------------	--

Withholding Periods:	<p>MEAT: DO NOT USE less than 21 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE in lactating cows or ewes where milk or milk products may be used for human consumption. DO NOT USE less than 21 days before calving or lambing in cows or ewes where milk or milk products from treated animals may be used for human consumption. Where cows or ewes are accidentally treated within this period or should any cows or ewes calve or lamb earlier than 21 days after treatment, milk will contain residues. This milk must not be used for human consumption or supplied for processing for at least 21 days following treatment. Calves fed this milk should not be slaughtered for human consumption within 10 days. Lambs fed this milk should not be slaughtered for human consumption within 7 days.</p>
----------------------	--

Trade Advice:	<p>TRADE ADVICE—Export Slaughter Interval: DO NOT USE less than 56 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Jurox Pty Limited on 1800 023 312 or the APVMA website (www.apvma.gov.au/residues/).</p>
---------------	---

Safety Directions:	<p>Harmful if absorbed by skin contact or swallowed. Avoid contact with skin. Wash hands after use.</p>
--------------------	---

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	
Additional User Safety:	Refer to Safety Data Sheet for additional information.
Environmental Statements:	Dangerous to fish. Do not contaminate dams, rivers or streams with the drench or used containers.
Disposal:	Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30°C (room temperature). Store in the original container, tightly closed, in a cool, dry place.

Tremacide 120 Flukicide for Cattle and Sheep
Dose Volume Tables

CATTLE				
Bodyweight	Dose	5 L treats	10 L treats	20 L treats
Up to 100 kg	10 mL	500	1000	2000
101 – 150 kg	15 mL	333	666	1333
151 – 200 kg	20 mL	250	500	1000
201 – 250 kg	25 mL	200	400	800
251 – 300 kg	30 mL	166	333	666
301 – 350 kg	35 mL	142	285	571
351 – 400 kg	40 mL	125	250	500
401 – 450 kg	45 mL	111	222	444
451 – 500 kg	50 mL	100	200	400
501 – 550 kg	55 mL	90	181	363
551 – 600 kg	60 mL	83	166	333
601 – 650 kg	65 mL	76	153	307

SHEEP				
Bodyweight	Dose	5 L treats	10 L treats	20 L treats
Up to 12 kg	1 mL	5000	10000	20000
13 – 24 kg	2 mL	2500	5000	10000
25 – 36 kg	3 mL	1666	3333	6666
37 – 48 kg	4 mL	1250	2500	5000
49 – 60 kg	5 mL	1000	2000	4000
61 – 72 kg	6 mL	833	1666	3333