



Product Name: Bovipour DT for control of internal and external parasites on beef and dairy cattle
APVMA Approval No: 86116/116201



Label Name:	Bovipour DT for control of internal and external parasites on beef and dairy cattle
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	5 mg/mL DORAMECTIN
Claims:	For the treatment and control of doramectin sensitive internal and external parasites of cattle. For the treatment and persistent control of gastrointestinal roundworms, lungworms and buffalo fly in cattle. For the control of eye worms, sucking and biting lice, mange mites and cattle ticks.
Directions for Use:	Bovipour DT for control of internal and external parasites on beef and dairy cattle contains doramectin, a member of the macrocyclic lactones family of chemicals. Highly effective in the treatment and control of the following species: - Gastrointestinal roundworms (adults and forth stage larvae) - Ostertagia ostertagi - Small brown stomach worm. - Haemonchus placei - Barber's pole worm - Trichostrongylus axei - Stomach hair worm - T. colubriformis - Black scour worm - Cooperia spp. - Small intestinal worm - Cooperia oncophora - Bunostomum phlebotomum - Hook worm - Oesophagostomum radiatum - Nodule worm - Nematodirus spathiger - Thin necked intestinal worm - Thelazia spp. - Eye worm - Thichuris spp. – Whipworm Adult and Immature Lungworm - Dictyocaulus viviparus

	<p>Sucking Lice</p> <ul style="list-style-type: none"> - Linognathus vituli - Haematopinus eurysternus - Solenopotes capillatus <p>Biting Lice</p> <ul style="list-style-type: none"> - Bovicola bovis <p>Buffalo Fly</p> <ul style="list-style-type: none"> - Haematobia irritans exigua <p>Cattle Tick</p> <ul style="list-style-type: none"> - Rhipicephalus (Boophilus) microplus <p>Mange Mites</p> <ul style="list-style-type: none"> - Chorioptes bovis - Sarcoptes scabiei var. bovis <p>When applied as directed this product can persistently protect against various gastrointestinal roundworms for 21 to 35 days; against lungworms for up to 28 days and against buffalo flies for up to 14 days.</p> <p>Extensive studies have also demonstrated the following levels of persistent protection against reinfection from important worm species:</p> <ul style="list-style-type: none"> - Ostertagia ostertagi - Up to 35 days - Haemonchus placei - Up to 35 days - Trichostrongylus axei - Up to 35 days - Cooperia spp. - Up to 35 days - Cooperia oncophora - Up to 21 days - Dictyocaulus viviparus - Up to 28 days - Oesophagostomum radiatum - Up to 21 days - Bunostomum phlebotomum - Up to 21 days <p>Worm parasites may develop resistance to any anthelmintic. If this</p>
--	--

Net Contents:	<p>250 mL</p> <p>1 L</p> <p>2.5 L</p> <p>3L</p> <p>5 L</p> <p>10 L</p> <p>20 L</p> <p>22 L</p>
---------------	--

Directions for Use:	
---------------------	--

Restrains:	Re-treatment interval: DO NOT re-treat animals for 56 days (8 weeks) after treatment.
------------	---

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

<p>Dosage and Administration:</p>	<p>For external use only. Apply topically along the mid-line of the back in a narrow strip between the withers and tailhead.</p> <p>Use the dose rate, according to the weight of the animal in the weight/dose table.</p> <p>DOSE RATE: 1mL per 10 kg bodyweight.</p> <p>To obtain optimal benefit from the persistent-action against internal parasite, it is recommended that cattle which are set-stocked should be treated twice at eight week intervals. By using in this way, Bovipour DT for control of internal and external parasites on beef and dairy cattle will reduce the build-up of infective larvae on pasture and protect such as animals from parasitic gastro-enteritis and parasitic bronchitis by reducing pasture larval challenge.</p> <p>Refer to the instructions of the dosing cup/applicator on the inner carton for the method of administration.</p> <p>This section contains file attachment.</p>
<p>General Directions:</p>	<p>NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.</p> <p>For best results, do not treat cattle when their hide or hair is wet.</p> <p>Do not apply to areas of skin which are contaminated with mud or manure.</p>
<p>Withholding Periods:</p>	<p>MEAT - DO NOT USE less than 42 days before slaughter for human consumption. MILK - Zero (0) days.</p>
<p>Trade Advice:</p>	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from ZAGRO AUSTRALIA Pty Ltd on 1800 144 884 or the APVMA website (www.apvma.gov.au/residues)</p>
<p>Safety Directions:</p>	<p>Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with skin and eyes. When using the product wear face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash face shield or goggles.</p>
<p>First Aid Instructions:</p>	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.</p> <p>If skin contact occurs, remove contaminated clothing and wash skin thoroughly.</p>
<p>First Aid Warnings:</p>	
<p>Additional User Safety:</p>	<p>Use only in well ventilated surroundings.</p> <p>Flammable - Keep away from heat, sparks, open flame and other sources of ignition.</p> <p>Additional information is listed in the Material Safety Data Sheet.</p>

Environmental Statements:	<p>ENVIRONMENT PROTECTION</p> <p>Doramectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.</p> <p>When used according to label instructions, the product is not likely to have an unintended effect that is harmful to the environment or have an overall adverse impact on dung beetle populations. Increased mortality and impaired development of larvae may be expected, but for a limited period after treatment.</p>
---------------------------	--

Disposal:	<p>For 250 mL and 1 L pack sizes: Dispose of container by wrapping with paper and putting in garbage.</p> <p>For 2.5 L, 3L, 5 L, 10 L, 20L and 22 L pack sizes: Triple-rinse container. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
-----------	--

Storage:	Store at temperatures below 30°C (Room Temperature). Protect from direct sunlight.
----------	--

DOSE RATE: 1 mL per 10 kg bodyweight

WEIGHT (kg)	DOSE (mL)
40 - 50	5
51 - 75	7.5
76 – 100	10
101 – 150	15
151 – 200	20
201 – 250	25
251 – 300	30
301 – 350	35
351 - 400	40
401 - 450	45
451 - 500	50
501 - 550	55
551 – 600	60
601 – 650	65

* Cattle heavier than 650 kg should be dosed at 1 mL per 10 kg. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing.