

SF00-SF06-0008

BOVIPOUR DT

For Control of Internal and
External Parasites On Beef and
Dairy Cattle



Active Content & Action

Doramectin 5 g/L. Doramectin is part of the Macrocylic Lactones group of chemicals. Doramectin works by interrupting the nervous systems of parasites, leading to paralysis and eventual death.

Use

A ready to use pour on to treat doramectin sensitive internal and external parasites in beef and dairy cattle; Including buffalo fly, sucking and biting lice, mange mites, gastrointestinal roundworms, lungworms, and cattle ticks.

Bovipour DT is highly effective in the treatment and control of the following species:

Gastrointestinal Roundworms (adults and fourth-stage larvae)

<i>Ostertagia ostertagi</i> ¹	- Small brown stomach worm
<i>Haemonchus placei</i>	- Barber's pole worm
<i>Trichostrongylus axei</i>	- Stomach hair worm
<i>T. colubriformis</i>	- Black scour worm
<i>Cooperia spp</i> ¹	- Small intestinal worm
<i>Cooperia oncophora</i> ¹	
<i>Bunostomum phlebotomum</i>	- Hook worm ²
<i>Oesophagostomum radiatum</i>	- Nodule worm
<i>Nematodirus spathiger</i>	- Thin necked intestinal worm ²
<i>Thelazia spp</i>	- Eye worm ²

(¹including inhibited larvae, ²adults only)

Adult and Immature Lungworm

Dictyocaulus viviparus

Sucking Lice

Linognathus vituli, *Haematopinus eurysternus*, *Solenopotes capillatus*

Biting Lice

Bovicola bovis

Buffalo Fly

Haematobia irritans exigua.

Cattle Tick³

Rhipicephalus (Boophilus) microplus

Mange Mites

Chorioptes bovis,

(³ Engorged female ticks that drop off for up to 3 days after treatment may lay viable eggs)

Indications

Nil

Doses

100 doses per 5L pack
(average weight 500kg).

Pack Size(s) Available

5L

Australian Poisons Schedule


S6

Presentation

5L - 3 per box.

APVMA Approval No.:

86116



Period of Protection

Studies have also demonstrated the following levels of persistent protection against reinfection from important worm species:

<i>Ostertagia ostertagi</i>	- Up to 35 days
<i>Haemonchus placei</i>	- Up to 35 days
<i>Trichostrongylus axei</i>	- Up to 35 days
<i>Cooperia spp.</i>	- Up to 35 days
<i>Cooperia oncophora</i>	- Up to 21 days
<i>Dictyocaulus viviparus</i>	- Up to 28 days
<i>Oesophagostomum radiatum</i>	- Up to 21 days
<i>Bunostomum phlebotomum</i>	- Up to 21 days

Bovipour DT controls buffalo flies (*Haematobia irritans exigua*) for up to 14 days after treatment.

Worm parasites may develop resistance to any anthelmintic. If this is suspected contact your local veterinary practitioner or advise the manufacturer.

Restrictions

Re-treatment interval: DO NOT re-treat animals for 56 days (8 weeks) after treatment.

Dosage and Administration

FOR EXTERNAL USE ONLY.

Apply topically along the mid-line of the back in a narrow strip between the withers and tailhead. Use the dose rate, according to the weight of the animal in the weight/dose table.

Dose Rate: 1 mL per 10 kg bodyweight.

To obtain optimal benefit from the persistent-action against internal parasite, it is recommended that cattle which are set-stocked should be treated twice at eight-week intervals. By using in this way, **BOVIPOUR DT** will reduce the build-up of infective larvae on pasture and protect such as animals from parasitic gastro-enteritis and parasitic bronchitis by reducing pasture larval challenge.

Cattle heavier than 650Kg should be dosed at 1 mL per 10Kg. A representative sample of animals should be weighed before treatment either with scales or a weigh band. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves, etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight to avoid excessive overdosing.

Treatment of very young animals (less than 40 kg bodyweight) is generally not warranted.

Withholding Periods

Meat	: DO NOT USE less than 42 days before slaughter for human consumption.
Milk	: Zero (0) days

Trade Advice

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. Before using this product, confirm the current ESI from ZAGRO AUSTRALIA Pty Ltd on 1800 144 884 or the APVMA website (www.apvma.gov.au/residues)

Accessories

This product can be used with any pour on applicator designed to apply pour on products to cattle. Contact your local reseller for the different options.

Weight (kg)	Dose (mL)
40 - 50	5
51 - 75	7.5
76 - 100	10
101 - 150	15
151 - 200	20
201 - 250	25
251 - 300	30
301 - 350	35
351 - 400	40
401 - 450	45
451 - 500	50
501 - 550	55
551 - 600	60
601 - 650	65

