

PHARMACHEM

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

PHARMACHEMICAL

MALDISON 50 INSECTICIDE

ACTIVE CONSTITUENT: 500g/L MALDISON (an anti-cholinesterase compound)
SOLVENT: 493g/L hydrocarbon liquid

For the control of a wide range of insect pests in dogs, poultry, pigs,
horses and cattle and their housing as per the Directions For Use Table.



500mL

DIRECTIONS FOR USE: Dilute to recommended application strength with water before applying.

WITHHOLDING PERIODS: MEAT: 7 days; MILK: 5 hours. Not to be fed to puddy calves; EGGS: NIL
(See Directions for Use Table for complete instructions. If the instructions have been detached, contact Pharmachem for a replacement).

SAFETY DIRECTIONS: Harmful if swallowed. Will damage eyes and will irritate the nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and a face shield. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. If product is in eyes, wash it out immediately with water. If product is on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield and contaminated clothing.

FIRST AID: If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre. Phone Australia 13 11 26 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT: Do not contaminate dams, rivers, streams or waterways with the chemical or used containers.

STORAGE: Store below 30°C (Room Temperature) in the closed original container. Do not store in direct sunlight.

DISPOSAL: Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.

APVMA Approval No: 33021/0210

Batch No:

Expiry Date:

pharmachem 
CARING FOR YOUR CREATURE'S COMFORTS

Unit 6, 70 Fison Ave West, Eagle Farm QLD 4009

www.pharmachem.com.au

FOR FULL INSTRUCTIONS
PEEL BACK LABEL

PEEL
BACK



SCAN ME

DIRECTIONS FOR USE TABLE



DIRECTIONS FOR USE:

ANIMAL SITUATION		PEST	DILUTION RATE OF PRODUCT IN WATER		CRITICAL COMMENTS	MIXING MALDISON 50
			SPRAY/HIGH VOLUME	ALTERNATIVE APPLICATION		
Dogs over 3months	Fleas & Lice		50mL/10L	Saturate the animal. Repeat in 7 days. Use the residue for spraying bedding etc. Do not immerse the animal.		MALDISON 50 gives a spontaneous emulsion in water and may be added directly to the water in the spray vat with the agitators running. Mixes readily with hard or soft water.
	Adult Brown Dog Tick & Sarcoptic Mange		50mL/5L			
Poultry Pests	Lice & Mites (Poultry Houses)		500mL/25L	Thoroughly treat nesting boxes, litter and walls with spray.	NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION	WITHHOLDING PERIODS: MEAT: DO NOT USE less than 7 days before slaughter for human consumption. MILK: Milk collected from treated animals within 5 hours following treatment must not be used for human consumption or processing. This milk should not be fed to paddy calves. EGGS: NIL
			500mL/8L	Paint onto roosts. The above applications should be repeated 8-14 days later to destroy lice hatching from eggs present at the first treatment.		
	Lice or Tropical Fowl Mite	500mL/100L	Spray birds at the rate of 50L/1000 birds.			
	Ticks		500mL/8L	Treat the walls, nesting boxes and roosts of poultry houses very thoroughly with a pressure sprayer, paying particular attention to cracks and crevices in timber where the lice hide.		
Pigs	Pig Lice & Sarcoptic Mange		250mL/20L	Thoroughly spray all pigs and sheds.		
Horses	Lice, Ticks, Sandflies		250mL/20L	To spray 10 horses (i.e. 2 Litres of spray per horse).		
Cattle	Lice	1L/100L		Spraying at the rate of 4.5 Litres per head is usually enough.		
Fly Control			500mL/40L	The addition of sugar or molasses to the spray will result in a better kill.		TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For ESI advice, please contact the manufacturer on (07 3868 0333) before using this product.
			250mL/20L			
Mosquito Control	Larvae	1L/ha		Dilute with water as required. Treat breeding areas as soon as larvae are seen.		APVMA Approval No: 33021/0210
	Adults	500mL/ha		Dilute with water as required. Application should be timed for periods of major emergence of adults.		