



Product Name: eLabel PHARMACHEMICAL MALDISON 50 INSECTICIDE

APVMA Approval No: 33021/133564

Label Name:	PHARMACHEMICAL MALDISON 50 INSECTICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	500 g/L MALDISON 493 g/L LIQUID HYDROCARBON
Claims:	For the control of a wide range of insect pests in dogs, poultry, pigs, horses and cattle as per the Directions for Use Table.
Net Contents:	250 mL 500 mL 5 L
Directions for Use:	This section contains file attachment.
Restrains:	
Contraindications:	
Precautions:	
Side Effects:	
Dosage and Administration:	This section contains file attachment.

General Directions:	
Withholding Periods:	MEAT: DO NOT USE less than 7 days before slaughter for human consumption MILK: Milk collected from treated animals within 5 hours following treatment must not be used for human consumption or processing. This milk should not be fed to bobby calves. EGGS: NIL.
Trade Advice:	Export Slaughter Interval (ESI): This product does not have an ESI established. For ESI advice, please contact the manufacturer on 07 38680333 before using this product.
Safety Directions:	Harmful swallowed. Will damage eyes and will irritate the nose and throat and skin. Avoid contact with eyes and skin. Do not inhale vapour. Repeated exposure may cause allergic disorders. When opening the container and preparing the spray wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and a face shield. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. If product is in eyes, wash it out immediately with water. If product is on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and face shield and contaminated clothing.
First Aid Instructions:	If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre. Phone Australia 131 126 or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.
First Aid Warnings:	
Additional User Safety:	ADDITIONAL USER SAFETY INFORMATION. Additional information is listed in the safety data sheet which can be obtained from the supplier.
Environmental Statements:	ENVIRONMENTAL PROTECTION: Do not contaminate dams, rivers, streams or waterways with the chemical or used containers.
Disposal:	Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.
Storage:	Store below 30 °C (room temperature) in the closed original container. Do not store in direct sunlight.

Pharmachemical Maldison 50 Insecticide				
DIRECTIONS FOR USE:				
Animal Situation	Pest	Dilution rate of product in water		Critical Comments
		Spray/High Volume	Alternative Application	
Dogs over 3 months	Fleas & Lice		50mL/10L	Saturate the animal. Repeat in 7 days. Use the residue for spraying bedding etc. Do not immerse the animal.
	Adult Brown Dog Tick & Sarcoptic Mange		50mL/5L	Saturate the animal. Repeat in 7 days. Use the residue for spraying bedding etc. Do not immerse the animal.
Poultry Pests	Lice & Mites (Poultry Houses)		500mL/25L (ie 1% Maldison 50)	Thoroughly treat nesting boxes, litter and walls with spray.
			500mL/8L (ie 3% Maldison 50)	Paint onto roosts. The above applications should be repeated 8-14 days later to destroy lice hatching from eggs present in the first treatment.
	Lice or Tropical Fowl Mite	500mL/100L		Spray birds at the rate at 50 L/1000 birds
	Ticks		500mL/8L (ie 3% Maldison 50)	Treat the walls, nesting boxes and roosts of poultry houses very thoroughly with a pressure sprayer, paying particular attention to cracks and crevices in timber where the ticks hide
Pigs	Pig Lice & Sarcoptic Mange		250mL/20L	Thoroughly spray all pigs and sheds
Horses	Lice, Ticks, Sandflies		250mL/20L	To spray 10 horses (ie 2 L of spray per horse)
Cattle	Lice	1L/100L		Spraying at the rate of 4.5 L per head is usually enough.
Fly Control		500mL/40L		The addition of sugar or molasses to the spray will result in a better kill.
			250mL/20L	
Mosquito Control	Larvae	1L/ha		Dilute with water as required. Treat breeding areas as soon as larvae are seen.
	Adults	500mL/ha		Dilute with water as required. Application should be timed for periods of major emergence of adults.

MIXING – MALDISON 50 gives a spontaneous emulsion in water and may be added directly to the water in the spray vat with the agitators running. Mixes readily with hard or soft water

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIODS:

MEAT: DO NOT USE less than 7 days before slaughter for human consumption

MILK: Milk collected from treated animals within 5 hours following treatment must not be used for human consumption or processing. This milk should not be fed to bobby calves.

EGGS: NIL

TRADE ADVICE: EXPORT SLAUGHTER INTERVAL (ESI): This product does not have an ESI established. For ESI advice, please contact the manufacturer on (07 38680333) before using this product.