



MOXISTAR[®]

LONG ACTING INJECTION FOR CATTLE

ACTIVE CONSTITUENT: 100 MG/ML MOXIDECTIN

PREMIUM
CHOICE FOR TICK
MANAGEMENT

PERSISTENT
CONTROL OF
CATTLE WORMS

NO KNOWN
EFFECT ON
DUNG BEETLES

Moxistar Long Acting Injection has been specially formulated to provide a prolonged release of moxidectin in cattle. It is ideal for long term protection against key internal and external parasites including cattle tick. With persistent activity against internal parasites for up to 120 days (see below table for details) or cattle tick for at least 51 days, no other product offers longer term protection ensuring optimum health, performance and fertility.

MOXISTAR LONG ACTING INJECTION FOR CATTLE is effective against sensitive strains of the following parasites:

Internal parasites: Mature (adult) and immature (L4): Barber's pole worm, *Ostertagia ostertagi* / *Ostertagia lyrata* (including inhibited larvae), Black scour worms, Small intestinal worm, Nodule worm, Hookworm, Whipworm, Lungworm and Adult nematodes.

External parasites: Lice (Sucking lice). Aids in the control of *Bovicola bovis* (biting lice), Mites and Ticks.

PROTECTION PERIOD

When MOXISTAR LONG ACTING INJECTION FOR CATTLE is used at the recommended dose rate as a single subcutaneous injection, it prevents re-infection of cattle with parasites as in the following table:

Parasite Species	Persistent Protection Period
<i>Ostertagia</i> spp.	112 days
<i>Haemonchus</i> spp.	120 days
<i>Trichostrongylus axei</i>	72 days
<i>Cooperia</i> spp.	21 days
<i>Dictyoacaulus viviparus</i>	120 days
<i>Oesophagostomum radiatum</i>	120 days
<i>Linognathus vituli</i>	133 days
<i>Rhipicephalus (Boophilus)</i>	51 days

APPLICATOR

5ml Vaccinator

WITHHOLDING PERIODS

MEAT: 56 days.

MILK: DO NOT USE in lactating cows or within 80 days of calving where milk may be used or processed for human consumption.

ESI: 108 days. DO NOT USE in cattle that may be exported live.

STORAGE

Store below 30°C (room temperature). Protect from light.


COMPARABLE PRODUCT

Virbac Cydectin Long Acting Injection for Cattle

DOSING & ADMINISTRATION

Dose Rate: 1 mg moxidectin/kg liveweight (1 mL/100 kg liveweight)

INJECT ONLY by subcutaneous injection into the back of the ear (as directed below). Do not inject anywhere else in the animal.



Weight range (kg)	Dose volume (mL)	Treatments per 500 mL
* 100	1.0	500
101-150	1.5	333
151-200	2.0	250
201-250	2.5	200
251-300	3.0	166
301-350	3.5	142
351-400	4.0	125
401-450	4.5	111
451-500†	5.0	100

* DO NOT USE IN CATTLE LESS THAN 100 KG

† DO NOT USE IN CATTLE GREATER THAN 500 KG

For more information please visit hunterriverco.com.au

Virbac & Cydectin are registered trade marks of Virbac (Australia) Pty Ltd.

