

## WITHHOLDING PERIODS

**MEAT:** Zero (0) days

**MILK:** Zero (0) days

## TRADE ADVICE

**EXPORT SLAUGHTER INTERVAL (ESI):**

**Zero (0) days.** Before using this product, confirm the current ESI from Bayer on 1800 678 368 or APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)).

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## LIMITATION OF LIABILITY

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

## DISPOSAL

Dispose of empty container by wrapping with paper and placing in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labeled 'sharps' container.

FOR ANIMAL TREATMENT ONLY



# HIDEJECT®

## Vitamin A, D3 and E Injection

### ACTIVE CONSTITUENTS:

**18mg/mL (60 000IU/mL) VITAMIN A as PALMITATE (RETINYL PALMITATE), 12.5mg/mL (500 000IU/mL) VITAMIN D3 (CHOLECALCIFEROL), 25mg/mL (25 IU/mL) VITAMIN E (DI-ALPHA TOCOPHEROL ACETATE)**

As an aid in the prevention of milk fever in cows, ewes and sows.



100mL

## STORAGE

Store below 30°C (room temperature).  
Protect from light.

Toxic if inhaled. Causes damage to organs through prolonged or repeated exposure if swallowed.

## CUSTOMER INFORMATION LINE

**1800 678 368**

from anywhere in Australia  
9am to 4pm (Eastern Time) Monday to Friday  
[www.farmadvisor.com.au](http://www.farmadvisor.com.au)

# Bayer

Bayer Australia Ltd, ACN 000 138 714  
875 Pacific Highway, Pymble NSW 2073

HIDEJECT® is a registered trade mark of the Bayer Group.

APVMA Approval No. 35995/105384



9 310160 824639

## DIRECTIONS FOR USE

Use the contents within 24 hours of first broaching the vial. Discard the unused portion.

## DOSAGE AND ADMINISTRATION

By deep intramuscular injection.

**Cows:** To aid in the prevention of milk fever. Administer 5mL 4–6 weeks before calving when induction methods are used and 8 days prior to natural calving.

**Ewes:** 1–4mL 6–8 weeks before lambing.

**Sows:** 3–4mL 4 weeks before farrowing.

## CAUTION: AVOID CARCASS DAMAGE

1. Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. Use needles of appropriate gauge and length.
5. As far as possible, avoid injection of animals during wet weather or under dusty conditions.
6. This product should be injected only into muscle tissue.
7. If possible, inject into muscle tissue on side of neck.