

Product Name: Fastrac Pro Blox Rodenticide
 APVMA Approval No: 88449/132270



Label Name:	Tomcat with Bromethalin Blox Rodenticide
-------------	--

Signal Headings:	<p>POISON</p> <p>KEEP OUT OF REACH OF CHILDREN</p> <p>READ SAFETY DIRECTIONS BEFORE OPENING OR USING</p>
------------------	--

Constituent Statements:	<p>ACTIVE CONSTITUENT: 0.1 g/kg BROMETHALIN</p> <p>Contains DENATONIUM BENZOATE (BITTERING AGENT)</p>
-------------------------	---

Mode of Action:	
-----------------	--

Statement of Claims:	<p>Label A -Ready to use block bait to control rats and mice in damp or dry situations in and around industrial, commercial and agricultural buildings. Rats and Mice usually consume a lethal dose in a single night's feeding</p> <p>Label B- DISPOSABLE non-refillable Bait Station</p> <p>[28 g] A pre-filled ready to use bait station to control mice in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings. Mice usually consume a lethal dose in a single night's feeding</p> <p>[113 g] A pre-filled ready to use bait station to control rats and mice in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings. Rats and Mice usually consume a lethal dose in a single night's feeding</p>
----------------------	--

Net Contents:	150g - 10kg
---------------	-------------

Restrains:	<p>Label A</p> <p>DO NOT use in domestic situations.</p> <p>DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.</p> <p>DO NOT place baits more than 30 m from buildings or permanent structures.</p> <p>DO NOT place baits in the open.</p> <p>DO NOT apply baits in crops.</p> <p>DO NOT place baits in areas liable to flooding.</p> <p>DO NOT place bait in areas where there is a possibility of contaminating water, or surfaces that come into direct contact with water.</p>
------------	--

	<p>Securable, tamper resistant bait stations MUST be used at all locations to prevent access to non-target animals and children.</p> <p>Label B - non-refillable DISPOSABLE Bait Station DO NOT exceed 3 m between bait stations for mice, or 9 m for rats. DO NOT place baits more than 30 m from buildings or structures. DO NOT place baits in the open. DO NOT apply baits in crops. DO NOT place baits in areas liable to flooding. DO NOT place bait in areas where there is a possibility of contaminating water, or surfaces that come into direct contact with water.</p>
--	--

Directions for Use:	This section contains file attachment.
---------------------	--

Other Limitations:	
--------------------	--

Withholding Periods:	
----------------------	--

Trade Advice:	
---------------	--

General Instructions:	<p>Tomcat with Bromethalin Blox Rodenticide is a highly palatable block bait for the control of rats and mice including those resistant to anticoagulants. For use in and around buildings. [Optional statements: [Norway rats, Roof rats and House mice usually consume a lethal dose in a single night's feeding, but it may take two or more days from the time of bait consumption for these rodents to die]. [Rats and Mice cease feeding after consuming a toxic dose.]]</p> <p>Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls or rafters, by gnawed openings, beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Replace contaminated or spoilt bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed.</p>
-----------------------	---

Resistance Warning:	
---------------------	--

Precautions:	<p>RE-HANDLING Cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves and a disposable dust facemask, covering mouth and nose, must be worn during cleaning/disposal operations of bait/bait stations (for example, sweeping/pan broom, washing/hosing) and when rehandling bait, bait waste or rodent carcasses. Rodent carcasses should be collected and sealed in plastic bags prior to disposal in accordance with the disposal statement. Clothing must be laundered after each day's use.</p> <p>PRECAUTIONS DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed, or surfaces that come into direct contact with food.</p>
--------------	---

Protections:	<p>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</p> <p>Toxic to birds and other native wildlife. Place baits so as to prevent access to non-target animals. Securable, tamper resistant bait stations must be used at all locations to prevent access to non-target animals. DO NOT use on sites where vulnerable, threatened, endangered or critically endangered species are known to occur. Any incidents where it is suspected that non-target animals may have been poisoned should be reported to State or Territory authorities.</p> <p>Very toxic to aquatic life. DO NOT contaminate wetlands or watercourses with this product or used containers.</p>
--------------	--

Storage and Disposal:	<p>Store product in original container, tightly closed and away from feed or foodstuffs. Bait storage facility to be locked at all times when not in use. Store in a locked storage facility away from children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Untaken baits should be recovered at the end of a baiting campaign and delivered to an approved waste management facility.</p> <p>To the extent possible, animal carcasses should be recovered during and for 14 days after a baiting campaign and be destroyed by burning or burial according to the requirements of the State or Territory in which use has occurred.</p>
-----------------------	---

Safety Directions:	<p>Harmful if swallowed. May irritate the eyes and skin. Do not cut individual baits. Avoid contact with eyes and skin. Do not touch the bait. If on skin and after each baiting, wash thoroughly with soap and water. When using the product do not eat, drink or smoke. When using the product wear cotton overalls, buttoned to the neck and wrist (equivalent clothing) and gauntlet length (chemical resistant) gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves and contaminated clothing.</p>
--------------------	---

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126; New Zealand 0800 764 766. If poisoning occurs, get to a doctor or hospital quickly.</p>
-------------------------	---

First Aid Warnings:	<p>ADDITIONAL USER SAFETY INFORMATION</p> <p>WARNING – Skin contact may be dangerous. Take every precaution to avoid contact. Securable, tamper resistant bait stations MUST be used at all locations to prevent access by children.</p> <p>Wash off after spillage and after use. Should not be used in areas accessible to children. Possible risk of irreversible effects.</p>
---------------------	--

DIRECTIONS FOR USE

LABEL A

Situation: In and around industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.

Pest

Mice (*Mus musculus*)

Rats (*Rattus norvegicus*, *Rattus rattus*)

For 20 g and 28 g block:

Rate

Mice: 1-2 blocks per bait station.

Rats: 2-20 blocks per bait station.

For mice: use single blocks. In very high infestation or fast uptake of the bait use 2 blocks.

For rats: use 2 or more blocks depending on level of infestation. In very high infestation or fast uptake use more blocks per bait station.

Baiting strategy

Set out baits in places frequented by mice and rats but not more than 30 m from buildings or permanent structures. Bait stations must be secured. For rats: place 2 or more blocks per bait station at intervals of 5-9 m depending on level of uptake and infestation. For mice: place 1-2 blocks per bait station at 2-3 m intervals. In very high infestations, use the shorter distance between bait stations.

For 15 g block:

Rate

Mice: 2 - 6 blocks per bait station.

Rats: 4 - 40 blocks per bait station.

For mice: use 2 blocks. In very high infestation or fast uptake of the bait use up to 6 blocks.

For rats: use 4 or more blocks depending on level of infestation. In very high infestation or fast uptake use more blocks per bait station.

Baiting strategy

Set out baits in places frequented by mice and rats but not more than 30 m from buildings or structures. Bait stations must be secured. For rats: place 4 or more blocks per bait station at intervals of 5-9 m depending on level of uptake and infestation. For mice: place 2-6 blocks per bait station at 2-3 m intervals. In very high infestations use the shorter distance between bait stations.

For 5 g and 10 g blocks:

Rate

Mice: 3-6 blocks per bait station.

Rats: 6-60 blocks per bait station.

For mice: use 3 or more blocks. In very high infestation or fast uptake of the bait use up to 6 blocks.

For rats: use 6 or more blocks depending on level of infestation. In very high infestation or fast uptake use more blocks per bait station.

Baiting strategy

Set out baits in places frequented by mice and rats but not more than 30 m from buildings or permanent structures. Bait stations must be secured. For rats: place 6 or more blocks per bait station at intervals of 5-9 m depending on level of uptake and infestation. For mice: place 3 or more blocks per bait station at 2-3 m intervals. In very high infestations use the shorter distance between bait stations.

For 225 g blocks:

Rate

Mice: 1 block per bait station.

Rats: 1-2 blocks per bait station.

For rats: use 1 or more blocks depending on level of infestation. In very high infestation or fast uptake use 2 blocks per bait station.

Baiting strategy

Set out baits in places frequented by mice and rats but not more than 30 m from buildings or permanent structures. Bait stations must be secured. For rats: place 1 or 2 blocks per bait station at intervals of 5-9 m depending on level of uptake and infestation. For mice: place 1 block per bait station at 2-3 m intervals. In very high infestations use the shorter distance between bait stations.

Critical Comments [for all sizes]

Eliminate, as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Securable, tamper resistant bait stations must be used at all locations to prevent access to non-target animals. Inspect bait stations on a regular basis. Replace eaten or old/deteriorated baits. If eaten quickly, increase the number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks may be necessary to reduce rat/mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

LABEL B – Non refillable DISPOSABLE Bait Station

Situation: In and around domestic homes, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.

Pest

Mice (*Mus musculus*)

Rats (*Rattus norvegicus*, *Rattus rattus*)

28 g Non-refillable bait station (Mice only)

Baiting strategy

Set out baits in places frequented by mice but not more than 30 m from buildings or permanent structures. Bait stations must be secured. Place the bait stations at 2-3 m intervals. In very high infestations use more bait stations and use the shorter distance between bait stations.

113g Non-refillable bait station

Baiting strategy

Set out baits in places frequented by mice and rats but not more than 30 m from buildings or permanent structures. Bait stations must be secured. For rats: place bait station at intervals of 5-9 m depending on level of uptake and infestation. For mice: place bait station at 2-3 m intervals. In very high infestations use more bait stations and use the shorter distance between bait stations.

Critical Comments

Eliminate, as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Inspect bait stations on a regular basis. Replace eaten or old/deteriorated baits. If eaten quickly, increase the number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks may be necessary to reduce rat/mouse numbers.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.