

NO IMAGE

BACK

DIRECTIONS FOR USE

Use the contents within 24 hours of first broaching of the pack. Discard the unused portion.

Dosage: (By subcutaneous injection)

Cattle: Small Cows 500mL
Heavier Cows 500-1000mL

Sheep: Small Breeds 40-70kg: 60-100mL
Heavier Breeds 70-100kg: 100-150mL

GENERAL DIRECTIONS

Caution: Avoid carcass damage

1. If possible warm to body temperature prior to injection.
2. Maintain cleanliness at all times.
3. Remove the fluid bag from the overpouch.
4. Snap off the tip of the needle connector at the end of the delivery tube ensuring the solution can flow freely and connect the sterile needle provided.
5. Disinfect the injection site and insert the needle.
6. Subcutaneous injections should be given under the skin of the neck or over the ribs. To assist the flow of the injection, gently squeeze the bag to inject the contents.
7. Large volumes should be distributed over more than one injection site.
8. If complete recovery does not occur within 3-4 hours repeat injection and seek veterinary advice. Use of this product intravenously may be hazardous to the animal and advice from a registered veterinarian should be sought.

WITHHOLDING PERIODS

MEAT: Zero (0) days

MILK: Zero (0) days

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): Zero (0) days.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

LIMITATION OF LIABILITY

Failure to use the product strictly as directed may be illegal, prove dangerous or render the product ineffective. Notwithstanding anything else contained on this label, and subject to any rights you have under the Australian Consumer Law and any other applicable law, Bayer Australia Limited's liability is limited to replacing this product or supplying equivalent product or paying the cost of replacing this product, at its discretion.

DISPOSAL

Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' container. Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE

Store below 25°C (air conditioning). Protect from light.

Bayer

Bayer Australia Ltd

875 Pacific Highway

PYMBLE NSW 2073

ACN 000 138 714

FLOPAK® is a registered trademark of


Bayer AG, Leverkusen, Germany.

APVMA Approval No. 47310/6/1197

CUSTOMER INFORMATION LINE
1800 678 368
TOLL FREE from anywhere in Australia
 9am to 4pm (Eastern Time) Monday to Friday
www.farmedvisor.com.au



81902569
FLO1207-PCH



500mL



Sterile needle and delivery tube included

Ready to use bag

STERILE NON-PYROGENIC INJECTION FLUID

For the treatment of milk fever, grass tetany and associated metabolic conditions in cattle and sheep

ACTIVE CONSTITUENTS: CALCIUM (as Calcium Gluconate)


27.5g/L (equivalent to 330g/L Calcium Borogluconate),

MAGNESIUM (as Magnesium Hypophosphite) 4.7g/L,

PHOSPHORUS (as Hypophosphite) 12.2g/L,

GLUCOSE 182.0g/L

**Calcium and Magnesium solution
with Phosphorus and Glucose**



FLOPAK® PLUS
(4 IN 1)

FOR ANIMAL TREATMENT ONLY

FRONT

NO IMAGE

10mm
15x5mm
10mm