

Product Name: POLE-AXE SHEEP DRENCH
APVMA Approval No: 84466/110473



Label Name:	POLE-AXE SHEEP DRENCH
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	800 g/kg NAPHTHALOPHOS (An anticholinesterase compound)
Claims:	<p>For the control of organophosphate susceptible strains of gastrointestinal mature and immature roundworms in sheep and lambs.</p> <p>POLE-AXE contains naphthalophos, a member of the organophosphorus family of chemicals. It is effective against organophosphate susceptible strains of mature and immature stages of Barber's Pole worm (<i>Haemonchus contortus</i>), Stomach Hair worm (<i>Trichostrongylus axei</i>), Black Scour worm (<i>Trichostrongylus</i> spp.), Small Brown Stomach worm (<i>Ostertagia</i> spp.), and mature stages of Thin Necked Intestinal worm (<i>Nematodirus</i> spp.) in sheep and lambs.</p>
Net Contents:	750 g 1.5 kg
Directions for Use:	
Restrictions:	DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.
Contraindications:	This product is contraindicated for use in severely debilitated, stressed or drought affected animals or those affected by lupinosis. This product should not be used on lambs less than 6kg.
Precautions:	This product must not be used at the same time as any other drench without veterinary advice.

	<p>Do not treat thirsty, exhausted sheep or those suffering from poor nutrition. Allow animals access to water after treatment.</p> <p>Use care when drenching. Do not misdirect dosage into the lungs.</p> <p>Check the accuracy of drench gun before starting to treat.</p>
--	---

Side Effects:	
---------------	--

Dosage and Administration:	
----------------------------	--

General Directions:	<p>Mixing instructions: POLE-AXE mixes readily with all types of water. Its high suspendibility ensures a consistent uniform suspension during drenching. If product is left standing, shake well before using.</p> <p>Remove large cap and half fill container slowly with water. Replace cap and shake well. Then remove cap and top up with water to the fill-line shown. Replace cap and shake well for 3 minutes. POLE-AXE may be mixed with benzimidazole and levamisole based drench product under veterinary advice. POLE-AXE Sheep Drench is compatible with copper sulphate.</p> <p>POLE-AXE will control strains resistant to other drench groups but resistance may develop to any drench product. Advice on internal parasite management to reduce development of resistance should be obtained from your veterinary practitioner or animal health consultant. It is advisable that a resistance test be conducted before any parasite treatment is used. If drench failure is suspected, seek veterinary confirmation and notify the manufacturer</p>
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 7 days before slaughter for human consumption.</p> <p>MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.</p>
----------------------	---

Trade Advice:	<p>EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export. Before using this product, confirm the current ESI from the registration holder on (02) 9499 7370 or the APVMA website (www.apvma.gov.au/residues).</p>
---------------	---

Safety Directions:	<p>Very Dangerous. Product and drench are poisonous if absorbed by skin contact, inhaled or swallowed. May irritate the eyes and skin. Repeated exposure may cause allergic disorders. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhale dust. When opening the container and preparing drench, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), PVC or rubber apron, elbow length chemical resistant gloves and a disposable dust face mask covering mouth and nose. When using the prepared drench wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), PVC or rubber apron, and elbow-length chemical resistant gloves. If clothing becomes contaminated with product remove clothing immediately. If product or drench on skin, immediately wash area with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use wash gloves and contaminated clothing.</p>
--------------------	--

First Aid Instructions:	<p>If swallowed, splashed on skin or in eyes, or inhaled, contact a Poisons Information Centre (Phone Australia 131126) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	
---------------------------	--

Disposal:	Thoroughly empty container into backpack. Triple rinse container. Dispose of rinsate, and bury containers below 500mm, in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.
-----------	--

Storage:	Store below 30°C (Room Temperature) in original, tightly closed container, in a safe place.
----------	---

**SHEEP SHOULD BE WEIGHED. SELECT THE CORRECT DOSE FOR EACH SHEEP.
DO NOT UNDERDOSE OR OVERDOSE.**

Bodyweight Kg	Dose Volume mL of prepared drench	Number of Sheep treated per pack [750 g]	Number of Sheep treated per pack [1.5 kg]
6 – 7.5	2	2500	5000
7.6 – 9	2.5	2000	4000
9.1 – 12	3	1666	3333
12.1 – 16	4	1250	2500
16.1 – 20	5	1000	2000
20.1 – 28	6.5	769	1538
28.1 – 40	9	555	1111
40.1 – 52	13	384	769
52.1 – 68	17	294	588
68.1 – 90	22	227	454