



Product Name: WSD SPURT OFF-SHEARS, POUR-ON SHEEP LICE CONTROL  
APVMA Approval No: 39065/ 120525



Label Name:	WSD SPURT OFF-SHEARS, POUR-ON SHEEP LICE CONTROL
Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	25 g/L CYPERMETHRIN SOLVENT: DIETHYLENE GLYCOL MONOBUTYL ETHER 935 g/L
Claims:	For the control of pyrethroid susceptible strains of body lice on off-sheers sheep
Net Contents:	1L 20L 5L
Directions for Use:	
Restrains:	
Contraindications:	
Precautions:	
Side Effects:	
Dosage and Administration:	Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams and lambs). DO not underdose. A representative sample of animals should be weighed before treatment either with scales or weighband. Where there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing.

	<p>Sheep in excess of 75kg dose at 1mL per 5kg</p> <p>For maximum benefit and efficiency, pour-ons need to be applied correctly in accordance with their instructions. A) Prior to the start of shearing, check your SPURT applicator for correct operation and calibration. B) Apply SPURT to shorn sheep up to 24 hours off shears using the correct and full dose. Control is achieved in 5-6 weeks. C) Apply the SPURT dose as a single backline strip (40mm to 60mm) along the centre of back – start at the shoulder and finish at the rump (20 cm to 40 cm). D) Ensure all shorn, treated sheep go into a cleanly mustered paddock and are not mixed with untreated sheep.</p> <p>Applied correctly SPURT puts a full strength dose of lousicide on every sheep and works well even under such adverse conditions as wet weather or wet sheep. A scorable blue marker dye is included in the product for specific identification of treated animals.</p> <p>This section contains file attachment.</p>
--	---

General Directions:	<p>Shake or stir contents before using this product.</p> <p>Must only be used within 24 hours of shearing. Do not treat ewes less than 6 weeks before lambing commences as live lice can persist and the lambs. Do not treat lambing ewes or ewes with lambs at foot. Do not mix treated sheep with untreated sheep until 6 weeks after treatment. Do not use on sheep which have not been cleanly shorn. DO not use on female sheep which are producing or may in the future produce milk or milk products for human consumption.</p> <p>Not to be used for any purpose, or in any manner, contrary to this label unless authorised under appropriate legislation.</p>
---------------------	---

Withholding Periods:	<p>Meat: do not use less than three days before slaughter for human consumption. Milk: Do not use on ewes which are producing or may in the future produce milk that may be used or processed for human consumption. WOOL HARVEST INTERVAL: DO NOT use less than 2 months before shearing or fibre collection.</p>
----------------------	--

Trade Advice:	<p>Export slaughter interval (ESI) Do not use less than 14 days before slaughter for export. The ESI on this label was correct at the time of printing of label approval. Before using this product confirm the current ESI from the manufacturer website <a href="http://www.wsdagribusiness.com">www.wsdagribusiness.com</a> Use of this product may result in wool residues that may not comply with European Union environmental quality standards.</p> <p>Wool treated with this product may contain detectable residues; adequate treatment records should be kept and made available, if requested by wool buyers.</p>
---------------	---

Safety Directions:	<p>Product is poisonous if swallowed. It will irritate the eyes and skin. Facial skin contact may cause temporary facial numbness. Sensitive workers should use protective clothing. Avoid contact with eyes and skin. Wear elbow length PVC gloves. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use wash gloves and contaminated clothing.</p>
--------------------	---

First Aid Instructions:	<p>If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor. If poisoning occurs contact a doctor or Poisons Information Centre (Phone 13 11 26).</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	<p>Sheep Rehandling Interval: Two weeks. If sheep must be handled during this interval, cotton overalls buttoned to the neck (or equivalent clothing) should be worn.</p> <p>Material Safety Data Sheet (MSDS): Additional information is listed in the Material Safety Data Sheet, which can be obtained from the supplier.</p>
Environmental Statements:	<p>Environmental protection</p> <p>Dangerous to fish and bees. Do not contaminate streams, rivers and waterways with the product or used containers.</p>
Disposal:	<p>Triple or (preferably) pressure rinse container. Dispose of rinsate in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Do not dispose of undiluted chemical on-site. Return clean container for recycling where this is an option or for disposal at landfill authorized to accept that waste. If neither of these options are available, bury the container below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers should not be burned.</p>
Storage:	<p>Store below 30°C (room temperature) in the closed, original container in a dry, well-ventilated area and protect from sunlight.</p>

## Dosage Table

Sheep	Dose
Under 10Kg	2mL
11-20Kg	4mL
21-30Kg	6mL
31-40Kg	8mL
41-50Kg	10mL
51-60Kg	12mL
61-70Kg	14mL
71-75Kg	15mL

Sheep in excess of 75Kg dose at 1mL per 5Kg