



Product Name: IVOMEK LV Pour-On Ivermectin Endectocide for Cattle  
APVMA Approval No: 92705/136369

Label Name:	IVOMEK LV Pour-On Ivermectin Endectocide for Cattle
-------------	---

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	10 mg/mL Ivermectin
-------------------------	---------------------

Claims:	<p>For the treatment and control of ivermectin sensitive internal and external parasites of beef and dairy cattle.</p> <p>Ivomec LV Pour-On contains Ivermectin, a member of the macrocyclic lactones. It is effective against ivermectin sensitive strains of internal and external parasites when applied to the skin along the backline of cattle.</p> <p><b>INDICATIONS</b> For the treatment and control of adult and some immature stages of:</p> <p>Gastro-intestinal Roundworms Barber's Pole Worm (<i>Haemonchus placei</i>) Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (including inhibited stages) Small Intestinal Worm (<i>Cooperia oncophora</i>) (<i>Cooperia punctata</i>) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Hookworm (<i>Bunostomum phlebotomum</i>) Large Bowel Worm (<i>Oesophagostomum radiatum</i>) Whipworm (<i>Trichuris</i> spp.) Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>) Intestinal Threadworm (<i>Strongyloides papillosus</i>)</p> <p>Lungworms (<i>Dictyocaulus viviparus</i>)</p> <p>Sucking and Biting Lice Long Nosed Sucking Lice (<i>Linognathus vituli</i>) Short Nosed Sucking Lice (<i>Haematopinus eurysternus</i>) Biting Lice (<i>Bovicola bovis</i>) Little Sucking Lice (<i>Solenopotes capillatus</i>) Mites (<i>Chorioptes bovis</i>, <i>Sarcoptes scabiei</i> var <i>bovis</i>) Buffalo Fly (<i>Haematobia irritans exigua</i>) Cattle Tick (<i>Rhipicephalus (Boophilus) microplus</i>)</p>
---------	--

Maximum tick efficacy is achieved 4-5 days after treatment. Engorged female ticks that drop from cattle in the initial days after treatment may lay viable eggs. This should be taken into account when planning a strategic tick control program. Ivomec LV Pour-On should be alternated with conventional dips or pour-ons when the program requires repeat treatments.

#### Sustained Activity

At the recommended dose rate, Ivomec LV Pour-On effectively controls infection with *Ostertagia* spp. (Small Brown Stomach Worm) acquired up to 14 days after treatment, with *Dictyocaulus viviparus* (Lungworm) acquired up to 28 days after treatment and *Haematobia irritans exigua* (Buffalo Fly) up to 21 days after treatment.

Resistance may develop to any chemical.

Net Contents: 250 mL, 1 L, 2.5 L, 3 L, 5 L, 5.5 L, 15 L, 20 L.

Directions for Use:

Restrains: Re-treatment interval: DO NOT re-treat with Ivomec LV Pour-On within the meat withholding period of 21 days.

Contraindications:

Precautions: Use with caution as rainfall on cattle less than 2 hours after dosing may result in reduced efficacy.

Side Effects: Temporary hair loss occasionally occurs along the application site. This resolves without treatment.

#### Dosage and Administration:

Ivomec LV Pour-On dose rate is 1 mL/20 kg body weight. Apply along the backline of the animal.

Live weight (kg)	Dose Volume (mL)	250 mL pack treats	1 Litre pack treats	2.5 Litre pack treats	3 Litre pack treats	5 Litre pack treats	5.5 Litre pack treats	15 Litre pack treats	20 Litre pack treats
51-100	5	50	200	500	600	1000	1100	3000	4000
101-150	7.5	33	133	333	400	666	733	2000	2666
151-200	10	25	100	250	300	500	550	1500	2000
201-250	12.5	20	80	200	240	400	440	1200	1600
251-300	15	16	66	166	200	333	366	1000	1333
301-350	17.5	14	57	142	171	285	314	857	1142
351-400	20	12	50	125	150	250	275	750	1000
401-450	22.5	11	44	111	133	222	244	666	888
451-500	25	10	40	100	120	200	220	600	800
501-550	27.5	9	36	90	109	181	200	545	727
551-600	30	8	33	83	100	166	183	500	666
601-650	32.5	7	30	76	92	153	169	461	615

Over 650 kg dose at rate of 1 mL/20 kg body weight.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose according to the heaviest animal by liveweight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

General Directions: NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION

Withholding Periods:	MEAT: DO NOT USE less than 21 days before slaughter for human consumption. MILK: Zero (0) days.
Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 21 days before slaughter for export. Before using this product confirm the current ESI from Boehringer Ingelheim Animal Health on 1800 808 691 or the APVMA website ( <a href="http://www.apvma.gov.au/residues">www.apvma.gov.au/residues</a> ).
Safety Directions:	Poisonous if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.
First Aid Warnings:	
Additional User Safety:	Additional information is in the Safety Data Sheet.
Environmental Statements:	<b>ENVIRONMENTAL PROTECTION</b> Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.
Disposal:	[250 mL & 1 L containers:] Dispose of container by wrapping with paper and putting in garbage.  [Containers larger than 1 L:] Triple-rinse container and dispose of rinsate in compliance with relevant local, state or territory government regulations. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. The cap should not be replaced, but may be recycled separately with the container. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.
Storage:	Store below 30 °C (room temperature).