

LIME SULPHUR
FUNGICIDE/INSECTICIDE

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Kendon
LIME SULPHUR
FUNGICIDE/INSECTICIDE

ACTIVE CONSTITUENT:
200g/L Sulphur (S) as Polysulphide Sulphur

For the control of certain diseases, insects and mites on pome fruit, stone fruit, citrus, grapes, roses, hedges, vegetables as specified in the Directions for Use.

CONTENTS: 5L, 20L, 200L

Kendon Chemical & MNFG. CO. PTY LTD 71 McClure Street, Thornbury, VIC 3071

AN AUSTRALIAN OWNED COMPANY EST. 1950

N.R.A.	
APPROVED LABEL (TEXT)	
Label No:	53131/1201
Date:	4/1/02
Signature:	Atall

LIME SULPHUR
FUNGICIDE/INSECTICIDE

DIRECTIONS FOR USE:

CROP	PEST/DISEASE	STATE	RATE/100L WATER	CRITICAL COMMENTS
Citrus Fruit Oranges Lemons Grapefruit Mandarins, etc	White Louse Scale	NSW,ACT, Vic, SA, Tas Only	10.0L	Winter Spray: Apply in July and thoroughly wet all parts of the tree.
		Qld Only	3.0 – 6.5L	
	Maori Mite	NSW,ACT, Vic, SA, WA, Tas only	2.5L	Summer Spray: Spray to thoroughly wet leaves
Grapes	Bud mite Erinose mite Rust mite Vine bunch mite	Vic, SA, WA, Tas only	10.0L	Apply as near as possible to budburst. Note: Spread of mites is checked by sulphur treatments used for powdery mildew.
		Qld, NSW, ACT only	5.0L	
	Post harvest treatment for mites	SA only	2.0L	As soon as possible after harvest.
Pome Fruit Apples	Blackspot (Scab), Powdery mildew	All States	5.0L	Apply during dormant to late budswell. Note: Lime Sulphur used after late green tip will restrict growth and depress yield. DO NOT use on Delicious or Cox's Orange Pippin apples.
	San Jose Scale	Qld, NSW, ACT, Vic, SA, WA only		
Pears	Blackspot Byrobia mite Pear leaf blister mite	All States	5.0L	Apply during dormant to late budswell. Note: Lime Sulphur used after late green tip will restrict growth and depress yield. DO NOT use on Delicious or Cox's Orange Pippin apples.
Stone Fruit Peaches Nectarines Prunes Plums Apricots Almonds	Byrobia mite Shot hole Leaf curl Freckle Rust	All States	5.0L	Spray while tree is dormant to budswell stage.
	Brown rot (Blossom blight) San Jose Scale	Qld, NSW, ACT, Vic, SA Tas only		
	Frosted scale	Qld, Vic, SA, WA, Tas only		
Summer Spray Roses Hedges Vegetables Tomatoes	Two spotted mite Tomato mite Powdery mildew Rust	All States	1.0L	Spray to wet leaves and repeat as necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS: DO NOT apply during the growing period of Apricots, Raspberries, Cucurbits, Peaches or other sulphur sensitive plants.

Mixing: Shake contents well before using. Add the required amount of product to the water in the spray tank with stirring or agitation. Agitate while spraying. DO NOT mix the product with acidic water or pH below 4.

Application: Spray in the cool of the day. The spray mixture should be used on the same day it is mixed with water. DO NOT apply when freezing weather is predicted or the air temperature is over 32 degrees celsius. DO NOT apply product within 2 weeks of an oil spray except as indicated under the "Compatibility" statement below.

COMPATIBILITY: This product may be mixed with Kendon White Oil for use on Pome and Stone fruit during the dormant to semi-dormant stages only. For details of this mixture consult the label for Kendon White Oil.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT: DO NOT
contaminate streams, rivers or waterways with the chemical or used containers.

STORAGE AND DISPOSAL: Store in the closed, original container in a well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS: Avoid contact with skin and eyes. When preparing the spray and using the prepared spray, wear elbow length PVC gloves and face shield or goggles. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each days use, wash gloves and face shield or goggles and contaminated clothing.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre (131126).

MATERIAL SAFETY DATA SHEET

Additional information is listed in Kendon Material Safety Data Sheet No A565

Batch No:

Date of Manufacture

NRA Approval No.:

53131/1201

KENDON
LIME SULPHUR FUNGICIDE / INSECTICIDE

ACTIVE CONSTITUENT:
200g/L SULPHUR(S) AS POLYSULPHIDE SULPHUR

For the control of certain diseases, insects and mites
on apples, pears, stone fruit, grapes,
roses, hedges and vegetables in the home garden

250mL / 500mL

Kendon Chemical & MNFG. CO. PTY. LTD.
71 McClure St., Thornbury, VIC 3071
(03)94972822

AN AUSTRALIAN OWNED COMPANY EST 1950

HOW TO USE

DO NOT apply when freezing weather is predicted

DO NOT apply when the air temperature is over 32 °C

DO NOT apply within 2 weeks of an oil spray

PLANT	DISEASE or INSECT	RATE	HOW TO APPLY
Apples	Black Spot and Powdery Mildew	50ml per litre of water	During dormant to late bud swell period DO NOT use on Delicious or Cox's Orange Pippin Apples
	San Jose Scale QLD, NSW, ACT, VIC, SA, WA only		
Pears	Black Spot, Bryobia mite, Pear leaf blister mite		
Apricots Almonds Nectarines Peaches Plums Prunes	Leaf Curl, Shot Hole, Freckle, Rust	50ml per litre of water	Spray whilst tree is dormant to bud swell stage
	Brown Rot (Blossom Blight) San Jose Scale QLD, NSW, ACT, VIC, SA, TAS only		
	Frosted scale QLD, VIC, SA, WA, TAS only		
Citrus	White Louse Scale NSW, ACT, Vic SA Tas only	100ml per litre of water	Winter spray. Apply in July and thoroughly wet all parts of the tree.
	White Louse-Scale Qld only	30 –65ml per litre of water	
	Maori Mite NSW, ACT, Vic, SA, WA Tas only	25ml per litre of water	Summer spray – spray to thoroughly wet leaves
Grapes	Mites VIC, SA, WA, TAS only	100ml per litre of water	Apply as near as possible to budburst
Roses, Hedges	Two Spotted Mite, Powdery Mildew, Rust	10ml per litre of water	Apply in Spring, Summer & Autumn. Spray to wet leaves and repeat as necessary.
Tomatoes, Vegetables	Tomato mite, Two Spotted mite, Powdery Mildew, Rust		

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED

HOW TO PREPARE

Shake contents well before using

Add the required amount of product to the water and mix

Use only when freshly mixed – do not store prepared spray

APPLICATION

Spray in the cool of the day

CAUTION

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

STORAGE and DISPOSAL

Store in the original container in a cool, dry place out of the reach of children. Do not store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (131126). .

B/N:

Date of Manufacture

NRA Approval No: 53131/1201.

: