

SF00-SF06-0002

# HERCULES

Pour-On for Lice



## Active Content & Action

Imidacloprid 35 g/L. Imidacloprid is part of the neonicotinoid class of chemicals. Neonicotinoids disrupts nerve transmission in target parasite.

## Use

A ready to use pour on to control neonicotinoid susceptible body lice (*Bovicola ovis*) on sheep.

## Indications

Can be applied to shorn sheep up to 7 days off shears and on unshorn lambs up to 2 months of age.

## Period of Protection

The product can be expected to protect sheep against re-infestation with lice for 4 weeks after treatment when applied within 24 hours of shearing.

## Dosage and Administration FOR EXTERNAL USE ONLY.

Apply the entire dose to sheep under 30 kg as a single broadband stripe from poll to tail butt. For sheep over 30 kg, apply half the dose as a broadband stripe either side of the midline from the mid neck to the butt of the tail. The stripes should be no further than 10cm apart. Refer to the diagrams (right).

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group (e.g. lambs), dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

Add 10mL for each 10kg above 100kg.

## Withholding Periods

Meat : 21 Days  
Milk : DO NOT USE on female sheep which are producing or may in the future produce milk or milk products for human consumption  
Wool : 6 months

## Trade Advice

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 63 days before slaughter for export. Before using this product, confirm the current ESI from ZAGRO AUSTRALIA Pty Ltd on 1800 144 884 or the APVMA website ([www.apvma.gov.au/residues](http://www.apvma.gov.au/residues))

## Accessories

This product can be used with any pour on applicator designed to apply pour on products to sheep. Contact your local reseller for the different options.

Bodyweight (kg)	Dose (mL)	Method	Apply in the pattern below
Below 6kg	8	Single stripe	
6.0 - 8.0	12		
8.1 - 10.0	15		
10.1 - 12.5	20		
12.6 - 15	25		
15.1 - 20	30		
20.1 - 30	40	Double stripe	
30.1 - 55	60		
55.1 - 80	80		
80.1 - 100	90		

## Doses

333 doses per 20L pack (average weight 50kg).

Pack Size(s) Available  
20L

Australian Poisons Schedule  
S5

## Presentation

40 drums to a pallet

APVMA Approval No.:  
88745

