

Product Name: Farmalinx Shredda 100 EC Herbicide  
APVMA Approval No: 85520/140939



Label Name:	ProForce Fast Draw 100 EC Herbicide
-------------	-------------------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	ACTIVE CONSTITUENT: 100 g/L BUTAFENACIL  SOLVENTS: 660.6 g/L LIQUID HYDROCARBONS 98.6 g/L N-METHYL-2-PYRROLIDONE
-------------------------	---

Mode of Action:	GROUP 14 HERBICIDE
-----------------	--------------------

Statement of Claims:	For the improvement of the control of various broadleaved weeds in commercial, industrial and public service areas and agricultural buildings and yards in a tank mixture with knockdown herbicides as specified in the Directions for Use table.
----------------------	---

Net Contents:	1-1000 L
---------------	----------

Restraints:	DO NOT apply by boomspray. DO NOT apply by aircraft. DO NOT apply if heavy rains or storms are forecast within 48 hours. DO NOT irrigate to the point of runoff for at least 48 hours after application. DO NOT make more than one application per season.  SPRAY DRIFT RESTRAINTS Specific definitions for terms used in this section of the label can be found at <a href="http://www.apvma.gov.au/spraydrift">www.apvma.gov.au/spraydrift</a> DO NOT allow bystanders to come into contact with the spray cloud. DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever
-------------	--

possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.  
DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.  
DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

Directions for Use: This section contains file attachment.

Other Limitations:

Withholding Periods: WITHHOLDING PERIODS:  
For use with Glyphosate: DO NOT GRAZE OR CUT FOR STOCK FEED FOR 6 WEEKS AFTER APPLICATION.  
  
REFER TO THE WITHHOLDING PERIOD OF PRODUCTS MIXED WITH FAST DRAW 100 EC.

Trade Advice:

General Instructions: GENERAL INSTRUCTIONS  
FAST DRAW 100 EC is an emulsifiable concentrate and is to be mixed with water and applied as per the labelled Directions for Use. FAST DRAW 100 EC is a post-emergence herbicide to be added to knockdown herbicides to improve the control of certain broadleaf weeds including marshmallow in commercial, industrial and public service areas, around agricultural buildings, yards and other farm situations.

FAST DRAW 100 EC is a fast acting contact herbicide and aids in control of weeds through a process of membrane disruption. The foliar uptake of FAST DRAW 100 EC is rapid and plant desiccation can occur within 4 days of application. Application of FAST DRAW 100 EC should target small actively growing weeds. Subsequent germinations will not be controlled.

FAST DRAW 100 EC is rapidly absorbed through the foliage of plants. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation, and in subsequent days necrosis and death of the plant.

Mixing  
FAST DRAW 100 EC is an emulsifiable concentrate herbicide which mixes readily with water. Add half the required volume of water to spray tank and start agitation. Add any partner WG herbicide next if it should be added before an EC or knockdown herbicide. Add the measured amount of FAST DRAW 100 EC next, followed by the knockdown herbicide (if not previously added). Add balance of water to tank and add any adjuvant or additive if recommended for use with the knockdown herbicide. Maintain good agitation at all times until spraying is completed. DO NOT store spray mixture overnight.

Application  
Spray equipment should be carefully calibrated before use. Complete coverage of weed seedling is required for maximum knockdown effect. A spray volume of 50 to 100 L/ha is recommended.

	<p><b>Cleaning Equipment After Application</b> Thoroughly clean the sprayer using the following procedure when you have finished spraying highly active materials such as butafenacil products. In addition to the following procedure, ensure proper equipment clean-out for any other products mixed with FAST DRAW 100 EC as specified on the other product labels.</p> <p><b>IMPORTANT:</b> More complete cleaning can be achieved if the spray equipment is cleaned immediately following each use. Mix only as much herbicide spray solution as needed at the time. <b>DO NOT</b> store the sprayer for any extended period of time, especially overnight, with FAST DRAW 100 EC spray solution remaining in the tank, spray lines, spray boom plumbing, spray nozzles or strainers.</p> <p><b>Preparation of the Cleaning Solution:</b> Prepare a spray equipment cleaning solution by mixing alkaline detergent eg "OMO" or "SPREE", "SURF" or Absolute Boomer at a rate of 100 g for every 100 L of clean water used.</p>
--	--

Resistance Warning:	<p><b>Resistant Weeds Warning</b> <b>GROUP 14 HERBICIDE</b> FAST DRAW 100 EC is a member of the pyrimidindiones groups of herbicides. The product has the inhibitors of protoporphyrinogen oxidase mode of action. For weed resistance management, the product is a Group 14 herbicide. Some naturally occurring weed biotypes resistant to the product and other Group 14 herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by this product or other Group 14 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, Indigo Specialty Products Pty Ltd accepts no liability for any losses that may result from the failure of this product to control resistant weeds. Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Indigo Specialty Products representative.</p>
---------------------	---

Precautions:	<p><b>RE-ENTRY</b> <b>DO NOT</b> allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.</p> <p>The user should also observe any re-entry statements listed on other products used in the tank mix.</p>
--------------	--

Protections:	<p><b>PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS</b> <b>DO NOT</b> apply or drain or flush equipment on or near by trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots. <b>DO NOT</b> apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants, crops, cropping lands or pastures.</p> <p><b>PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT</b> Very toxic to aquatic life. <b>DO NOT</b> contaminate wetlands or watercourses with this product or used containers.</p>
--------------	---

Storage and Disposal:	<p>Store in the closed, original container in a dry, well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Triple rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site. If recycling,</p>
-----------------------	--

replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulation. Do not burn empty containers or product.

**Safety Directions:**

Will irritate the eyes and skin. May irritate the nose and throat. Avoid contact with eyes and skin. Do not inhale vapour. When opening the container and preparing spray wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and half face-piece respirator. If product on skin, immediately wash area with soap and water. Wash hands after use. After each day's use, wash gloves, goggles, respirator and if rubber wash with detergent and water, and contaminated clothing.

**First Aid Instructions:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766. If swallowed do not induce vomiting. If in eyes, wash out immediately with water.

**First Aid Warnings:**

## DIRECTIONS FOR USE

CROP	WEEDS CONTROLLED	RATE/ha	CRITICAL COMMENTS
<p>To assist in weed control in Commercial, Industrial and public service areas, around Agricultural buildings and yards.</p>	<p>Annual Ryegrass (<i>Lolium rigidum</i>), Barley Grass (<i>Hordium leporium</i>), Brome Grass (<i>Bromus diandrus</i>), Canola (<i>Brassica napus</i>), Capeweed (<i>Arctotheca calendula</i>), Common Storksbill (max. 4 leaves) (<i>Erodium cicutarium</i>), Deadnettle (<i>Lamium amplexicaule</i>), Dock (seedling) (<i>Rumex crispus</i>), Doublegee/Three Cornered Jack/Spiny Emex (<i>Emex australis</i>), Fleabane (<i>Conyza</i> sp.) suppression, Goosefoot (<i>Chenopodium pumilio</i>), Lupins (<i>Lupinus</i> spp.), Marshmallow (<i>Malva parviflora</i>), Medic (<i>Medicago</i> spp.), Paterson's Curse (<i>Echium plantagenium</i>), Prickly paddy melon (<i>Cucumis myriocarpus</i>), Sorrel (<i>Rumex acetosella</i>), Sowthistle (<i>Sonchus oleraceus</i>), Stink Grass (<i>Eragrostis cilianensis</i>), Sub Clover (<i>Trifolium</i> spp.), Turnip Weed (<i>Rapistrum rugosum</i>), Wild Oats (<i>Avena fatua</i>), Volunteer Cereals, Volunteer canola (conventional variety) (<i>Brassica napus</i>) Volunteer narrow-leafed lupin (<i>Lupinus angustifolius</i>) Wild Radish (<i>Raphanus raphanistrum</i>), Wireweed (<i>Polygonum heterophyllum</i>) Wild Turnip (<i>Brassica tournfortii</i>).</p> <p><b>Refer also to the product label for the knockdown herbicide used. If one of the above weeds is the dominant weed, and there is no specific rate for it in the knockdown herbicide's label, consult the label's generic annual weed rate-range. Select from within this range to suit the weed-stage, weed-density, conditions (etc) of your situation</b></p>	<p><b>100 mL plus a registered rate of glyphosate</b></p>	<p>Treat only actively growing weeds not under stress from low moisture, frost, cold, disease or waterlogging. If heavy grazing occurred allow regrowth before spraying.</p> <p>Addition of FAST DRAW 100 EC to glyphosate will increase the speed at which treated broadleaved weeds in general develop visible symptoms (compared to results achieved with knockdown herbicides applied alone) and may improve final control of broadleaved weeds including certain hard-to-kill weeds, marshmallow in particular. Where broadleaved weeds are a problem such as heliotrope and paddy melon the addition of Rebel (440 g/L 2,4-D Acid) or other formulations of 2,4-D may assist control. Apply only as a tank mix with recommended rates of knockdown herbicides. Refer to the appropriate label for weed sizes and follow all label directions. Addition of Hasten at 0.5% may be beneficial when applying FAST DRAW 100 EC with a glyphosate herbicide.</p> <p>The addition of FAST DRAW 100 EC to glyphosate significantly increases efficacy on fleabane</p>

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**