

Product Name: PRON8URE PASTE
APVMA Approval No: 58902/131950



Label Name:	PRON8URE PASTE
-------------	----------------

Signal Headings:	FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	Each gram contains 180 million CFU as: Lactobacillus acidophilus Lactobacillus delbrueckii subspecies bulgaricus Lactobacillus plantarum Lactobacillus rhamnosus Bifidobacterium bifidum Enterococcus faecium Streptococcus salivarius subspecies thermophilus.
-------------------------	--

Claims:	Concentrated Multi-Strain Probiotic for Animals
---------	---

Net Contents:	30 g 100 g 1 kg 4 kg
---------------	-------------------------------

Directions for Use:	
---------------------	--

Restrains:	PRON8URE® PASTE should not be used in conjunction with Penicillins, Cephalosporins, Tetracyclines, Avoparcin, Gentamycin, Lincomycin, Streptomycin, Tiamulin or Tylosin, but may be used immediately following these antibiotics.
------------	---

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	
---------------	--

Dosage and Administration:	This section contains file attachment.
General Directions:	
Withholding Periods:	Meat: Zero (0) days Milk: Zero (0) days
Trade Advice:	Export Slaughter Interval (ESI): Zero (0) days. Before using this product, confirm the current ESI from INTERNATIONAL ANIMAL HEALTH PRODUCTS PTY LTD on (02) 9672 7944 or the APVMA website (www.apvma.gov.au/residues).
Safety Directions:	Avoid contact with eyes.
First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Australia 131 126.
First Aid Warnings:	
Additional User Safety:	
Environmental Statements:	
Disposal:	<p>For 30 g, 100 g and 1 kg pack sizes: Dispose of syringe by wrapping with paper and putting in garbage.</p> <p>For 4 kg pack size: Triple-rinse container into the medicated water, dip, drench, etc. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p>
Storage:	<p>STORE BELOW 25°C (Air Conditioning). KEEP OUT OF REACH OF CHILDREN. If this product is stored under refrigerated conditions (less than 8°C), the stated expiry may be extended by 12 months.</p> <p>For 1 and 4 kg packs, KEEP CONTAINER TIGHTLY CLOSED.</p>

Dosage and Administration:

- For 1 kg and 4 kg pack size, mix the product well before withdrawing for each use
- Select a clear marked syringe and draw the desired dose.
- Insert the syringe into the mouth and ensure paste is administered on the rear of the tongue.
- Hold animal's head raised for a few seconds after dosing to ensure the paste is swallowed.
- Clean syringe after each dose use.

Ensure that the lid is placed firmly on the container after every use.

Use at first signs of stress*. Dose once a day for a minimum 5 days as listed in the table:

SPECIES	DOSAGE	SPECIES	DOSAGE	SPECIES	DOSAGE
Horses	10-20 g	Cattle, Sheep, Goats, Pigs	2-4 g	Dogs/Cats	1-3 g to 2-6 g
Foals/Ponies	5-10 g	Calves, Lambs, Kids, Piglets	1-2 g	Pups/Kittens	1-2 g

* (associated with pregnancy, birth, weaning, and transportation, loss of appetite, scouring and temperature extremes, medication, surgery, training and exercise).